arXiv:1201.2648v1 [cond-mat.str-el] 12 Jan 2012

Symmetry-protected topological orders for interacting fermions – fermionic topological non-linear σ-models and a group super-cohomology theory

Zheng-Cheng Gu[1] and Xiao-Gang Wen[2, 3]

[1]*Kavli Institute for Theoretical Physics, University of California, Santa Barbara, CA 93106, USA*
[2]*Department of Physics, Massachusetts Institute of Technology, Cambridge, Massachusetts 02139, USA*
[3]*Institute for Advanced Study, Tsinghua University, Beijing, 100084, P. R. China*

Symmetry-protected topological (SPT) phases are gapped quantum phases with a symmetry, which can be smoothly connected to the trivial product states only if we break the symmetry. For a given symmetry, we can have many different SPT phases. But how to describe/construct those different SPT phases that can not be distinguished by their symmetry? It has been shown that different bosonic SPT phases in any dimensions and for any symmetry groups can be described/constructed using group cohomology theory of the symmetry group. In this paper, we introduce a group super-cohomology theory which is a generalization of the standard group cohomology theory. Using the group super-cohomology theory, we can describe/construct different interacting fermionic SPT phases, in any dimensions and for symmetry groups where the fermions form 1D representations of the symmetry group. Just like the boson case, our systematic construction is based on constructing discrete fermionic topological non-linear σ-models from the group super-cohomology theory. Our discrete fermionic topological non-linear σ-model, when defined on a space-time with boundary, can be viewed as a "non-local" boundary effective Lagrangian, which is a fermionic and discrete generalization of the bosonic continuous Wess-Zumino-Witten term. Thus we believe that the boundary excitations of a non-trivial SPT phase are gapless if the symmetry is not broken. As an application of this general result, we construct three non-trivial SPT phases in 3D, for interacting fermionic superconductors with coplanar spin order (which have $T^2 = 1$ time-reversal Z_2^T and fermion-number-parity Z_2^f symmetries described by a full symmetry group $Z_2^T \times Z_2^f$). We also construct three interacting fermionic SPT phases in 2D with a full symmetry group $Z_2 \times Z_2^f$. Those 2D fermionic SPT phases all have central-charge $c = 1$ gapless edge excitations, if the symmetry is not broken.

PACS numbers: 71.27.+a, 02.40.Re

Contents

I. Introduction 2

II. Physical pictures of some generic results 4

III. Grassmann tensor network 6
- A. Discretized space-time 6
- B. Physical variables on discretized space-time 6
- C. A constraint on physical variables 7
- D. Grassmann tensor on a single d-dimensional simplex 7
- E. Evaluation of the Grassmann tensors on a complex 8
- F. An example for evaluation of Grassmann tensor network 9

IV. Grassmann tensor network and fermionic path integral 10

V. Topologically invariant Grassmann tensor network and fermionic topological non-linear σ-model 10
- A. (1+1)D case 10
- B. (2+1)D case 11
- C. (3+1)D case 12
- D. The fixed-point action on a closed complex and the unitary condition 12
- E. Symmetry and stability of the fixed-point action amplitude 13

VI. Ideal ground state wave function 14
- A. Construction of 2D wave function 15
- B. No intrinsic topological orders 16

VII. Examples 17
- A. 2D fermionic SPT state with Z_2 symmetry 17
- B. 3D fermionic SPT state with $T^2 = 1$ time-reversal symmetry 19

VIII. Summary 19

IX. acknowledgements 19

A. Topological invariance of the partition amplitudes 20
- 1. $1 + 1$D 20
- 2. $2 + 1$D 21
- 3. $3 + 1$D 24

B. The fixed-point action on a closed complex and the unitary condition 26

C. Group super-cohomology 27
- 1. Graded structure of a group G_b 27
- 2. G-module 28
- 3. Fermionic d-cochain 28
- 4. Fermionic d-cocycle 28

5. An "linearized" representation of group super-cohomology 29
6. Fermionic d-cohomology class 29
7. Group super-cohomology classes and fermionic SPT phases 31
8. A simplified "linear" equations for group super-cohomology 31

D. The mapping $n_d \to f_{d+2}$ induces a mapping $\mathcal{H}^d(G_b, \mathbb{Z}_2) \to \mathcal{H}^{d+2}(G_b, \mathbb{Z}_2)$ 32

E. Calculate $\mathcal{H}^d(G_b, \mathbb{Z}_2)$ – the graded structure 33
1. Calculate $\mathcal{H}^d(Z_n, \mathbb{Z}_2)$ 33
2. Calculate $\mathcal{H}^d(G, \mathbb{Z}_2)$ from $\mathcal{H}^d(G, \mathbb{Z})$ using Künneth formula for group cohomology 34

F. Calculate the group super-cohomology classes $\mathscr{H}^d[G_f, U_T(1)]$ 35
1. A general discussion 35
2. Calculate $\mathscr{H}^d[Z_1 \times Z_2^f, U(1)]$ 36
3. Calculate $\mathscr{H}^d[Z_{2k+1} \times Z_2^f, U(1)]$ 36
4. Calculate $\mathscr{H}^d[Z_2 \times Z_2^f, U(1)]$ 37
5. Calculate $\mathscr{H}^d[Z_2^T \times Z_2^f, U_T(1)]$ 38

G. Properties of the constructed SPT states 38
1. Equivalence between ground state wave functions 38
   a. A special equivalent relation 39
   b. Generic equivalent relations 39
2. Symmetry transformations and generic SPT states 40
3. Entanglement density matrix 41
4. Low energy effective edge Hamiltonian 43

H. Ideal Hamiltonian 44

References 48

I. INTRODUCTION

Landau symmetry breaking theory[1–3] allows us to understand a large class of phases and phase transitions. In 1989, it was realized that many quantum phases can contain a new kind of orders – topological order – which cannot be described by Landau symmetry breaking theory.[4–6] Recently, it was found that topological orders are related to long range entanglements.[7,8] In fact, we can regard topological orders as patterns of long range entanglements[9] defined through local unitary (LU) transformations.[10–12]

The notion of topological orders and long range entanglements leads to the following more general and also more detailed picture of phases and phase transitions (see Fig. 1).[9] For gapped quantum systems without any symmetry, their quantum phases can be divided into two classes: short range entangled (SRE) states and long range entangled (LRE) states.

SRE states are states that can be transformed into direct product states via LU transformations. All SRE states can be transformed into each other via LU transformations. So all SRE states belong to the same phase (see Fig. 1a), *ie* all SRE states can continuously deform into each other without closing energy gap and without phase transition.

LRE states are states that cannot be transformed into direct product states via LU transformations. It turns out that, in general, LRE states may not be connected to each other through LU transformations. The LRE states that are not connected via LU transformations belong to different classes and represent different quantum phases. Those different quantum phases are nothing but the topologically ordered phases.

Fractional quantum Hall states[13,14], chiral spin liquids,[15,16] Z_2 spin liquids,[17–19] non-Abelian fractional quantum Hall states,[20–23] *etc* are examples of topologically ordered phases. The mathematical foundation of topological orders is closely related to tensor category theory[9,10,24,25] and simple current algebra.[20,26] Using this point of view, we have developed a systematic and quantitative theory for non-chiral topological orders in 2D interacting boson and fermion systems.[9,10,25] Also for chiral 2D topological orders with only Abelian statistics, we find that we can use integer K-matrices to describe them.[27–30]

For gapped quantum systems with symmetry, the structure of phase diagram is even richer (see Fig. 1b). Even SRE states now can belong to different phases. The Landau symmetry breaking states belong to this class of phases. However, there are more interesting examples in this class. Even SRE states that have the same symmetry can belong to different phases. The 1D Haldane phase for spin-1 chain[31–34] and topological insulators[35–40] are non-trivial examples of phases with short range entanglements that do not break any symmetry. Those phases are



FIG. 1: (Color online) (a) The possible gapped phases for a class of Hamiltonians $H(g_1, g_2)$ without any symmetry restriction. (b) The possible gapped phases for the class of Hamiltonians $H_{\text{symm}}(g_1, g_2)$ with a symmetry. The shaded regions in (a) and (b) represent the phases with long range entanglement. Each phase is labeled by its entanglement properties and symmetry breaking properties. SRE stands for short range entanglement, LRE for long range entanglement, SB for symmetry breaking, SY for no symmetry breaking. SB-SRE phases are the Landau symmetry breaking phases, which are understood by introducing group theory. The SY-SRE phases are the SPT phases, which can be understood by introducing group cohomology theory.

Symm. G	$d_{sp}=0$	$d_{sp}=1$	$d_{sp}=2$	$d_{sp}=3$
$U(1)\rtimes Z_2^T$	$\mathbb{Z}$	$\mathbb{Z}_2$	$\mathbb{Z}_2$	$\mathbb{Z}_2^2$
$U(1)\times Z_2^T$	$\mathbb{Z}_1$	$\mathbb{Z}_2^2$	$\mathbb{Z}_1$	$\mathbb{Z}_2^3$
$U(1)$	$\mathbb{Z}$	$\mathbb{Z}_1$	$\mathbb{Z}$	$\mathbb{Z}_1$
$SO(3)$	$\mathbb{Z}_1$	$\mathbb{Z}_2$	$\mathbb{Z}$	$\mathbb{Z}_1$
$SO(3)\times Z_2^T$	$\mathbb{Z}_1$	$\mathbb{Z}_2^2$	$\mathbb{Z}_2$	$\mathbb{Z}_2^3$
Z_2^T	$\mathbb{Z}_1$	$\mathbb{Z}_2$	$\mathbb{Z}_1$	$\mathbb{Z}_2$
Z_n	$\mathbb{Z}_n$	$\mathbb{Z}_1$	$\mathbb{Z}_n$	$\mathbb{Z}_1$
$Z_2\times Z_2$	$\mathbb{Z}_2^2$	$\mathbb{Z}_2$	$\mathbb{Z}_2^3$	$\mathbb{Z}_2^2$
$Z_n\times Z_2^T$	$\mathbb{Z}_{(2,n)}$	$\mathbb{Z}_2\times\mathbb{Z}_{(2,n)}$	$\mathbb{Z}_{(2,n)}^2$	$\mathbb{Z}_2\times\mathbb{Z}_{(2,n)}^2$

TABLE I: (Color online) Bosonic SPT phases in d_{sp}-spatial dimensions protected by some simple symmetries (represented by symmetry groups G). Here $\mathbb{Z}_1$ means that our construction only gives rise to the trivial phase. $\mathbb{Z}_n$ means that the constructed non-trivial SPT phases plus the trivial phase are labeled by the elements in $\mathbb{Z}_n$. Z_2^T represents time-reversal symmetry, $U(1)$ represents $U(1)$ symmetry, Z_n represents cyclic symmetry, *etc* . Also (m,n) is the greatest common divisor of m and n. The red rows are for bosonic topological insulators and the blue rows bosonic topological superconductors.

beyond Landau symmetry breaking theory since they do not break any symmetry. On the other hand, their very existence requires symmetry in Hamiltonian. We will call this kind of phases Symmetry Protected Topological (SPT) phases.[33,34] Some examples of 2D SPT phases protected by translation and some other symmetries were discussed in Ref. 41–43.

It turns out that there is no gapped LRE state in 1D.[11] So all 1D gapped states are either symmetry breaking states or SPT states. This realization led to a complete classification of all 1D gapped quantum phases.[42,43,48]

In Ref. 49,50, the classification of 1D SPT phase is generalized to any dimensions: *For gapped bosonic systems in d_{sp}-spatial dimension with an on-site symmetry G, we can construct distinct SPT phases that do not break the symmetry G from distinct elements in $\mathcal{H}^{d+1}[G,U_T(1)]$ – the $(1+d)$-cohomology class of the symmetry group G with G-module $U_T(1)$ as coefficient.* Note that the above result does not require the translation symmetry. The results for some simple on-site symmetry groups are summarized in Table I. In particular, the interacting bosonic topological insulators/superconductors are included in the table.

In Ref. 25, the LU transformations for fermionic systems were introduced, which allow us to define and study gapped quantum phases and topological orders in fermionic systems. In particular, we developed a fermionic tensor category theory to classify non-chiral intrinsic fermionic topological orders defined through the fermionic LU transformations without any symmetry. In this paper, we are going to use the similar line of thinking to study fermionic gapped phases with symmetries. To begin, we will study the simplest kind of them – short range entangled fermionic phases with symmetries. Those phases are called fermionic SPT phases which can be defined through the fermionic LU transformations with symmetry. Fermionic SPT phases are $T=0$ gapped *symmetry unbroken* phases of fermionic Hamiltonians with a certain symmetry. But if we break the symmetry of the Hamiltonians, all those symmetry unbroken phases can be smoothly connected to the trivial product states. Note that in 0 spatial dimension, the trivial product states can have even or odd numbers of fermions. So the trivial product states in 0D can belong to two different phases. In higher dimensions, all the trivial product states belong to one phase (if there is

Interacting fermionic SPT phases					
$G_f\ \backslash d_{sp}$	0	1	2	3	Example
“none”$=Z_2^f$	$\mathbb{Z}_2$	$\mathbb{Z}_1$	$\mathbb{Z}_1$	$\mathbb{Z}_1$	superconductor
$Z_2\times Z_2^f$	$\mathbb{Z}_2^2$	$\mathbb{Z}_2$	$\mathbb{Z}_2^2$	$\mathbb{Z}_1$	
$Z_2^T\times Z_2^f$	$\mathbb{Z}_2$	$\mathbb{Z}_2^2$	$\mathbb{Z}_1$	$\mathbb{Z}_2^2$	supercond. with coplanar spin order
$Z_{2k+1}\times Z_2^f$	$\mathbb{Z}_{4k+2}$	$\mathbb{Z}_1$	$\mathbb{Z}_{2k+1}$	$\mathbb{Z}_1$	

Interacting fermionic gapped phases					
$G_f\ \backslash d_{sp}$	0	1	2	3	Example
“none”$=Z_2^f$	$\mathbb{Z}_2$	$\mathbb{Z}_2$	?	?	superconductor
$Z_2\times Z_2^f$	$\mathbb{Z}_2^2$	4	?	?	
$Z_2^T\times Z_2^f$	$\mathbb{Z}_2$	8	?	?	supercond. with coplanar spin order
$Z_{2k+1}\times Z_2^f$	$\mathbb{Z}_{4k+2}$	$\mathbb{Z}_2$	?	?	
$Z_{2k}\times Z_2^f$	$\mathbb{Z}_{2k}\times\mathbb{Z}_2$	4	?	?	

Non-interacting fermionic SPT phases					
$G_f\ \backslash d_{sp}$	0	1	2	3	Example
“none”$=Z_2^f$	$\mathbb{Z}_2$	$\mathbb{Z}_1$	$\mathbb{Z}_1$	$\mathbb{Z}_1$	superconductor
$Z_2\times Z_2^f$	$\mathbb{Z}_2^2$	$\mathbb{Z}_2$	$\mathbb{Z}$	$\mathbb{Z}_1$	
$Z_2^T\times Z_2^f$	$\mathbb{Z}_2$	$2\mathbb{Z}$	$\mathbb{Z}_1$	$\mathbb{Z}_1$	supercond. with coplanar spin order

Non-interacting fermionic gapped phases					
$G_f\ \backslash d_{sp}$	0	1	2	3	Example
“none”$=Z_2^f$	$\mathbb{Z}_2$	$\mathbb{Z}_2$	$\mathbb{Z}$	$\mathbb{Z}_1$	superconductor
$Z_2\times Z_2^f$	$\mathbb{Z}_2^2$	$\mathbb{Z}_2^2$	$\mathbb{Z}^2$	$\mathbb{Z}_1$	
$Z_2^T\times Z_2^f$	$\mathbb{Z}_2$	$\mathbb{Z}$	$\mathbb{Z}_1$	$\mathbb{Z}_1$	supercond. with coplanar spin order

TABLE II: (Color online) Fermionic SPT phases in d_{sp}-spatial dimensions constructed using group super-cohomoloy for some simple symmetries (represented by the full symmetry groups G_f). The red symmetry groups can be realized by electron systems. Here $\mathbb{Z}_1$ means that our construction only gives rise to the trivial phase. $\mathbb{Z}_n$ means that the constructed non-trivial SPT phases plus the trivial phase are labeled by the elements in $\mathbb{Z}_n$. Note that fermionic SPT phases always include the bosonic SPT phases with the corresponding bosonic symmetry group $G_b\equiv G_f/Z_2^f$ as special cases. The blue entries are complete constructions which become classifications. Z_2^T represents time-reversal symmetry, $U(1)$ represents $U(1)$ symmetry, *etc* . As a comparison, the results for non-interacting fermionic gapped/SPT phases,[44] as well as the interacting ones in 1D,[45–48] are also listed. Note that the symmetric interacting and non-interacting fermionic gapped phases can be the SPT phases or intrinsically topologically ordered phases. This is why the lists for gapped phases and for SPT phases are different. Here a number n means that there are $n-1$ non-trivial phases and one trivial phase. $2\mathbb{Z}$ means that the phases are labeled by even integers. Note that all the phases listed above respect the symmetry G_f.

no symmetry). This is why when there is no symmetry, there are two fermionic SPT phases in 0D and only one trivial fermionic SPT phase in higher dimensions.

After introducing the concept of fermionic SPT phase, we find that we can generalize our construction of bosonic SPT orders to fermion systems, by generalizing the group cohomology theory[50,51] to group super-cohomology theory. This allows us to systematically construct fermionic SPT phases for interacting fermions in any dimensions and for a certain type of symmetries where the fermions form an 1D representation of the symmetry group. The constructed fermionic SPT phases for some simple symmetry groups are given in Table II. The rows correspond to different symmetries for the fermion systems.

We note that, sometimes in literature, when we describe the symmetry of a fermion system, we do not include the fermion-number-parity transformation $P_f = (-)^N$ in the symmetry group. In this paper and in Table II, we use the full symmetry group G_f which includes the fermion-number-parity transformation to describe the symmetry of fermion systems.[44] So the full symmetry group of a fermion system with no symmetry is $G_f = Z_2^f$ generated by P_f. The bosonic symmetry group G_b is given by $G_b = G_f/Z_2^f$, which correspond to the physical symmetry of the fermion system that can be broken. In fact, the full symmetry group is a projective symmetry group discussed in Ref. 52, which is a Z_2^f extension of the physical symmetry group G_b.

The columns correspond to different spatial dimensions. In 0D and 1D, our results reproduce the exact results obtained from previous studies.[45–48] The results for 2D and 3D are new.

Each entry indicates the number of non-trivial phases plus one trivial phase. For example, $\mathbb{Z}_2$ means that there is one non-trivial SPT phase labeled by 1 and one trivial phase labeled by 0. Also, say, 8 means that there are seven non-trivial SPT phases and one trivial phase. For each non-trivial fermionic SPT phase, we can construct the ideal ground state wave function and the ideal Hamiltonian that realizes the SPT phase, using group super-cohomology theory.

We believe that each non-trivial SPT phase has gapless boundary excitations protected by the same symmetry that protects the bulk SPT phase. This belief is supported by the discussion in the next section, and by the fact that the boundary excitations are described by a "non-local" Lagrangian which is the fermionic and discrete generalization of the bosonic continuous Wess-Zumino-Witten (WZW) term.[53,54]

In the rest of this paper, we will first compare the results in Table II with some known results for 1D and for free fermion systems. This will give us some physical understanding of Table II. We then define fermionic path integral on discrete space-time via Grassmann tensor network. Next we discuss the conditions under which the fermionic path integral becomes a fixed-point theory under the coarse-graining transformation of the space-time complex. Such a fixed-point theory is a fermionic topological non-linear σ-model. The fixed-point path integral describes a fermionic SPT phase. We then construct the ground state wave function from the fixed-point path integral. Some simple explicit examples of fermionic SPT phases and their boundary excitations are discussed. In appendix, we develop a group super-cohomology theory, and calculate the Table II from the group super-cohomology theory. We also construct the exactly soluble Hamiltonian of the fermionic SPT phases from the fixed-point path integral.

II. PHYSICAL PICTURES OF SOME GENERIC RESULTS

Before describing how to obtain the generic results in Table II, in this section, we will compare some of the generic results with known results for 1D interacting systems and 2D/3D non-interacting systems. The comparison will give us a physical understanding for some of the interacting fermionic SPT phases.

First, from the Table II, we find that fermion systems with $T^2 = 1$ time-reversal symmetry (or the full symmetry group $G_f = Z_2^T \times Z_2^f$) can have three non-trivial fermionic SPT phases in $d_{sp} = 1$ spatial dimension and three non-trivial fermionic SPT phases in $d_{sp} = 3$ spatial dimensions.

For 0D free or interacting fermion systems with $T^2 = 1$ time-reversal symmetry Z_2^T, the possible symmetric gapped phases are the 1D representations of $Z_2^T \times Z_2^f$. Since the time-reversal transformation T is anti unitary, Z_2^T has only one trivial 1D representation. Z_2^f has two 1D representations. Thus 0D fermion systems with $T^2 = 1$ time-reversal symmetry are classified by $\mathbb{Z}_2$ corresponding to even fermion states and odd fermion states.

For 1D free fermion systems with $T^2 = 1$ time-reversal symmetry Z_2^T, the possible gapped phases are classified by $\mathbb{Z}$.[55,56] For the phase labeled by $n \in \mathbb{Z}$, it has n Majorana zero modes at one end of the chain.[57] Those boundary states form a representation of n Majorana fermion operators $\eta_1, ..., \eta_n$ which all transform the same way under the time-reversal transformation T. (If, say η_1 and η_2 transform differently under T: $\eta_1 \to \eta_1$ and $\eta_2 \to -\eta_2$, $i\eta_1\eta_2$ will be invariant under T and such a term will gap out the η_1 and η_2 modes.) Some of those 1D gapped states have intrinsic topological orders. Only those labeled by even integers (which have even numbers of Majorana boundary zero modes) become the trivial phase after we break the time-reversal symmetry Z_2^T. Thus the free fermion SPT phases are labeled by even integers.

For interacting 1D systems with $T^2 = 1$ time-reversal symmetry, there are eight possible gapped phases that do not break the time-reversal symmetry.[45–48] Four of them have intrinsic topological orders (which break the fermion-number-parity symmetry in the mapped bosonic model)[48] and the other four are fermionic SPT phases given in Table II (three of them are non-trivial fermionic SPT phases and the fourth one is the trivial phase).

For 3D free fermion systems with $T^2 = 1$ time-reversal symmetry, there is no non-trivial gapped phase.[55,56] In contrast, for 3D interacting fermion systems with $T^2 = 1$ time-reversal symmetry, there are (at least) three non-trivial fermionic SPT phases. One of the non-trivial 3D fermionic SPT phases is actually a bosonic SPT phase formed by tightly bounded fermion pairs. The other two fermionic SPT phases cannot be obtained from bosonic models and are totally new. So the non-trivial 3D fermionic SPT phases protected by $T^2 = 1$ time-reversal symmetry cannot be constructed from free fermion systems.

We also find that fermion systems with symmetry group $G_b = Z_2$ (or the full symmetry group $G_f = Z_2 \times Z_2^f$) can have one non-trivial fermionic SPT phases in $d_{sp} = 1$ spatial dimension and three non-trivial fermionic SPT phases in $d_{sp} = 2$ spatial dimensions. (One of the non-trivial 3D fermionic SPT phases is actually a bosonic SPT phase while the other two are intrinsically fermionic.)

To compare the above result with the known free fermion result, let us consider free fermion systems with symmetry group $G_b = Z_2$ (or the full symmetry group $G_f = Z_2 \times Z_2^f$), which contain two kinds of fermions: one carries the Z_2-charge 0 and the other carries Z_2-charge 1. For such free fermion systems, their gapped phases are classified by[44]

$$\begin{array}{rcccccccc} d_{sp}: & 0 & 1 & 2 & 3 & 4 & 5 & 6 & 7 \\ \text{gapped phases}: & \mathbb{Z}_2^2 & \mathbb{Z}_2^2 & \mathbb{Z}^2 & 0 & 0 & 0 & \mathbb{Z}^2 & 0 \end{array} \tag{1}$$

The four $d_{sp} = 0$ phases correspond to the ground state with even or odd Z_2-charge-0 fermions and even or odd Z_2-charge-1 fermions. The four $d_{sp} = 1$ phases correspond to the phases where the Z_2-charge-0 fermions are in the trivial or non-trivial phases of Majorana chain[57] and the Z_2-charge-1 fermions are in the trivial or non-trivial phases of Majorana chain. A $d_{sp} = 2$ phase labeled by two integers $(m, n) \in \mathbb{Z}^2$ corresponds to the phase where the Z_2-charge-0 fermions form m $(p_x + \mathrm{i}p_y)$ states with m right moving Majorana chiral modes and the Z_2-charge-1 fermions form n $(p_x + \mathrm{i}p_y)$ states with n right moving Majorana chiral modes. (if m and/or n are negative, the fermions then form the corresponding number of $(p_x - \mathrm{i}p_y)$ states with the corresponding number of left moving Majorana chiral modes.)

Some of the above non-interacting gapped phases have intrinsic fermionic topological orders. (Those phases are stable and have intrinsic fermionic topological orders even after we turn on interactions.) So only a subset of them are non-interacting fermionic SPT phases:

$$\begin{array}{rcccccccc} d_{sp}: & 0 & 1 & 2 & 3 & 4 & 5 & 6 & 7 \\ \text{SPT phases}: & \mathbb{Z}_2^2 & \mathbb{Z}_2 & \mathbb{Z} & 0 & 0 & 0 & \mathbb{Z} & 0 \end{array} \tag{2}$$

The four $d_{sp} = 0$ phases correspond to the ground states with even or odd numbers of fermions and 0 or 1 Z_2-charges. The two $d_{sp} = 1$ phases correspond to the phases where the Z_2-charge-0 fermions and the Z_2-charge-1 fermions are both in the trivial or non-trivial phases of Majorana chain. The $d_{sp} = 2$ phase labeled by one integer $n \in \mathbb{Z}$ corresponds to the phase where the Z_2-charge-0 fermions have n right moving Majorana chiral modes and the Z_2-charge-1 fermions have n left moving Majorana chiral modes. We see that in 0D and 1D, the non-interacting fermionic SPT phases are the same as the interacting fermionic SPT phases.

However, in two dimensions, the non-interacting fermionic SPT phases are quite different from the interacting fermionic SPT phases. Despite very different phase diagram, it appears that the above three non-trivial interacting fermionic SPT phases in 2D can be realized by free fermion systems. In fact, it appears that there are seven non-trivial interacting 2D fermionic SPT phases protected by Z_2 symmetry[58]. All the seven non-trivial SPT phases can be represented by free fermion SPT phases. This suggests that our current construction is incomplete, since we only obtain four fermionic SPT phases (including the trivial one) with the $G_f = Z_2 \times Z_2^f$ symmetry. One possible reason of the incompleteness may be due to our limiting requirement that fermions only form 1D representations of G_f.

One way to understand why there can only be seven non-trivial interacting 2D fermionic SPT phases protected by Z_2 symmetry is using the idea of duality between intrinsic topological orders and SPT orders discovered recently[59]. The key observation in such a duality map is that for any SPT orders associated with a (discrete) global symmetry G, we can always promote the global symmetry to a local (gauge) symmetry. For different SPT phases, the corresponding promoted (discrete) gauge models describe different intrinsic topological orders. For fermionic SPT phases protected by Z_2 symmetry, we can let Z_2-charge-1 fermion couples to a Z_2 gauge field. In this way, different intrinsic topological ordered phases can be characterized by different statistics of the Z_2-flux. According to Kitaev's classification[60] for different types of vortices in superconductors[69], we know that there are seven non-trivial cases. Thus, we see that although in free 2D fermionic systems, SPT phases protected by Z_2 symmetry is classified by an integer (Chern number), the interactions dramatically changes the classifications.

Following Kitaev's idea[60], we can have a very simple way to understand the seven non-trivial types of vertices by counting the number of Majorana modes in the vertex core. In the corresponding free fermion model, the number of Majoran modes n corresponds to the Chern number of the Z_2-charge-1 free fermion. The seven non-trivial SPT phases are described by $n = 1, 2, \cdots, 7$.

Now, let us try to understand why four of the non-trivial fermionic SPT phases protected by the Z_2 symmetry require fermions to form high dimensional representations of $G_f = Z_2 \times Z_2^f$. It is easy to see that when $n = even$, the free fermion models are described by $n/2$ Z_2-charge-1 complex fermions per site and these fermions form 1D representation of $G_f = Z_2 \times Z_2^f$. However, the situations are more complicated when $n = odd$.



FIG. 2: (Color online) Two branched simplices with opposite orientations. (a) A branched simplex with positive orientation and (b) a branched simplex with negative orientation.

For example, when $n = 1$, the free fermion Hamiltonian describes one Z_2-charge-0 Majorana fermion and one Z_2-charge-1 Majorana fermion per site, labeled as $\gamma_{i,\uparrow}$ and $\gamma_{i,\downarrow}$. Under the Z_2 action, $\gamma_{i,\uparrow}$ does not change while $\gamma_{i,\downarrow}$ changes to $-\gamma_{i,\downarrow}$. Thus the symmetry operation U_{Z_2} can be constructed as $U_{Z_2} = \mathrm{i}^{(N-1)N/2}\prod_{i=1}^{N}\gamma_{i,\uparrow}$, where N is the number of sites. It is easy to check that $U_{Z_2}\gamma_{i,\uparrow}U_{Z_2}^\dagger = \gamma_{i,\uparrow}$, $U_{Z_2}\gamma_{i,\downarrow}U_{Z_2}^\dagger = -\gamma_{i,\downarrow}$ and $U_{Z_2}^2 = 1$. However, the symmetry operator U_{Z_2} can be regarded as the Z_2 symmetry transformation only when $N =$ even, in that case U_{Z_2} contains an even number of fermion operators. But in our construction, the Z_2 symmetry can be realized as bosonic unitary transformation regardless $N =$ even or $N =$ odd. Thus, we can only construct four fermionic SPT phases that are labeled by even n.

Another example requiring fermions to carry high dimensional representations is the well known time reversal symmetry with $T^2 = P_f$. We believe that the principle/framework developed in this paper should be applicable for all these interesting cases and the results will be presented in our future work.

III. GRASSMANN TENSOR NETWORK

After having some physical understanding of the fermionic SPT phases protected by $Z_2^T \times Z_2^f$ or $Z_2 \times Z_2^f$ symmetries, in the following, we will discuss a generic construction that allow us to obtain fermionic SPT phases protected by more general symmetries in any dimensions.

One way to obtain a systematic construction of bosonic SPT phases is through a systematic construction of *topological* bosonic path integral[50] using a tensor network representation of the path integral.[33,61] Here we will use a similar approach to obtain a systematic construction of fermionic SPT phases through *topological* fermionic path integral using a Grassmann tensor network[25,62] representation of the fermionic path integral. The Grassmann tensor network and the associated fermion path integral are defined on a discretized space-time. So first, let us describe the structure of a discretized space-time.

A. Discretized space-time

A d-dimensional discretized space-time is a d-dimensional complex Σ_d formed by many d-dimensional simplices $\Sigma_{[i_0...i_d]}$. We will use $i, j, ...$ to label the vertices of complex Σ_d. $\Sigma_{[i_0...i_d]}$ is the simplex with vertices $i_0, ..., i_d$.

However, in order to define Grassmann tensor network (and the associated fermion path integral) on the discretized space-time Σ_d, the space-time complex must have a so called branching structure.[50,63] A branching structure is a choice of orientation of each edge in the d-dimensional complex so that there is no oriented loop on any triangle (see Fig. 2).

The branching structure induces a *local order* of the vertices on each d-simplex. The first vertex of a d-simplex is the vertex with d outgoing edges, and the second vertex is the vertex with $d-1$ outgoing edges and 1 incoming edge, *etc* . So the simplex in Fig. 2a has the following vertex ordering: $0, 1, 2, 3$.

The branching structure also gives the simplex (and its sub simplices) an orientation. Fig. 2 illustrates two 3-simplices with opposite orientations. The red arrows indicate the orientations of the 2-simplices which are the subsimplices of the 3-simplices. The black arrows on the edges indicate the orientations of the 1-simplices.

B. Physical variables on discretized space-time

To define a path integral on a discretized space-time (*ie* on a complex Σ_d with a branching structure), we associate each vertex i (a 0-simplex) in Σ_d with a variable g_i. So g_i is a local physical dynamical variable of our system. The allow values of g_i form a space which is denoted as G_b. At the moment, we treat G_b as an arbitrary space. Later, we will assume that G_b is the space of a group.

We also associate each $(d-1)$-simplex $(i_0..\hat{i}_i..i_d)$ of a d-simplex $[i_0...i_d]$ with a Grassmann number. Here the $(d-1)$-simplex $(i_0..\hat{i}_i..i_d)$ is a subsimplex of the d-simplex $[i_0...i_d]$ that does not contain the vertex i_i. Relative to the orientation of the d-simplex $[i_0...i_d]$, the $(d-1)$-simplex $(i_0..\hat{i}_i..i_d)$ can have a "+" or "−" orientation (see Fig. 2). We associate $(d-1)$-simplex $(i_0..\hat{i}_i..i_{d-1})$ with a Grassmann number $\theta_{(i_0..\hat{i}_i..i_d)}$ if the $(d-1)$-simplex has a "+" orientation and with a Grassmann number $\bar\theta_{(i_0..\hat{i}_i..i_d)}$ if the $(d-1)$-simplex has a "−" orientation. However, the Grassmann number $\theta_{(i...k)}$ (or $\bar\theta_{(i...k)}$) may or may not present on the $(d-1)$-simplex $(i...k)$. So on each $(d-1)$-simplex $(i...k)$ we also have a local physical dynamical variable $n_{i...k} = 0, 1$ or $\bar n_{i...k} = 0, 1$, indicating whether $\theta_{(i...k)}$ or $\bar\theta_{(i...k)}$ is present or not. So each $(d-1)$-simplex $(i...k)$ is really associated with a dynamical Grassmann variable $\theta_{(i...k)}^{n_{i...k}}$ or $\bar\theta_{(i...k)}^{\bar n_{i...k}}$.

We see that each $(d-1)$-simplex $(i_0...i_{d-1})$ on the surface of the d-complex Σ_d is associated with two types of



FIG. 3: The 2-simplex (012) is associated with $\bar\theta_{(012)}$ which belong to the top 3-simplex [0123]. The 2-simplex (012) is also associated with $\theta_{(012)}$ which belong to the bottom 3-simplex [0124]. A rank 4-tensor $\mathcal{V}_3$ on the top tetrahedron is given by $\mathcal{V}_3^+(g_0, g_1, g_2, g_3)$ and on the lower tetrahedron by $\mathcal{V}_3^-(g_0, g_1, g_2, g_4)$.

Grassmann variables $\theta^{n_{i_0\cdots i_{d-1}}}_{(i_0...i_{d-1})}$ and $\bar\theta^{\bar n_{i_0\cdots i_{d-1}}}_{(i_0...i_{d-1})}$, each belongs to the d-simplex on each side of the $(d-1)$-simplex $(i_0...i_{d-1})$ (see Fig. 3). We also see that each d-simplex has its own set of Grassmann variables that do not overlap with the Grassmann variables of other simplices.

C. A constraint on physical variables

In general, the physical dynamical variables g_i, $n_{i...k}$, and $\bar n_{i...k}$ are independent. We can formulate fermionic topological theory and obtain fermionic topological phases by treating g_i, $n_{i...k}$, and $\bar n_{i...k}$ as independent variables. This is essentially the approach used in Ref. 25 in (2+1)D, and we have obtained a system of non-linear algebraic equations, whose solutions describe various 2D fermionic topological phases. However, the non-linear algebraic equations are very hard to solve, despite they only describe 2D fermionic topological orders. As a result, we have obtained very few fermionic solutions.

In this paper, we are going to study a simpler and less general problem, by putting a constraint between the physical dynamical variables g_i and $n_{i...k}$ $(\bar n_{i...k})$:

$$n_{i...k} = n(g_i, ..., g_k), \qquad \bar n_{i...k} = n(g_i, ..., g_k). \tag{3}$$

That is when $n_{i...k} \neq n(g_i, ..., g_k)$ or $\bar n_{i...k} \neq n(g_i, ..., g_k)$, there will be a huge energy cost. So at low energies, the system always stays in the subspace that satisfy the above constraint. In this case $n_{i...k}$, and $\bar n_{i...k}$ are determined from g_i's.

It turns out that the constraint indeed simplifies the mathematics a lot, which allows us to systematically describe fermionic topological phases in any dimensions. The resulting non-linear algebraic equations also have nice structures which can be solved more easily. So we can also find many solutions in any dimensions and hence obtain many new examples of topological phases in any dimensions. However, the constraint limits us to describe only SPT phases. To describe phases with intrinsic topological orders, we have to use more general unconstraint formulism, which is discussed in Ref. 25. In this paper, we will only consider the simple constraint cases.

The above constraint can be generalized a little bit. In general, we may have several kinds of fermionic, or even bosonic, variables in the $(d-1)$-simplices. So we can associate $\big(\theta^\alpha_{(i_0...i_{d-1})}\big)^{n^\alpha_{i_0\cdots i_{d-1}}}$ or $\big(\bar\theta^\alpha_{(i_0...i_{d-1})}\big)^{\bar n^\alpha_{i_0\cdots i_{d-1}}}$ with each $(d-1)$-simplex. Here $n^\alpha_{i_0...i_{d-1}} = 0, 1$ indicating the presence or the absence of the α^{th} fermionic variable in the $(d-1)$-simplex $(i_0...i_{d-1})$. We can also associate bosonic variables $\big(\phi^a_{(i_0...i_{d-1})}\big)^{l^a_{i_0\cdots i_{d-1}}}$ or $\big(\bar\phi^a_{(i_0...i_{d-1})}\big)^{\bar l^a_{i_0\cdots i_{d-1}}}$ with each $(d-1)$-simplex, where ϕ^a and $\bar\phi^a$ are complex phases. Here $l^a_{i_0...i_{d-1}} = 0, 1, 2, ...$ indicating the number of the a^{th} bosonic variable in the $(d-1)$-simplex $(i_0...i_{d-1})$. Also, $\alpha, a = 1, 2, ...$ label different kinds of fermionic or bosonic variables, which may be in different 1D representations of the symmetry.

In this more general situation, we impose the following constraint between the physical dynamical variables g_i and $n^\alpha_{i...k}$ $(\bar n^\alpha_{i...k})$:

$$\begin{aligned}
& n^{\alpha(g_i,...,g_k)}_{i...k} = n(g_i, ..., g_k), \qquad \bar n^{\alpha(g_i,...,g_k)}_{i...k} = n(g_i, ..., g_k),\\
& \text{other } n^\alpha_{i...k}, \bar n^\alpha_{i...k} = 0,\\
& \alpha(gg_i, ..., gg_k) = \alpha(g_i, ..., g_k).
\end{aligned} \tag{4}$$

So for each fixed set $g_i, ..., g_k$, only one kind of fermionic variable with $\alpha = \alpha(g_i, ..., g_k)$ can appear. We also impose the following constraint between the physical dynamical variables g_i and $l^a_{i...k}(\bar l^a_{i...k})$:

$$\begin{aligned}
& l^a_{i...k} = l^a(g_i, ..., g_k), \qquad \bar l^a_{i...k} = \bar l^a(g_i, ..., g_k),\\
& a(gg_i, ..., gg_k) = a(g_i, ..., g_k).
\end{aligned} \tag{5}$$

So for each fixed set $g_i, ..., g_k$, the number of the bosonic fields for each kind cannot fluctuate. They are given by fixed values $l^a(g_i, ..., g_k)$ and $\bar l^a(g_i, ..., g_k)$. This weaker constraint is enough to simplify the mathematics, since after the constraint, only g_i on vertices can fluctuate independently at low energies. The degrees of freedom on the $(d-1)$-simplices are tied to g_i's and cannot fluctuate independently.

We note that different degrees of freedom are attached to the $(d-1)$-simplices for different choices of $g_i, ..., g_k$. Those degrees of freedom may carry different 1D representations of the symmetry. Later, we will use this picture to construct path integrals that respect the symmetry.

D. Grassmann tensor on a single d-dimensional simplex

In the low energy constraint subspace, we can associate each d-simplex of "+" orientation with a Grassmann rank $(1+d)$-tensor $\mathcal{V}_d^+$, and each d-simplex of "−" orientation with a different Grassmann rank $(1+d)$-tensor $\mathcal{V}_d^-$.

Let us first assume that a d-simplex $[0...d]$ has a "+" orientation, and discuss the structure of the Grassmann tensor $\mathcal{V}_d^+$. The Grassmann tensor $\mathcal{V}_d^+$ is a map from space $G_b^{1+d} \to M_f$:

$$\mathcal{V}_d^+(g_0, g_1, ..., g_d) \in M_f, \quad g_i \in G_b. \tag{6}$$

The order of the variables $g_0, g_1, ..., g_d$ in $\mathcal{V}_d$ is the same as the order of the branching structure: vertex-0 $<$ vertex-1 $<$ ... $<$ vertex-d.

Here an elements in M_f is a complex number times an even number of Grassmann numbers. Thus $\mathcal{V}_d^+(g_0, g_1, ..., g_d)$ has a form

$$\begin{aligned}\mathcal{V}_d^+(g_0, g_1, ..., g_d) &= \nu_d^+(g_0, g_1, g_2, g_3, g_4, ..., g_d)\times \\ &\theta_{(1234...d)}^{n_{d-1}(g_1,g_2,g_3,g_4,...,g_d)} \theta_{(0134...d)}^{n_{d-1}(g_0,g_1,g_3,g_4,...,g_d)} ...\times \\ &\bar\theta_{(0234...d)}^{n_{d-1}(g_0,g_2,g_3,g_4,...,g_d)} \bar\theta_{(0124...d)}^{n_{d-1}(g_0,g_1,g_2,g_4...,g_d)} ...\end{aligned} \tag{7}$$

where $\nu_d^+(g_0, g_1, ..., g_d)$ is a pure phase factor and $\theta_{(ij...k)}$ or $\bar\theta_{(ij...k)}$ is the Grassmann number associated with $(d-1)$-dimensional subsimplex on Σ_d^0, $(ij...k)$. The constraint is implemented explicitly by writing, say, $\theta_{(1234...d)}^{n_{1234...d}}$ as $\theta_{(1234...d)}^{n_{d-1}(g_1,g_2,g_3,g_4,...,g_d)}$.

Note that the vertices on the branched simplex Σ_d^0 have a natural order: vertex-0 $<$ vertex-1 $<$ ... $<$ vertex-d. This leads to the ordering of the Grassmann numbers: the first Grassmann number is associated with the subsimplex that does not contain the vertex-0, the second Grassmann number is associated with the subsimplex that does not contain the vertex-2, *etc* . Then it is followed by the Grassmann number associated with the subsimplex that does not contain the vertex-1, the Grassmann number associated with the subsimplex that does not contain the vertex-3, *etc* .

The above example is for simplices with a "+" orientation. For branched simplex Σ_d^0 with a "−" orientation, we have

$$\begin{aligned}\mathcal{V}_d^-(g_0, g_1, ..., g_d) &= \nu_d^-(g_0, g_1, g_2, g_3, g_4, ..., g_d)\times \\ &...\theta_{(0124...d)}^{n_{d-1}(g_0,g_1,g_2,g_4...,g_d)} \theta_{(0234...d)}^{n_{d-1}(g_0,g_2,g_3,g_4,...,g_d)} \times \\ &...\bar\theta_{(0134...d)}^{n_{d-1}(g_0,g_1,g_3,g_4,...,g_d)} \bar\theta_{(1234...d)}^{n_{d-1}(g_1,g_2,g_3,g_4,...,g_d)}\end{aligned} \tag{8}$$

where the order of the Grassmann numbers is reversed and θ's are switched with $\bar\theta$'s.

The number of the Grassmann numbers on the $(d-1)$-dimensional simplex $(ij...k)$ is given by

$$n_{d-1}(g_i, g_j, ..., g_k) = 0, 1. \tag{9}$$

which depends on d variables $g_i, g_j, ..., g_k$, and must satisfy

$$\sum_{i=0}^{d} n_{d-1}(g_0, ..., \hat g_i, ..., g_d) = \text{ even}, \tag{10}$$

so that the Grassmann rank $(1+d)$-tensor always has an even number of Grassmann numbers. Here the sequence $g_0, ..., \hat g_i, ..., g_d$ is the sequence $g_0, ..., g_d$ with g_i removed.

All the solution of eqn. (10) can be obtained using an integer function $m_{d-2}(g_1, g_2, ..., g_{d-1}) = 0, 1$ with $d-1$ variables (while $n_{d-1}(g_1, g_2, ..., g_d)$ is an integer function of d variables):

$$\begin{aligned}&n_{d-1}(g_1, g_2, ..., g_d) \\ &= \sum_{i=1}^{d} m_{d-2}(g_1, ..., \hat g_i, ..., g_d) \text{ mod } 2.\end{aligned} \tag{11}$$

For example, a rank 1-tensor has a form

$$\mathcal{V}_0(g_0) = \nu_0(g_0) \tag{12}$$

which contains no Grassmann number. A rank 2-tensor has a form

$$\mathcal{V}_1^+(g_0, g_1) = \nu_1^+(g_0, g_1)\theta_{(1)}^{n_0(g_1)} \bar\theta_{(0)}^{n_0(g_0)} \tag{13}$$

where $n_0(g)$ has only two consistent choices

$$n_0(g) = 0, \quad \forall g \in G_b, \quad \text{and} \quad n_0(g) = 1, \quad \forall g \in G_b. \tag{14}$$

As another example, a rank 4-tensor $\mathcal{V}_3^+$ for the tetrahedron in Fig. 2a is given by

$$\begin{aligned}\mathcal{V}_3^+(g_0, g_1, g_2, g_3) &= \nu_3^+(g_0, g_1, g_2, g_3)\times \\ &\theta_{(123)}^{n_2(g_1,g_2,g_3)} \theta_{(013)}^{n_2(g_0,g_1,g_3)} \bar\theta_{(023)}^{n_2(g_0,g_2,g_3)} \bar\theta_{(012)}^{n_2(g_0,g_1,g_2)}.\end{aligned} \tag{15}$$

Note that the four triangles of the tetrahedron have different orientations: the triangles (123) and (013) point outwards and the triangles (023) and (012) point inwards. We have used θ and $\bar\theta$ for triangles with different orientations.

For the tetrahedron in Fig. 2b, which has an opposite orientation to that of the tetrahedron in Fig. 2a, we have a rank 4-tensor $\mathcal{V}_3^-$ which is given by

$$\begin{aligned}\mathcal{V}_3^-(g_0, g_1, g_2, g_3) &= \nu_3^-(g_0, g_1, g_2, g_3)\times \\ &\theta_{(012)}^{n_2(g_0,g_1,g_2)} \theta_{(023)}^{n_2(g_0,g_2,g_3)} \bar\theta_{(013)}^{n_2(g_0,g_1,g_3)} \bar\theta_{(123)}^{n_2(g_1,g_2,g_3)}\end{aligned} \tag{16}$$

Note the different order of the Grassmann numbers $\theta_{(012)}^{n_2(g_0,g_1,g_2)} \theta_{(023)}^{n_2(g_0,g_2,g_3)} \bar\theta_{(013)}^{n_2(g_0,g_1,g_3)} \bar\theta_{(123)}^{n_2(g_1,g_2,g_3)}$.

E. Evaluation of the Grassmann tensors on a complex

Let Σ_d be a d-complex with a branching structure which is formed by several simplices $[ab...c]$. As discussed in the last section, a simplex is associated with one of the two Grassmann tensors $\mathcal{V}_d^\pm$. Let us use $\int_{\text{in}(\Sigma_d)} \prod_{[ab...c]} \mathcal{V}_d^{s(a,b,...,c)}$ to represent the evaluation of rank $(1+d)$ Grassmann tensors $\mathcal{V}_d^\pm$ on the d-complex

Σ_d:

$$\begin{aligned}&\int_{\text{in}(\Sigma_d)} \prod_{[ab...c]} \mathcal{V}_d^{s(a,b,...,c)} \\ &\equiv \int \prod_{(ij...k)} \mathrm{d}\theta_{(ij...k)}^{n_{d-1}(g_i,g_j,...,g_k)} \mathrm{d}\bar\theta_{(ij...k)}^{n_{d-1}(g_i,g_j,...,g_k)} \times \\ &\prod_{\{xy...z\}} (-)^{m_{d-2}(g_x,g_y,...,g_z)} \prod_{[ab...c]} \mathcal{V}_d^{s(a,b,...,c)}(g_a,g_b,...,g_c)\end{aligned} \tag{17}$$

where $\prod_{[ab...,c]}$ multiplies over all the d-dimensional simplices $[ab...c]$ in Σ_d and $\mathcal{V}_d^{s(a,b,...,c)}(g_a,g_b,...,g_c)$ is a Grassmann rank $(1+d)$-tensor associated with the simplex $[ab...c]$. $s(a,b,...,c) = \pm$ depending on the orientation of the d-dimensional simplex $[ab...c]$. Also $\prod_{(ij...k)}$ multiplies over all the interior $(d-1)$-dimensional simplices, $(ij...k)$, of the complex Σ_d [*ie* those $(d-1)$-dimensional simplices that are not on the surface of Σ_d]. $\prod_{\{xy...z\}}$ multiplies over all the interior $(d-2)$-dimensional simplices [*ie* those $(d-2)$-dimensional simplices that are not on the surface of Σ_d], $\{xy...z\}$, of the complex Σ_d. We note that, when Σ_d is a single simplex $[ab...c]$, $\int_{\text{in}(\Sigma_d)} \prod_{[ab...c]} \mathcal{V}_d^{s(a,b,...,c)}$ is given by $\mathcal{V}_d^{s(a,b,...,c)}(g_a,g_b,...,g_c)$.

We see that in the formal notation, $\int_{\text{in}(\Sigma_d)} \prod_{[ab...c]} \mathcal{V}_d^{s(a,b,...,c)}$, $\int_{\text{in}(\Sigma_d)}$ represents a Grassmann integral over the Grassmann numbers on all the interior $(d-1)$-simplicies in the d-complex Σ_d. So $\text{in}(\Sigma_d)$ in $\int_{\text{in}(\Sigma_d)}$ actually represents the collection of all those interior $(d-1)$-simplicies in the d-complex Σ_d. The Grassmann numbers on the $(d-1)$-simplicies on the surface of the d-complex Σ_d are not integrated over. In fact $\int_{\text{in}(\Sigma_d)}$ has the following explicit form

$$\begin{aligned}\int_{\text{in}(\Sigma_d)} = \int \prod_{(ij...k)} \mathrm{d}\theta_{(ij...k)}^{n_{d-1}(g_i,g_j,...,g_k)} \mathrm{d}\bar\theta_{(ij...k)}^{n_{d-1}(g_i,g_j,...,g_k)} \times \\ \prod_{\{xy...z\}} (-)^{m_{d-2}(g_x,g_y,...,g_z)}\end{aligned} \tag{18}$$

We note that $\mathrm{d}\theta$ always appears in front of $\mathrm{d}\bar\theta$. We also note that the integration measure contains a non-trivial sign factor $(-)^{m_{d-2}(g_x,g_y,...,g_z)}$ on the interior $(d-2)$ simplices $\{xy...z\}$.

The sign factor $\prod_{\{xy...z\}}(-)^{m_{d-2}(g_x,g_y,...,g_z)}$ is included to help us to define topological fermionic path integral later. Choosing orientation dependent tensors $\mathcal{V}_d^\pm$ also help us to define topological fermionic path integral. Adding the sign factor $\prod_{\{xy...z\}}(-)^{m_{d-2}(g_x,g_y,...,g_z)}$ and choosing orientation dependent tensors $\mathcal{V}_d^\pm$ appear to be very unnatural. However, they are two extremely important features of our approach. We cannot obtain topological fermionic path integral without these two features. Realizing these two features is one of a few breakthroughs that allows us to obtain topological fermionic path integral on discrete space-time. The two features are related to our choice of the ordering convention of θ and $\bar\theta$ in the Grassmann tensors $\mathcal{V}_d^\pm$, and the ordering convention of $\mathrm{d}\theta$ and $\mathrm{d}\bar\theta$ in the integration measure. We have not find an ordering convention to remove the sign factor $\prod_{\{xy...z\}}(-)^{m_{d-2}(g_x,g_y,...,g_z)}$.

Now let us consider two d-complices Σ_d^1 and Σ_d^2, which do not overlap but may share part of their surfaces. Let Σ_d be the union of the two d-complices. From our definition of the Grassmann integral, we find that the Grassmann integral on Σ_d can be expressed as

$$\begin{aligned}&\int_{\text{in}(\Sigma_d)} \prod_{[ab...c]} \mathcal{V}_d^{s(a,b,...,c)} \\ &= \int_{\Sigma_d^1\cap\Sigma_d^2} \Big(\int_{\text{in}(\Sigma_d^1)} \prod_{[ab...c]} \mathcal{V}_d^{s(a,b,...,c)} \int_{\text{in}(\Sigma_d^2)} \prod_{[ab...c]} \mathcal{V}_d^{s(a,b,...,c)} \Big)\end{aligned} \tag{19}$$

where $\Sigma_d^1 \cap \Sigma_d^2$ is the intersection of the two complices, which contains only $(d-1)$-simplices on the shared surface of the two complices Σ_d^1 and Σ_d^2. More precisely

$$\begin{aligned}\int_{\Sigma_d^1\cap\Sigma_d^2} = \int \prod_{(ij...k)} \mathrm{d}\theta_{(ij...k)}^{n_{d-1}(g_i,g_j,...,g_k)} \mathrm{d}\bar\theta_{(ij...k)}^{n_{d-1}(g_i,g_j,...,g_k)} \times \\ \prod_{\{xy...z\}} (-)^{m_{d-2}(g_x,g_y,...,g_z)}\end{aligned} \tag{20}$$

where $\prod_{(ij...k)}$ is a product over all the $(d-1)$-simplices in $\Sigma_d^1\cap\Sigma_d^2$ and $\prod_{\{xy...z\}}$ is a product over all the interior $(d-2)$-simplices in $\Sigma_d^1\cap\Sigma_d^2$ (note that $\Sigma_d^1\cap\Sigma_d^2$ is a $(d-1)$-complex). Eqn. (19) describes the process to glue the two complices Σ_d^1 and Σ_d^2 together.

F. An example for evaluation of Grassmann tensor network

Let us consider an example to help us to understand the complicated expression (17). When Σ is formed by two tetrahedrons as in Fig. 3, the top tetrahedron with a "+" orientation is associated with a rank 4-tensor $\mathcal{V}_3^+$:

$$\begin{aligned}&\mathcal{V}_3^+(g_0,g_1,g_2,g_3) = \nu_3^+(g_0,g_1,g_2,g_3)\times \\ &\quad \theta_{(123)}^{n_2(g_1,g_2,g_3)} \theta_{(013)}^{n_2(g_0,g_1,g_3)} \bar\theta_{(023)}^{n_2(g_0,g_2,g_3)} \bar\theta_{(012)}^{n_2(g_0,g_1,g_2)}\end{aligned} \tag{21}$$

The lower tetrahedron with a "−" orientation is associated with a rank 4-tensor $\mathcal{V}_3^-$:

$$\begin{aligned}&\mathcal{V}_3^-(g_0,g_1,g_2,g_4) = \nu_3^-(g_0,g_1,g_2,g_4)\times \\ &\quad \theta_{(012)}^{n_2(g_0,g_1,g_2)} \theta_{(024)}^{n_2(g_0,g_2,g_4)} \bar\theta_{(014)}^{n_2(g_0,g_1,g_4)} \bar\theta_{(124)}^{n_2(g_1,g_2,g_4)}\end{aligned} \tag{22}$$

Note that $\mathcal{V}_3^+(g_0,g_1,g_2,g_3)$ and $\mathcal{V}_3^-(g_0,g_1,g_2,g_4)$ have independent Grassmann numbers $\theta_{(012)}$ and $\bar\theta_{(012)}$ on the shared face (012). $\bar\theta_{(012)}$ belongs to $\mathcal{V}_3^-(g_0,g_1,g_2,g_4)$ and $\theta_{(012)}$ belongs to $\mathcal{V}_3^+(g_0,g_1,g_2,g_3)$. Since $\text{in}(\Sigma) = (012)$, the evaluation of $\mathcal{V}_3^{\pm}$ on the complex Σ is given by

$$\begin{aligned}&\int_{(012)} \mathcal{V}_3^{+}(g_0,g_1,g_2,g_3)\mathcal{V}_3^{-}(g_0,g_1,g_2,g_4)\\ &= \int \mathrm{d}\theta_{(012)}^{n_2(g_0,g_1,g_2)}\mathrm{d}\bar\theta_{(012)}^{n_2(g_0,g_1,g_2)}\times\\ &\quad \mathcal{V}_3^{+}(g_0,g_1,g_2,g_3)\mathcal{V}_3^{-}(g_0,g_1,g_2,g_4) \qquad (23)\end{aligned}$$

which is a special case of eqn. (17).

IV. GRASSMANN TENSOR NETWORK AND FERMIONIC PATH INTEGRAL

Given a closed space-time manifold, M_{ST}, with a time direction, its triangularization Σ is a complex with no boundary. The time direction gives rise to a local order of the vertices. So the complex Σ has a branching structure.

Since Σ has no boundary, the evaluation of Grassmann tensors $\mathcal{V}_d^{\pm}$ on it gives rise to a complex number:

$$e^{-S} = \int_{\text{in}(\Sigma)} \prod_{[ab\dots c]} \mathcal{V}_d^{s(a,b,\dots,c)}. \qquad (24)$$

Such a complex number can be viewed as the action-amplitude in the imaginary-time path integral of a fermionic system. The different choices of the complex functions $\nu_d^{\pm}(g_0,\dots,g_d)$ and the integer functions $m_{d-2}(g_0,\dots,g_{d-2})$ correspond to different choices of Lagrangian of the fermionic system. The partition function of the imaginary-time path integral is given by

$$Z = \sum_{\{g_i\}} \int_{\text{in}(\Sigma)} \prod_{[ab\dots c]} \mathcal{V}_d^{s(a,b,\dots,c)}, \qquad (25)$$

where each vertex of the complex Σ is associated with a variable g_i. We note that

the fermionic partition function is determined from two $(1+d)$-variable complex functions: $\nu_d^{+}(g_i,g_j,\dots)$, $\nu_d^{-}(g_i,g_j,\dots)$, and one $(d-1)$-variable integer function $m_{d-2}(g_i,g_j,\dots)$.

V. TOPOLOGICALLY INVARIANT GRASSMANN TENSOR NETWORK AND FERMIONIC TOPOLOGICAL NON-LINEAR σ-MODEL

Now we would like to study the low energy effective theory of a gapped fermion system. The fixed point low energy effective action-amplitude of a gapped system must describe a topological quantum field theory. So the fixed point action-amplitude must be invariant under a renormalization group (RG) flow which generates coarse-graining transformation of the space-time complex Σ. In fact, such a fixed-point requirement under RG flow completely fixes the form of the fixed-point action-amplitude. The fixed-point action-amplitude, in turn, classifies the possible fermionic SPT phases.



FIG. 4: (a) Consider two triangles which are a part of a space-time complex Σ with a branching structure. The first type of the RG flow step changes the two triangles to other two triangles. The branching structure leads to the following vertex ordering $(0,1,2,3)$. (b) A different space-time complex Σ which corresponds to a different ordered vertices $(0,3,1,2)$. The corresponding complex does have a valid branching structure because some triangles overlap.

The detailed RG flow steps depend on the dimensions of the space-time. We will discuss different space-time dimensions and the corresponding fixed-point action-amplitude separately. Here we will only give a brief discussion. A more detailed discussion will be given in Appendix A.

A. (1+1)D case

We first consider fermion systems in $(1+1)$ space-time dimension. Its imaginary-time path integral is given by

$$Z = \sum_{\{g_i\}} \int_{\text{in}(\Sigma_2)} \prod_{[abc]} \mathcal{V}_2^{s(a,b,c)}, \qquad (26)$$

where Σ_2 is the (1+1)D space-time complex with a branching structure (see Fig. 4), and $\mathcal{V}_2^{\pm}$ are rank-3 Grassmann tensors.

The first type of the RG flow step that changes the space-time complex is described by Fig. 4a. If the Grassmann tensors $\mathcal{V}_2^{\pm}$ describe a fixed point action-amplitude, their evaluation on the two complices in Fig. 4a should be the same. This leads to the following condition

$$\begin{aligned}&\nu_2(g_0,g_1,g_3)\nu_2(g_1,g_2,g_3)\\ &= \nu_2(g_0,g_1,g_2)\nu_2(g_0,g_2,g_3) \qquad (27)\end{aligned}$$

where ν_2 is a complex phase related to $\nu_2^{\pm}$ and $m_0(g)$ through

$$\begin{aligned}\nu_2^{+}(g_0,g_1,g_2) &= \nu_2(g_0,g_1,g_2),\\ \nu_2^{-}(g_0,g_1,g_2) &= (-)^{m_0(g_1)}/\nu_2(g_0,g_1,g_2). \qquad (28)\end{aligned}$$



FIG. 5: (a) Consider three triangles which are a part of a space-time complex Σ with a branching structure. The second type of the RG flow step changes the three triangles to one triangle. The branching structure leads to the following vertex ordering $(0, 1, 3, 2)$. (b) The vertex order $(0, 1, 2, 3)$ does not correspond to a valid branching structure.



FIG. 6: (a) A 3D complex formed by two 3-simplices (g_0, g_1, g_2, g_3) and (g_1, g_2, g_3, g_4). The two 3-simplices share a 2-simplex (g_1, g_2, g_3). (b) A 3D complex formed by three 3-simplices (g_0, g_2, g_3, g_4), (g_0, g_1, g_3, g_4), and (g_0, g_1, g_2, g_4).

Note that the above condition is obtained for a particular vertex ordering $(0, 1, 2, 3)$ (a particular branching structure) as described in Fig. 4a. Different *valid* branching structures, in general, lead to other conditions on $\nu_2(g_0, g_1, g_2)$. It turns out that all those conditions are equivalent to eqn. (27). For example, for vertex ordering $(1, 3, 0, 2)$, we obtain

$$\begin{aligned}&\nu_2(g_1, g_0, g_2)\nu_2(g_1, g_3, g_0)\\ &= \nu_2(g_1, g_3, g_2)\nu_2(g_3, g_0, g_2)\end{aligned} \tag{29}$$

which is the same as eqn. (27) after replacing $g_1 \to g_0$, $g_3 \to g_1$, $g_0 \to g_2$, and $g_2 \to g_3$.

Some vertex orders do not correspond to valid branching structure due to the overlap of the simplices, and are not considered. One of the invalid ordering is described by Fig. 4b which has a vertex order $(0, 3, 1, 2)$. The invalid orderings can give rise to conditions on $\nu_2(g_0, g_1, g_2)$ that are *not* equivalent to eqn. (27).



FIG. 7: (a) A 3D complex formed by one 3-simplex (g_0, g_1, g_2, g_4). (b) A 3D complex formed by four 3-simplices (g_1, g_2, g_3, g_4), (g_0, g_2, g_3, g_4), (g_0, g_1, g_3, g_4), and (g_0, g_1, g_2, g_3).



FIG. 8: (Color online) (a) A 4D complex formed by two 4-simplices $(g_0, g_1, g_2, g_3, g_4)$ and $(g_1, g_2, g_3, g_4, g_5)$. The two 4-simplices share a 3-simplex (g_1, g_2, g_3, g_4) (color red). (b) A 4D complex formed by four 4-simplices $(g_0, g_2, g_3, g_4, g_5)$, $(g_0, g_1, g_3, g_4, g_5)$, $(g_0, g_1, g_2, g_4, g_5)$, and $(g_0, g_1, g_2, g_3, g_5)$. The two simplices $(g_0, g_1, g_2, g_4, g_5)$, and $(g_0, g_1, g_2, g_3, g_5)$ share a 3-simplex (g_0, g_1, g_2, g_5) (color blue).

The second type of the RG flow step that changes the space-time complex is described by Fig. 5. Different *valid* branching structures (see Fig. 5a for one example), lead to different conditions on $\nu_2(g_0, g_1, g_2)$ (which is defined in eqn. (28)). It turns out that all those conditions are equivalent to eqn. (27). (Some vertex orderings do not correspond to valid branching structures. One of the invalid ordering is described by Fig. 5b.)

Since $\nu_2(g_0, g_1, g_2)$ is related to $\nu_2^{\pm}(g_0, g_1, g_2)$ and $m_0(g)$ through eqn. (28), eqn. (27) is actually a condition on $\nu_2^{\pm}(g_0, g_1, g_2)$ and $m_0(g)$ that determines the fermion path integral. The fermion path integral described by $\nu_2^{\pm}(g_0, g_1, g_2)$ and $m_0(g)$ that satisfies eqn. (27) is a topological fermion path integral which is a fixed-point theory.

We would like to point out that eqn. (27) is the standard 2-cocycle condition of group cohomology.[50,51]

B. (2+1)D case

We next consider fermion systems in (2+1) space-time dimension described by imaginary-time path integral

$$Z = \sum_{\{g_i\}} \int_{\text{in}(\Sigma_3)} \prod_{[abcd]} \mathcal{V}_3^{s(a,b,c,d)}, \tag{30}$$

where Σ_3 is the (2+1)D space-time complex with a branching structure and $\mathcal{V}_3^{\pm}$ are rank-4 Grassmann tensors.

The first type of the RG flow step that changes the space-time complex is described by Fig. 6. In order for the Grassmann tensors $\mathcal{V}_3^{\pm}$ to describe a fixed point action-amplitude, their evaluations on the two complices



FIG. 9: (Color online) (a) A 4D complex formed by one 4-simplex (g_0,g_1,g_2,g_3,g_4). (b) A 4D complex formed by five 4-simplices (g_1,g_2,g_3,g_5,g_4), (g_0,g_2,g_3,g_5,g_4), (g_0,g_1,g_3,g_5,g_4), (g_0,g_1,g_2,g_5,g_4), and (g_0,g_1,g_2,g_3,g_5). The two simplices (g_0,g_1,g_2,g_5,g_4), and (g_0,g_1,g_2,g_3,g_5) share a 3-simplex (g_0,g_1,g_2,g_5) (color blue).

in Fig. 6 should be the same. This leads to the following condition

$$\begin{aligned}&\nu_3(g_1,g_2,g_3,g_4)\nu_3(g_0,g_1,g_3,g_4)\nu_3(g_0,g_1,g_2,g_3) = \\ &(-)^{n_2(g_0,g_1,g_2)n_2(g_2,g_3,g_4)}\nu_3(g_0,g_2,g_3,g_4)\nu_3(g_0,g_1,g_2,g_4)\end{aligned} \tag{31}$$

where ν_3 is a complex phase and is given by

$$\begin{aligned}\nu_3^+(g_0,g_1,g_2,g_3) &= (-)^{m_1(g_0,g_2)}\nu_3(g_0,g_1,g_2,g_3), \\ \nu_3^-(g_0,g_1,g_2,g_3) &= (-)^{m_1(g_1,g_3)}/\nu_3(g_0,g_1,g_2,g_3)\end{aligned} \tag{32}$$

Different vertex orders give rise to different branching structures. It turns out that all the different *valid* branching structures give rise to the same condition (31).

The second type of the RG flow step that changes the space-time complex is described by Fig. 7. If the Grassmann tensors $\mathcal{V}_3^\pm$ describe a fixed point action-amplitude, their evaluations on the two complices in Fig. 7 should also be the same. This leads to the same condition (31).

C. (3+1)D case

Last, we consider fermion systems in (3+1) space-time dimension, described by imaginary-time path integral

$$Z = \sum_{\{g_i\}} \int_{\text{in}(\Sigma_4)} \prod_{[abcde]} \mathcal{V}_4^{s(a,b,c,d,e)}, \tag{33}$$

where Σ_4 is the (3+1)D space-time complex with a branching structure and $\mathcal{V}_4^\pm$ are rank-5 Grassmann tensors.

The first type of the RG flow step that changes the space-time complex is described by Fig. 8. The second type of the RG flow step that changes the space-time complex is described by Fig. 9. If the Grassmann tensor $\mathcal{V}_4$ describes a fixed point action-amplitude, its evaluation on the two complices in Fig. 8 or Fig. 9 should be the same. This leads to the following condition

$$\begin{aligned}&\nu_4(g_1,g_2,g_3,g_4,g_5)\nu_4(g_0,g_1,g_3,g_4,g_5)\nu_4(g_0,g_1,g_2,g_3,g_5) \\ &= (-)^{n_3(g_0,g_1,g_2,g_3)n_3(g_0,g_3,g_4,g_5)+n_3(g_1,g_2,g_3,g_4)n_3(g_0,g_1,g_4,g_5)+n_3(g_2,g_3,g_4,g_5)n_3(g_0,g_1,g_2,g_5)}\times \\ &\quad \nu_4(g_0,g_2,g_3,g_4,g_5)\nu_4(g_0,g_1,g_2,g_4,g_5)\nu_4(g_0,g_1,g_2,g_3,g_4)\end{aligned} \tag{34}$$

where ν_4 is a complex phase and is given by

$$\begin{aligned}&\nu_4^+(g_0,g_1,g_2,g_3,g_4) = \\ &(-)^{m_2(g_0,g_1,g_3)+m_2(g_1,g_3,g_4)+m_2(g_1,g_2,g_3)}\nu_4(g_0,g_1,g_2,g_3,g_4), \\ &\nu_4^-(g_0,g_1,g_2,g_3,g_4) = (-)^{m_2(g_0,g_2,g_4)}/\nu_4(g_0,g_1,g_2,g_3,g_4)\end{aligned} \tag{35}$$

The sign factors in above relations between $\nu_4^\pm$ and ν_4, and the sign factor in eqn. (34) are very strange. Obtaining those strange sign factors is another break through that allows us to obtain topological fermionic path integral beyond (1+1) dimensions. It appears that those sign factors are related to a deep mathematical structure called Steenrod squares.[64,65] Different vertex orders give rise to different branching structures. It turns out that all the valid branching structures give rise to the same condition (34).

D. The fixed-point action on a closed complex and the unitary condition

We know that the fixed-point theory of a bosonic SPT phase is a discrete bosonic topological non-linear σ-model.[50] The bosonic topological non-linear σ-models are characterized by action-amplitudes that is equal to 1 on any closed complex. The action-amplitude being equal to 1 on any closed complex is the key reason why the models describe bosonic SPT phases without intrinsic topological orders.

In Section V, we have developed a fixed-point theory of fermionic SPT phases, and defined discrete fermionic topological non-linear σ-model. A discrete fermionic topological non-linear σ-model is described by a pair of functions $[\nu_d(g_0,...,g_d), m_{d-2}(g_0,...,g_{d-2})]$, that satisfy the conditions eqn. (27), eqn. (31), or eqn. (34). According to the branching moves discussed above, the evaluation on any closed manifold is reduced to an equivalent evaluation

$$\int \mathcal{V}_d^+ \mathcal{V}_d^- \tag{36}$$

where the symbol $\int$ means integrating out all the Grassmann variables on the corresponding $d-1$ simplices (see eqn. (18)). We note that the sign factor $\prod_{\{x,y,\dots,z\}}(-)^{m_{d-2}(g_x,g_y,\dots,g_z)}$ should be included in the fermionic path integral. In Appendix B we will show that $\int \mathcal{V}_d^+ \mathcal{V}_d^-$ is always equal to 1.

We also note that the bosonic topological non-linear σ-models are characterized by action-amplitudes that are pure $U(1)$ phases $|\nu_d(g_0,\dots,g_d)| = 1$ (where the action-amplitude on "+" oriented simplices is given by $\nu_d(g_0,\dots,g_d)$, and the action-amplitude on "−" oriented simplices is given by $\nu_d^{-1}(g_0,\dots,g_d) = \nu_d^*(g_0,\dots,g_d)$). The action-amplitude being a pure $U(1)$ phase ensures that the model defined by the path integral to be unitary theory.

But what is the analogue of the pure $U(1)$ phase condition on the Grassmann amplitude $\mathcal{V}_d^\pm$ (or on the pair of functions $[\nu_d(g_0,\dots,g_d), m_{d-2}(g_0,\dots,g_{d-2})]$)? Clearly, the Grassmann amplitude $\mathcal{V}_d^\pm$ are not even complex numbers. It is hard to say when $\mathcal{V}_d^\pm$ behave like $U(1)$ phases.

To address this issue, let us introduce complex conjugate of a quantity that contains Grassmann numbers:

$$\Big(\nu\theta_1\bar\theta_2\theta_3\dots\Big)^* \equiv \nu^*\dots\bar\theta_3\theta_2\bar\theta_1. \tag{37}$$

We see that under complex conjugate a) the complex coefficients are complex conjugated, b) the order of the Grassmann number is reversed, and c) θ and $\bar\theta$ is exchanged. With this definition, one is tempting to require $\mathcal{V}_d^+(\mathcal{V}_d^+)^* = 1$, in order for $\mathcal{V}_d^+$ to be a $U(1)$ phase. But, $\mathcal{V}_d^+(\mathcal{V}_d^+)^*$ still contains Grassmann numbers and we cannot require it to be 1. So we try to require $\int \mathcal{V}_d^+(\mathcal{V}_d^+)^* = 1$ where the Grassmann integral is defined in eqn. (18). Now $\int \mathcal{V}_d^+(\mathcal{V}_d^+)^*$ is a complex number. But $\int \mathcal{V}_d^+(\mathcal{V}_d^+)^*$ is not non-negative. So we cannot treat $\int \mathcal{V}_d^+(\mathcal{V}_d^+)^*$ as a norm-square of $\mathcal{V}_d^+$. After some considerations, we find that we need to define a different complex conjugate:

$$\begin{aligned}
[\mathcal{V}_1^\pm(g_0,g_1)]^\dagger &= [\mathcal{V}_1^\pm(g_0,g_1)]^*,\\
[\mathcal{V}_2^\pm(g_0,g_1,g_2)]^\dagger &= (-)^{m_0(g_1)}[\mathcal{V}_2^\pm(g_0,g_1,g_2)]^*,\\
[\mathcal{V}_3^\pm(g_0,..,g_3)]^\dagger &= (-)^{m_1(g_0,g_2)+m_1(g_1,g_3)}[\mathcal{V}_3^\pm(g_0,..,g_3)]^*,\\
[\mathcal{V}_4^\pm(g_0,\dots,g_4)]^\dagger &= [\mathcal{V}_4^\pm(g_0,\dots,g_4)]^* \times\\
&\quad (-)^{m_1(g_0,g_1,g_3)+m_1(g_1,g_3,g_4)+m_1(g_1,g_2,g_3)+m_1(g_0,g_2,g_4)}
\end{aligned} \tag{38}$$

Using the new complex conjugate and the evaluation of $\int \mathcal{V}_d^+ \mathcal{V}_d^-$ described above, we find that

$$\int \mathcal{V}_d^+[\mathcal{V}_d^+]^\dagger = \int \mathcal{V}_d^-[\mathcal{V}_d^-]^\dagger = \nu_d\nu_d^*. \tag{39}$$

Thus we would like to require the Grassmann amplitude $\mathcal{V}_d^\pm$ to satisfy

$$\int \mathcal{V}_d^\pm[\mathcal{V}_d^\pm]^\dagger = 1. \tag{40}$$

which is to require $|\nu_d(g_0,\dots,g_d)| = 1$. Eqn. (40) is the analogue of the pure $U(1)$ phase condition on the Grassmann amplitude $\mathcal{V}_d^\pm$.

For the Grassmann amplitude $\mathcal{V}_d^\pm$ that satisfy the pure $U(1)$ phase condition (40), we have

$$[\mathcal{V}_d^\pm]^\dagger = \mathcal{V}_d^\mp. \tag{41}$$

E. Symmetry and stability of the fixed-point action amplitude

We have seen that a fermionic system can be described by a fermionic path integral on the discretized space-time. In d dimensional space-time, the fermionic path integral is determined by two $(d+1)$-variable complex functions $\nu_d^\pm(g_0,\dots,g_d)$ and one $(d-1)$-variable integer function $m_{d-2}(g_0,\dots,g_{d-2})$. In the last few sections, we have shown that if $\nu_d^\pm(g_0,\dots,g_d)$ and $m_{d-2}(g_0,\dots,g_{d-2})$ satisfy some conditions, eqn. (27), eqn. (31), and eqn. (34), the fermionic path integral determined from $\nu_d^\pm(g_0,\dots,g_d)$ and $m_{d-2}(g_0,\dots,g_{d-2})$ is actually a fixed-point theory under the coarse-graining transformation of the space-time complex.

A fixed-point theory can be used to describe a phase if the fixed-point theory is stable. To see whether a fixed-point theory is stable or not, we perturb the fixed-point action amplitude $\nu_d^\pm(g_0,\dots,g_d)$ by a small amount $\nu_d^\pm(g_0,\dots,g_d) \to \nu_d^\pm(g_0,\dots,g_d)+\delta\nu_d^\pm(g_0,\dots,g_d)$. If $\nu_d^\pm(g_0,\dots,g_d)+\delta\nu_d^\pm(g_0,\dots,g_d)$ flows back to $\nu_d^\pm(g_0,\dots,g_d)$ under the coarse-graining transformation of space-time complex, the fixed-point theory is stable.

From this point of view, our constructed fixed-point theories described by $\nu_d^\pm(g_0,\dots,g_d)$ and $m_{d-2}(g_0,\dots,g_{d-2})$ are not stable. If we add a perturbation of form $\delta\nu_d^\pm(g_0,\dots,g_d) = \epsilon\prod_{i=0}^d \delta_{g_i,\bar g}$ to increase the weight of $\bar g$ state in the action amplitude, we believe that the action amplitude will flow to the simple form $\nu_d^\pm(g_0,\dots,g_d) = \prod_{i=0}^d \delta_{g_i,\bar g}$ where only the $\bar g$ state has a non-zero weight.

However, under a symmetry condition, our constructed fixed-point theories can be stable. To impose a symmetry condition, we assume that our fermion system has full symmetry group G_f. Let $G_b = G_f/Z_2^f$, we will take g_i in $\nu_d^\pm(g_0,\dots,g_d)$ and $m_{d-2}(g_0,\dots,g_{d-2})$ as a group element in G_b. From Section III C, we see that different degrees of freedom are attached to the $(d-1)$-simplices for different choices of $g_i,\dots,g_k$. The degrees of freedom on a $(d-1)$-simplex form a 1D representation u_{d-1}^g of the symmetry $g \in G_f$, which may depend on $g_i,\dots,g_k$:

$$\begin{aligned}
u_{d-1}^{g_1g_2}(g_i,g_j,\dots,g_k) &= u_{d-1}^{g_1}(g_i,g_j,\dots,g_k)u_{d-1}^{g_2}(g_i,g_j,\dots,g_k)\\
u_{d-1}^{g}(g'g_i,g_j,\dots,g'g_k) &= u_{d-1}^{g}(g_i,g_j,\dots,g_k),
\end{aligned} \tag{42}$$

We also require the variables with bar and without bar to carry opposite quantum number (*ie* their 1D representations are inverse of each other). In this case, the symmetry of the path integral (*ie* the invariance of the action amplitude) can be implemented by requiring:

$$\begin{aligned} \theta_{(ij\dots k)} &\to u_g^{-1}(g_i,g_j,\dots,g_k)\theta_{(ij\dots k)}, \\ \bar\theta_{(ij\dots k)} &\to u_g(g_i,g_j,\dots,g_k)\bar\theta_{(ij\dots k)}, \end{aligned} \tag{43}$$

and

$$\begin{aligned} &(\nu_d^\pm(gg_0,..,gg_d))^{s(g)} \\ &= \nu_d^\pm(g_0,..,g_d)\prod_i (u^g_{d-1})^{(-)^i}(g_0,..,\hat g_i,..,g_d), \\ &m_{d-2}(gg_0,..,gg_{d-2}) = m_{d-2}(g_0,..,g_{d-2}),\ g\in G_f. \end{aligned} \tag{44}$$

where $s(g) = -1$ if g contains the anti unitary time-reversal transformation and $s(g) = 1$ otherwise. If we view $\mathcal{V}_d^\pm(g_0,..,g_d))$ as a whole object, it is invariant under the symmetry action g:

$$\mathcal{V}_d^\pm(gg_0,..,gg_d))^{s(g)} = \mathcal{V}_d^\pm(g_0,..,g_d) \tag{45}$$

Indeed, the above symmetric $\nu_d^\pm(g_0,...,g_d)$ and $m_{d-2}(g_0,...,g_{d-2})$ describe a fermion system with G_f symmetry. If $\nu_d^\pm(g_0,...,g_d)$ and $m_{d-2}(g_0,...,g_{d-2})$ further satisfy eqn. (27), eqn. (31), or eqn. (34), we believe, they even describe a fixed-point theory that is stable against perturbations which do not break the symmetry. But further consideration suggests that such a symmetry condition is too strong. The fermionic systems that are described by the symmetric $\nu_d^\pm(g_0,...,g_d)$ and $m_{d-2}(g_0,...,g_{d-2})$ appear to be essentially bosonic. At least, their classification is the same as the classification of bosonic SPT phases.

So in this paper, we will use a weaker symmetry condition. We only require $\nu_d(g_0,...,g_d)$ and $n_{d-1}(g_0,...,g_{d-1})$ (which are certain combinations of $\nu_d^\pm(g_0,...,g_d)$ and $m_{d-2}(g_0,...,g_{d-2})$) to be symmetric

$$\begin{aligned} &\nu_d^{s(g)}(gg_0,..,gg_d) \\ &= \nu_d(g_0,..,g_d)\prod_i (u^g_{d-1})^{(-)^i}(g_0,..,\hat g_i,..,g_d), \\ &n_{d-1}(gg_0,..,gg_{d-1}) = n_{d-1}(g_0,..,g_{d-1}),\ g\in G_f. \end{aligned} \tag{46}$$

Since the fermion fields always change sign under the fermion-number-parity transformation $P_f\in G_f$, the 1D representations satisfy

$$u_{P_f}(g_0,..,g_{d-1}) = (-)^{n_{d-1}(g_0,...,g_{d-1})}. \tag{47}$$

Note that we only require $n_{d-1}(g_0,...,g_{d-1})$ to be symmetric. In general, $m_{d-2}(g_0,...,g_{d-2})$ is not symmetric $m_{d-2}(gg_0,...,gg_{d-2}) \neq m_{d-2}(g_0,...,g_{d-2})$. Although the fermionic path integral appears not to be symmetric under the weaker symmetry condition, it turns out that the symmetry breaking is only a boundary effect and the symmetry can be restored by adding an additional boundary action to the space-time path integral. In fact, we will show that the ground state wavefunction and the Hamiltonian obtained from the path integral with the weaker symmetry condition are indeed symmetric. So a fermion path integral determined by $\nu_d(g_0,...,g_d)$ and $n_{d-1}(g_0,...,g_{d-1})$ that satisfy eqn. (46) as well as eqn. (27), eqn. (31), or eqn. (34) does describe a symmetric stable fixed-point theory. Such $\nu_d(g_0,...,g_d)$, $n_{d-1}(g_0,...,g_{d-1})$, and $u^g_{d-1}(g_0,..,g_{d-1})$ describe a fermionic topological phase with symmetry.

Let us introduce

$$\begin{aligned} f_1(g_0,g_1) &= 0; \\ f_2(g_0,g_1,g_2) &= 0; \\ f_3(g_0,g_1,...,g_3) &= 0; \\ f_4(g_0,g_1,...,g_4) &= n_2(g_0,g_1,g_2)n_2(g_2,g_3,g_4); \\ f_5(g_0,g_1,...,g_5) &= \quad n_3(g_0,g_1,g_2,g_3)n_3(g_0,g_3,g_4,g_5) \\ &\quad + n_3(g_1,g_2,g_3,g_4)n_3(g_0,g_1,g_4,g_5) \\ &\quad + n_3(g_2,g_3,g_4,g_5)n_3(g_0,g_1,g_2,g_5). \end{aligned} \tag{48}$$

Using f_d, we can define a mapping from $(d+1)$-variable functions $\nu_d(g_0,...,g_d)$ to $(d+2)$-variable functions $(\delta\nu_d)(g_0,...,g_{d+1})$:

$$\begin{aligned} &(\delta\nu_d)(g_0,...,g_{d+1}) \\ &\equiv (-)^{f_{d+1}(g_0,...,g_{d+1})}\prod_{i=0}^{d+1}\nu_d^{(-)^i}(g_0,..,\hat g_i,..,g_{d+1}). \end{aligned} \tag{49}$$

Then using $(\delta\nu_d)$, we can rewrite the conditions, eqn. (27), eqn. (31), and eqn. (34), in a uniform way

$$(\delta\nu_d)(g_0,...,g_{d+1}) = 1. \tag{50}$$

Here $[\nu_d(g_0,...,g_d), n_{d-1}(g_0,...,g_{d-1}), u^g_{d-1}(g_0,...,g_{d-1})]$ that satisfies eqn. (46), eqn. (10), and eqn. (50), will be called a d-cocycle. The space of d-cocycle is denoted as $\mathscr{Z}^d[G_f, U_T(1)]$. The fermionic path integral obtained from a d-cocycle $(\nu_d, n_{d-1}, u^g_{d-1})$ (via $\nu_d^\pm(g_0,...,g_d)$ and $m_{d-2}(g_0,...,g_{d-2})$) will be called a fermionic topological non-linear σ-model.

In the following, we will show that we can construct an exactly soluble local fermionic Hamiltonian in d spatial dimensions from each $(d+1)$-cocycle (ν_{d+1}, n_d, u^g_d) in $\mathscr{Z}^{d+1}[G_f, U_T(1)]$. The Hamiltonian has a symmetry G_f. The ground state wave function of the constructed Hamiltonian can also be obtained exactly from the $(d+1)$-cocycle. Such a ground state does not break the symmetry G_f and describes a fermionic SPT phase.

VI. IDEAL GROUND STATE WAVE FUNCTION

We have shown that from each element (ν_{d+1}, n_d, u^g_d) of $\mathscr{Z}^{d+1}[G_f, U_T(1)]$, we can define a fermionic topological non-linear σ-model in $(d+1)$ space-time dimensions. The action amplitude $\mathcal{V}^\pm_{d+1}$ of the model [obtained from



FIG. 10: A 9-site triangular lattice on a torus, where each site i has physical states $|g_i\rangle$ where $g_i \in G_b$, and each triangle has fermionic states $|n_{ijk}\rangle$ where $n_{ijk} = 01$ is the fermion occupation number. The orientations on the edges give rise to a natural order of the three vertices of a triangle (i,j,k) where the first vertex i of a triangle has two outgoing edges on the triangle and the last vertex k of a triangle has two incoming edges on the triangle. The triangular lattice can be viewed as the surface of solid torus. A discretization of the solid torus can be obtained by adding a vertex-0 inside the solid torus. The branching structure of the resulting complex is indicated the arrows on the edges.

$(\nu_{d+1}, n_d, u_d^g)]$ is a fixed-point action amplitude under the coarse-graining transformation of the space-time complex. The fermionic path integral is supposed to give us a quantum ground state. We claim that such a quantum ground state is a fermionic SPT phase described by (ν_{d+1}, n_d, u_d^g). In the section, we will construct the ground state wave function, in (2+1)D as an example.

A. Construction of 2D wave function

We will assume that our 2D system lives on a triangular lattice which forms a 2D torus (see Fig. 10). On each lattice site i, we have physical states $|g_i\rangle$ labeled by $g_i \in G_b$. On each triangle (ijk), we have two states: no fermion state $|0\rangle$ and one fermion state $|1\rangle$.

The ideal ground state wave function can be obtained by viewing the 2D torus as the surface of a 3D solid torus. The fermionic path integral on the 3D solid torus with the action amplitude $\mathcal{V}_{d+1}^{\pm}$ obtained from the cocycle (ν_{d+1}, n_d, u_d^g) should give us the ground state wave function for the corresponding fermionic SPT state. To do the fermionic path integral, we need to divide the 3D solid torus into a 3D complex with a branching structure. Due to the topological invariance of our constructed action amplitude, the resulting wave function should not depend on how we divide the 3D solid torus into 3D complex. So we choose a very simple 3D complex which is formed by the triangular lattice on the surface of the 3D solid torus and one additional vertex-0 inside the 3D solid torus. The resulting 3D complex is formed by simplices $[0ijk]$, where ijk is a triangle on the surface (see Fig. 10). The branching structure of the 3D complex is given by the orientations on the edges. Those orientations for the edges on the surface are given in Fig. 10. For the edges inside the 3D solid torus, the orientation is always pointing from vertex-0 to the vertex on the surface. We note that the simplices associated with the down triangles in Fig. 10 have a "+" orientation while the simplices associated with the up triangles have a "−" orientation.

From each 3-cocycle (ν_3, n_2, u_2^g) in $\mathscr{Z}^3[G_f, U_T(1)]$, we can construct a fixed-point action amplitude. For simplices with "+" orientation, the fixed-point action amplitude is given by

$$\mathcal{V}_3^+(g_0,g_i,g_j,g_k) = \nu_3^+(g_0,g_i,g_j,g_k)\times \theta_{(123)}^{n_2(g_i,g_j,g_k)}\theta_{(013)}^{n_2(g_0,g_i,g_k)}\bar\theta_{(023)}^{n_2(g_0,g_j,g_k)}\bar\theta_{(012)}^{n_2(g_0,g_i,g_j)} \qquad (51)$$

and for simplices with "−" orientation given by

$$\mathcal{V}_3^-(g_0,g_i,g_j,g_k) = \nu_3^-(g_0,g_i,g_j,g_k)\times \theta_{(012)}^{n_2(g_0,g_i,g_j)}\theta_{(023)}^{n_2(g_0,g_j,g_k)}\bar\theta_{(013)}^{n_2(g_0,g_i,g_k)}\bar\theta_{(123)}^{n_2(g_i,g_j,g_k)}, \qquad (52)$$

where

$$\begin{aligned}\nu_3^+(g_0,g_i,g_j,g_k) &= (-)^{m_1(g_0,g_j)}\nu_3(g_0,g_i,g_j,g_k),\\ \nu_3^-(g_0,g_i,g_j,g_k) &= (-)^{m_1(g_i,g_k)}/\nu_3(g_0,g_i,g_j,g_k),\end{aligned} \qquad (53)$$

Since $n_2(g_0,g_1,g_2)$ is in $\mathcal{H}^2(G,\mathbb{Z}_2)$, we can always write $n_2(g_0,g_1,g_2)$ in terms of $m_1(g_0,g_1)$:

$$\begin{aligned}n_2(g_0,g_1,g_2) &= m_1(g_1,g_2)+m_1(g_0,g_2)+m_1(g_0,g_1) \text{ mod } 2,\\ m_1(g_0,g_1) &= 0,1.\end{aligned} \qquad (54)$$

However, in general $m_1(gg_0,gg_1) \neq m_1(g_0,g_1)$, even though $n_2(g_0,g_1,g_2)$ satisfies $n_2(gg_0,gg_1,gg_2) = n_2(g_0,g_1,g_2)$.

Now, the wave function is given by

$$\begin{aligned}&\Psi(\{g_i\},\{\theta_{(ijk)}\},\{\bar\theta_{(ijk)}\})\\ &= \int \prod_{(0ij)} \mathrm{d}\theta_{(0ij)}^{n_2(g_0,g_i,g_j)}\mathrm{d}\bar\theta_{(0ij)}^{n_2(g_0,g_i,g_j)} \prod_{\{0i\}}(-)^{m_1(g_0,g_i)}\times\\ &\quad\prod_{\triangle}\mathcal{V}_3^-(g_0,g_i,g_j,g_k)\prod_{\triangledown}\mathcal{V}_3^+(g_0,g_i,g_j,g_k),\end{aligned} \qquad (55)$$

where $\prod_{(0ij)}$ is the product over all links of the triangular lattice, $\prod_{\{0i\}}$ is the product over all sites of the triangular lattice, $\prod_{\triangle}$ is the product over all up-triangles, and $\prod_{\triangledown}$ is the product over all down-triangles. Clearly, the above wave function depends on g_i through ν_3, n_2 and m_1. The wave function also appears to depend on g_0, the variable that we assigned to the vertex-0 inside the solid torus. In fact, the g_0 dependence cancels out, and the wave function is independent of g_0. We can simply set $g_0 = 1$.

If all the m_1 dependence also cancels out, the wave function will have a symmetry described by G_f: $\Psi(\{gg_i\},\{\theta_{(ijk)}\},\{\bar\theta_{(ijk)}\}) = \Psi^{s(g)}(\{g_i\},\{\theta_{(ijk)}\},\{\bar\theta_{(ijk)}\})$. But does the m_1 dependence cancel out? Let us only write down the m_1

dependence of the wave function:

$$\begin{aligned}&\Psi(\{g_i\},\{\theta_{(ijk)}\},\{\bar\theta_{(ijk)}\})\\ &=\prod_{\{0i\}}(-)^{m_1(g_0,g_i)}\prod_{\triangle}(-)^{m_1(g_i,g_k)}\prod_{\triangledown}(-)^{m_1(g_0,g_j)}\dots\dots\\ &=\prod_{\triangle}(-)^{m_1(g_i,g_k)}\dots\dots \end{aligned} \tag{56}$$

where ijk around the up- and down-triangles are arranged in a way as illustrated in Fig. 10. We also have used the relation $\prod_{\{0i\}}(-)^{m_1(g_0,g_i)}\prod_{\triangledown}(-)^{m_1(g_0,g_j)}=1$. We see that m_1 does not cancel out and the wave function is not symmetric, since $m_1(gg_0,gg_1)\neq m_1(g_0,g_1)$.

This is a serious problem. But the symmetry breaking is only on the surface and it can be easily fixed: we simply redefine the wave function by including an extra factor $\prod_{\text{up-left}}(-)^{m_1(g_i,g_k)}$ on the surface:

$$\begin{aligned}&\Psi(\{g_i\},\{\theta_{(ijk)}\},\{\bar\theta_{(ijk)}\}) \\ &=\prod_{\text{up-left}}(-)^{m_1(g_i,g_k)}\int\prod_{(0ij)}\mathrm{d}\theta_{(0ij)}^{n_2(g_0,g_i,g_j)}\mathrm{d}\bar\theta_{(0ij)}^{n_2(g_0,g_i,g_j)}\times\\ &\prod_{\{0i\}}(-)^{m_1(g_0,g_i)}\prod_{\triangle}\mathcal{V}_3^-(g_0,g_i,g_j,g_k)\prod_{\triangledown}\mathcal{V}_3^+(g_0,g_i,g_j,g_k)\\ &=\int\prod_{(0ij)}\mathrm{d}\theta_{(0ij)}^{n_2(g_0,g_i,g_j)}\mathrm{d}\bar\theta_{(0ij)}^{n_2(g_0,g_i,g_j)}\times\\ &\qquad\prod_{\triangle}\nu_3^{-1}(g_0,g_i,g_j,g_k)\prod_{\triangledown}\nu_3(g_0,g_i,g_j,g_k)\times\\ &\qquad\prod_{\triangle}\theta_{(0ij)}^{n_2(g_0,g_i,g_j)}\theta_{(0jk)}^{n_2(g_0,g_j,g_k)}\bar\theta_{(0ik)}^{n_2(g_0,g_i,g_k)}\bar\theta_{(ijk)}^{n_2(g_i,g_j,g_k)}\times\\ &\qquad\prod_{\triangledown}\theta_{(ijk)}^{n_2(g_i,g_j,g_k)}\theta_{(0ik)}^{n_2(g_0,g_i,g_k)}\bar\theta_{(0jk)}^{n_2(g_0,g_j,g_k)}\bar\theta_{(0ij)}^{n_2(g_0,g_i,g_j)}\end{aligned} \tag{57}$$

where $\prod_{\text{up-left}}$ is a product over all the links with the up-left orientation (note that $\prod_{\text{up-left}}(-)^{m_1(g_i,g_k)}=\prod_{\triangle}(-)^{m_1(g_i,g_k)}$ and see Fig. 10). The redefined wave function is indeed symmetric, and is the wave function that corresponds to the fermionic SPT state described by the cocycle $(\nu_3,n_2)\in\mathscr{Z}^3[G_f,U_T(1)]$.

We note that the wave function $\Psi(\{g_i\},\{\theta_{(ijk)}\},\{\bar\theta_{(ijk)}\})$ described above contains Grassmann numbers. Indeed, $\Psi(\{g_i\},\{\theta_{(ijk)}\},\{\bar\theta_{(ijk)}\})$ can be regarded as the wave function in the fermion coherent state basis. After we expand the wave function in power of the Grassmann numbers, we obtain

$$\begin{aligned}&\Psi(\{g_i\},\{\theta_{(ijk)}\},\{\bar\theta_{(ijk)}\})\\ &=\sum_{n_{ijk}=0,1}\Phi(\{g_i\},\{n_{ijk}\})\prod_{\triangle}\bar\theta_{(ijk)}^{n_{ijk}}\prod_{\triangledown}\theta_{(ijk)}^{n_{ijk}}.\end{aligned} \tag{58}$$

Then $\Phi(\{g_i\},\{n_{ijk}\})$ is the amplitude of the ground state on the fermion number basis $\otimes_i|g_i\rangle\otimes_{(ijk)}|n_{ijk}\rangle$, where $|g_i\rangle$ is the state on site-i and $|n_{ijk}\rangle$ is the state on triangle (ijk) (where $n_{ijk}=0,1$ is the fermion occupation number on the triangle (ijk)). Note that the sign of $\Phi(\{g_i\},\{n_{ijk}\})$ will depend on how the Grassmann numbers are ordered in $\prod_{\triangle}\bar\theta_{(ijk)}^{n_{ijk}}\prod_{\triangledown}\theta_{(ijk)}^{n_{ijk}}$. We also note that the wave function vanishes if $n_{ijk}\neq n_2(g_i,g_j,g_k)$.

In Appendix C, we study the fermionic cocycles (ν_{d+1},n_d,u_d^g) systematically. In particular, we will study the equivalence relation between them. This will lead to the notion of group super-cohomology class $\mathscr{H}^{d+1}[G_f,U_T(1)]$. In Appendix G 1, we will show the elements in $\mathscr{H}^{d+1}[G_f,U_T(1)]$ correspond to distinct fermionic SPT phases. We will calculate $\mathscr{H}^{d+1}[G_f,U_T(1)]$ for some simple symmetry groups G_f in Appendix F. This allows us to construct several new fermionic SPT phases. The results are summarized in Table II.

In Appendix H, we further show that for each ground state wave function derived from (ν_{d+1},n_d,u_d^g), one can explicitly construct an exact solvable hermitian (local) Hamiltonian $H=-\sum_i H_i$. Similar as the bosonic case, H_i acts on site i as well as its 6 neighbor. However, since there are 6 more fermionic degrees of freedom in the 6 triangles surround i, H_i also acts on these 6 triangles. Moreover, all the H_i are hermitian (unconstraint)commuting projectors satisfying $H_i^2=H_i$ and $H_iH_j=H_jH_i$. Such a nice (frustration free)property make it exact solvable, with $\Psi(\{g_i\},\{\theta_{(ijk)}\},\{\bar\theta_{(ijk)}\})$ as its unique ground state.

B. No intrinsic topological orders

In this section, we are going to show that the wave function constructed in the last section $\Phi(\{g_i\},\{n_{ijk}\})$ contain no intrinsic topological orders as a fermion system. In other words, starting with the following pure bosonic direct product state

$$|\Phi_0\rangle=\otimes_i|\phi_i\rangle,\quad |\phi_i\rangle=|G_b|^{-1/2}\sum_{g_i\in G_b}|g_i\rangle, \tag{59}$$

we can obtain the fermionic state $|\Phi\rangle$ constructed in last section after a fermionic LU transformation defined in Ref. 25.

To show this, we start with an expand fermionic Hilbert space, where we have four fermionic orbitals within each triangles. We also have bosonic state $|g_i\rangle$ on each vertex (see Fig. 11). In the Grassmann number form, the constructed fermionic wave function is given by



FIG. 11: (Color online) A triangular lattice with four fermionic orbitals in each triangle (ijk). The fermions on the two solid dots in the yellow triangle are described by operators $c_{(ijk)}$ (the center one) and $c_{(0ik)}$ (the side one), and the fermions on the two open dots are described by operators $\bar c_{(0ij)}$ and $\bar c_{(0jk)}$. The fermions on the two open dots in the blue triangle are described by operators $\bar c_{(ijk)}$ (the center one) and $\bar c_{(0ik)}$ (the side one), and the fermions on the two solid dots are described by operators $c_{(0ij)}$ and $c_{(0jk)}$. Each vertex has bosonic states described by $|g_i\rangle$, $g_i \in G_b$.

eqn. (60):

$$\begin{aligned}
&\Psi(\{g_i\},\{\theta_{(ijk)}\},\{\bar\theta_{(ijk)}\}) \\
&= \int \prod_{(0ij)} \mathrm{d}\theta_{(0ij)}^{n_2(g_0,g_i,g_j)} \mathrm{d}\bar\theta_{(0ij)}^{n_2(g_0,g_i,g_j)} \times \\
&\quad \prod_{\triangle} \nu_3^{-1}(g_0,g_i,g_j,g_k) \prod_{\bigtriangledown} \nu_3(g_0,g_i,g_j,g_k) \times \\
&\quad \prod_{\triangle} \theta_{(0ij)}^{n_2(g_0,g_i,g_j)} \theta_{(0jk)}^{n_2(g_0,g_j,g_k)} \bar\theta_{(0ik)}^{n_2(g_0,g_i,g_k)} \bar\theta_{(ijk)}^{n_2(g_i,g_j,g_k)} \times \\
&\quad \prod_{\bigtriangledown} \theta_{(ijk)}^{n_2(g_i,g_j,g_k)} \theta_{(0ik)}^{n_2(g_0,g_i,g_k)} \bar\theta_{(0jk)}^{n_2(g_0,g_j,g_k)} \bar\theta_{(0ij)}^{n_2(g_0,g_i,g_j)}
\end{aligned} \tag{60}$$

We would like to point out that although the above expression contains g_0, the topological invariance of the fermion path integral ensures that the Grassmann wave function on the left-hand-side does not depend on g_0.

If we treat θ's and $\bar\theta$'s as the following complex fermion operators

$$\begin{aligned}
\theta_{(abc)} &= c^\dagger_{(abc)}, \quad \bar\theta_{(abc)} = \bar c^\dagger_{(abc)}, \\
\mathrm{d}\theta_{(abc)} &= c_{(abc)}, \quad \mathrm{d}\bar\theta_{(abc)} = \bar c_{(abc)},
\end{aligned} \tag{61}$$

the expression eqn. (60) can be viewed as an operator:

$$\begin{aligned}
\hat U = &\prod_{(0ij)} c_{(0ij)}^{n_2(g_0,g_i,g_j)} \bar c_{(0ij)}^{n_2(g_0,g_i,g_j)} \times \\
&\prod_{\triangle} \nu_3^{-1}(g_0,g_i,g_j,g_k) \prod_{\bigtriangledown} \nu_3(g_0,g_i,g_j,g_k) \times \\
&\prod_{\triangle} c_{(0ij)}^{\dagger n_2(g_0,g_i,g_j)} c_{(0jk)}^{\dagger n_2(g_0,g_j,g_k)} \bar c_{(0ik)}^{\dagger n_2(g_0,g_i,g_k)} \bar c_{(ijk)}^{\dagger n_2(g_i,g_j,g_k)} \times \\
&\prod_{\bigtriangledown} c_{(ijk)}^{\dagger n_2(g_i,g_j,g_k)} c_{(0ik)}^{\dagger n_2(g_0,g_i,g_k)} \bar c_{(0jk)}^{\dagger n_2(g_0,g_j,g_k)} \bar c_{(0ij)}^{\dagger n_2(g_0,g_i,g_j)}.
\end{aligned} \tag{62}$$

Again $\hat U$ is independent of g_0, despite the appearance of g_0 on the right-hand-side. Then the fermionic state constructed in the last section can be obtained from the bosonic product state $|\Phi_0\rangle$:

$$|\Psi\rangle = \hat U |\Phi_0\rangle. \tag{63}$$

Now we would like to point out that $\hat U$ itself is formed by several layers of *fermionic* LU transformations. Since ν_3 is a pure $U(1)$ phase, thus $\prod_\triangle \nu_3^{-1}(g_0,g_i,g_j,g_k) \prod_\bigtriangledown \nu_3(g_0,g_i,g_j,g_k)$ represents layers of bosonic LU transformations. Also the $\hat U$ has a property that when acting on $|\Phi_0\rangle$, $c^\dagger$ and $\bar c^\dagger$ always act on states with no fermion and c and $\bar c$ always act on states with one fermion. So those operators map a set of orthogonal states to another set of orthogonal states. In this case, an even numbers of c's and $\bar c$'s correspond a fermionic LU transformation. Therefore, the terms in $\hat U$, such as $c_{(ijk)}^{\dagger n_2(g_i,g_j,g_k)} c_{(0ik)}^{\dagger n_2(g_0,g_i,g_k)} \bar c_{(0jk)}^{\dagger n_2(g_0,g_j,g_k)} \bar c_{(0ij)}^{\dagger n_2(g_0,g_i,g_j)}$ and $c_{(0ij)}^{n_2(g_0,g_i,g_j)} \bar c_{(0ij)}^{n_2(g_0,g_i,g_j)}$, all represent fermionic LU transformation as defined in Ref. 25. Note that none of the above transformations changes g_i. So we can show those transformations to be unitary within each fixed set of $\{g_i\}$. Therefore, the state $|\Psi\rangle$ has no fermionic long range entanglement as defined in Ref. 25 (*ie* no fermionic intrinsic topological order). $|\Psi\rangle$ is the fermionic SPT state described by a cocycle (ν_3, n_2, u_2^g).

VII. EXAMPLES

A. 2D fermionic SPT state with Z_2 symmetry

In Appendix F, we have calculated the group super-cohomology classes $\mathscr{H}^3[Z_2\times Z_2^f, U(1)] = \mathbb{Z}_2^2$. This means that interacting fermion systems with a Z_2 symmetry can have (at least) 4 different SPT phases: a trivial one plus three non-trivial ones. This result is described by the $Z_2 \times Z_2^f$ row and $d_{sp} = 2$ column of Table II.

In the following, we will list the explicit fermionic 3-cocycles $[\nu_3(g_0,g_1,g_2,g_3), n_2(g_0,g_1,g_2)]$, $g_i \in G_b = G_f/Z_2^f$, that describe the above four phases. The first two SPT phases are given by

$$\begin{aligned}
& n_2(g_0,g_1,g_2) = 0, \\
& \nu_3(e,o,e,o) = \nu_3(o,e,o,e) = \pm 1, \qquad \text{other } \nu_3 = 1.
\end{aligned} \tag{64}$$

where we have assumed that the elements in $G_b = Z_2$ are described by $\{e, o\}$ with e being the identity element. The two 3-cocycles actually describe bosonic SPT phases [the Z_n row and $d_{sp} = 2$ column of Table I (with $n = 2$)]. The 3-cocycle $\nu_3(e,o,e,o) = \nu_3(o,e,o,e) = 1$ describes the trivial SPT phase and the 3-cocycle $\nu_3(e,o,e,o) = \nu_3(o,e,o,e) = -1$ describes the non-trivial bosonic SPT phase, protected by the on-site Z_2 symmetry

$$\hat W^g = \prod_{\boldsymbol{i}} |g_{\boldsymbol{i}}\rangle\langle g g_{\boldsymbol{i}}|, \qquad g, g_{\boldsymbol{i}} \in Z_2. \tag{65}$$

If we choose the bulk Hamiltonian to be the ideal Hamiltonian $H = \sum_{\boldsymbol{i}} H_{\boldsymbol{i}}$ with $H_{\boldsymbol{i}}$ given by eqn. (H9), the low energy edge excitations will have a zero energy (zero bandwidth). So we will add the following Z_2 symmetric edge Hamiltonian to give those zero energy edge excitations a non-trivial dynamics:

$$H_{\text{edge}} = \sum_i [J\sigma_i^z\sigma_{i+1}^z + h\sigma_i^x], \tag{66}$$

where σ_i^l are the Pauli matrices acting on the two states $|e\rangle$ and $|o\rangle$ on site-i. From eqn. (G23) and eqn. (G28), we find that the low energy effective edge Hamiltonian is given by

$$H_{\text{eff}} = \sum_i [J\sigma_i^z\sigma_{i+1}^z + h\sigma_i^x] \tag{67}$$

for the trivial SPT phase with $\nu_3(e,o,e,o) = \nu_3(o,e,o,e) = 1$; and by

$$H_{\text{eff}} = \sum_i [J\sigma_i^z\sigma_{i+1}^z + \frac{h}{2}\sigma_i^x(1 - \sigma_{i-1}^z\sigma_{i+1}^z)] \tag{68}$$

for the non-trivial SPT phase with $\nu_3(e,o,e,o) = \nu_3(o,e,o,e) = -1$.

The other two SPT phases are given by

$$\begin{aligned} n_2(e,o,e) = n_2(o,e,o) = 1, &\quad \text{other } n_2 = 0, \\ \nu_3(e,o,e,o) = \nu_3(o,e,o,e) = \pm\mathrm{i}, &\quad \text{other } \nu_3 = 1. \end{aligned} \tag{69}$$

They describe non-trivial intrinsic fermionic SPT phases. The ideal bulk Hamiltonian is given by $H = \sum_{\boldsymbol{i}} H_{\boldsymbol{i}}$ with $H_{\boldsymbol{i}}$ given by eqn. (H9).

Again, for the ideal bulk Hamiltonian, the low energy edge excitations all have a zero energy. Let us add the following Z_2 symmetric edge Hamiltonian to give the zero energy edge excitations a non-trivial dynamics:

$$\begin{aligned} H_{\text{edge}} = \sum_i [&-J\sigma_i^z\sigma_{i+1}^z + h_{00}\sigma_i^x \\ &+ (h_{+-}\sigma_i^x c^\dagger_{(i,i+1)}c_{(i-1,i)} + h_{++}\sigma_i^x c^\dagger_{(i,i+1)}c^\dagger_{(i-1,i)} + h.c.)], \end{aligned} \tag{70}$$

where a fermion lives on each link $(i, i+1)$. From eqn. (G23), eqn. (G28) and eqn. (G30), we find that the low energy effective edge Hamiltonian is given by

$$\begin{aligned} H_{\text{eff}} = \sum_i \Big[& J\sigma_i^z\sigma_{i+1}^z + \frac{h_{00}}{2}\sigma_i^x(\sigma_{i-1}^z - \sigma_{i+1}^z) \\ &+ [\frac{h_{+-}}{4}\sigma_i^-(1+\sigma_{i-1}^z)(1-\sigma_{i+1}^z) \\ &- \frac{h_{+-}}{4}\sigma_i^+(1-\sigma_{i-1}^z)(1+\sigma_{i+1}^z) + h.c.]\Big] \\ = \sum_i \Big[& -J\sigma_i^z\sigma_{i+1}^z + \frac{h_{00}+\mathrm{Re}(h_{+-})}{2}\sigma_i^x(\sigma_{i-1}^z - \sigma_{i+1}^z) \\ &+ \frac{\mathrm{Im}(h_{+-})}{2}(\sigma_i^y - \sigma_{i-1}^z\sigma_i^y\sigma_{i+1}^z)\Big] \end{aligned} \tag{71}$$

for both the fermionic SPT phases with $\nu_3(e,o,e,o) = \nu_3(o,e,o,e) = \pm i$; where

$$\sigma^- = \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix}, \quad \sigma^+ = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}. \tag{72}$$

Note that the h_{++} term vanishes in the low energy edge sector.

Let us call a link $(i, i+1)$ with $\sigma_i^z\sigma_{i+1}^z = -1$ a domain wall. We note that all the three terms in the above H_{eff} cannot change the numbers of domain walls. So the model has an effective $U(1)$ symmetry. It has two phases: gapped phase for large J where $\sigma^z = \pm 1$ and there are no domain walls; gapless phase for large h_{00} where the domain walls form a "superfluid". The gapped phase breaks the Z_2 symmetry while the gapless phase is described by a central-charge $c = 1$ conformal field theory.

From the expression eqn. (G22) of the low energy edge state, we see that the total number of fermions in the low energy edge states $|\{g_i\}_{\text{edge}}\rangle$ are given by

$$N_f = \sum_i n_2(g_a, g_{i+1}, g_i). \tag{73}$$

We note that the above sum does not depend on g_a and is only a function of g_i's. If we choose $g_a = e$, we see that only the domain wall $(g_i, g_{i+1}) = (e, o)$ has one fermion, while the domain wall $(g_i, g_{i+1}) = (o, e)$ has no fermion. Thus the number of fermions is equal to the half of the number of the domain walls.

To consider a more general edge effective Hamiltonian that does not conserve the domain wall numbers, we add the following edge terms:

$$H_{\text{edge}} = \sum_i h_{0+0+}\sigma_i^x\sigma_{i+2}^x c^\dagger_{(i,i+1)}c^\dagger_{(i+2,i+3)} + h.c..$$

The corresponding edge effective Hamiltonian is given by

$$\begin{aligned} H_{\text{edge}} = \sum_i \frac{\mathrm{i}\,\mathrm{Im}(h_{0+0+})}{2}&(\sigma_i^+\sigma_{i+2}^+ - \sigma_i^-\sigma_{i+2}^-)\times \\ &(1+\sigma_{i-1}^z\sigma_{i+1}^z)(1+\sigma_{i+1}^z\sigma_{i+3}^z). \end{aligned} \tag{74}$$

So the more generic edge effective Hamiltonian has a form

$$\begin{aligned} H_{\text{edge}} = \sum_i \Big[& -J\sigma_i^z\sigma_{i+1}^z + h_0\sigma_i^x(\sigma_{i-1}^z - \sigma_{i+1}^z) \\ &+ h_1(\sigma_i^y - \sigma_{i-1}^z\sigma_i^y\sigma_{i+1}^z) \\ &+ \mathrm{i}h_2(\sigma_i^+\sigma_{i+2}^+ - \sigma_i^-\sigma_{i+2}^-)(1+\sigma_{i-1}^z\sigma_{i+1}^z)(1+\sigma_{i+1}^z\sigma_{i+3}^z)\Big]. \end{aligned} \tag{75}$$

The new term can only change the domain wall number by ±4, which can be irrelavent. So the edge excitations still be gapless.

From the structure of $n_2(g_0, g_1, g_2)$ in eqn. (69), we find that the number of fermions on the edge is given by half of the domain wall number. So the fermion-number-parity conservation only allows the domain wall number to change by a multiple of 4.

We note that $\frac{1}{2}\sum_{i=1}^{N}(1-\sigma_i^z\sigma_{i+1}^z)$ counts the number of domain walls, where N is the number sites on the 1D edge. Therefore, the following unitary operator

$$S=\exp\left(\frac{\mathrm{i}\pi}{8}\sum_{i=1}^{N}(1-\sigma_i^z\sigma_{i+1}^z)\right)\prod_{i=1}^{N}\sigma_i^y \tag{76}$$

satisfies $S^2=1$ in the subspace where the domain wall number is zero mode 4. So we can regard S as a Z_2 transformation in such a subspace. One can also check that the J, h_0 and h_1 terms in eqn. (75) commute with both $\exp\left(\frac{\mathrm{i}\pi}{8}\sum_{i=1}^{N}(1-\sigma_i^z\sigma_{i+1}^z)\right)$ and $\prod_{i=1}^{N}\sigma_i^y$, while the h_2 term in eqn. (75) anti-commute with both $\exp\left(\frac{\mathrm{i}\pi}{8}\sum_{i=1}^{N}(1-\sigma_i^z\sigma_{i+1}^z)\right)$ and $\prod_{i=1}^{N}\sigma_i^y$. So the edge effective Hamiltonian H (75) is invariant under S. Similar as the bosonic Z_2 SPT case[49,59], it can be shown that any 1D system satisfies such a non-local Z_2 symmetry must be gapless if the symmetry is not broken(spontaneously or explicitly).

B. 3D fermionic SPT state with $T^2=1$ time-reversal symmetry

In Appendix F, we have also calculated the group super-cohomology classes $\mathscr{H}^4[Z_2^T\times Z_2^f,U(1)]=\mathbb{Z}_2^2$. So interacting fermion systems with a $T^2=1$ time-reversal symmetry can have (at least) 4 different SPT phases: a trivial one and three non-trivial ones. This is described by the $Z_2^T\times Z_2^f$ row and $d_{sp}=3$ column of Table II.

Let us list the fermionic 4-cocycles $[\nu_4(g_0,g_1,g_2,g_3,g_4),n_3(g_0,g_1,g_2,g_3)]$ that describe the above four phases. The first two SPT phases are given by

$$\begin{aligned} &n_3(g_0,g_1,g_2,g_3)=0, \\ &\nu_4(e,o,e,o,e)=\nu_4(o,e,o,e,o)=\pm1, \quad \text{other } \nu_4=1. \end{aligned} \tag{77}$$

Again, the two 4-cocycles actually describe bosonic SPT phases (the Z_2^T row and $d_{sp}=3$ column of Table I).

The other two SPT phases are given by

$$\begin{aligned} &n_3(e,o,e,o)=n_3(o,e,o,e)=1, \quad \text{other } n_3=0, \\ &\nu_4(e,o,e,o,e)=-\nu_4(o,e,o,e,o)=\pm\mathrm{i}, \quad \text{other } \nu_4=1. \end{aligned} \tag{78}$$

They describe non-trivial intrinsic fermionic SPT phases.

Using the above cocycles, we can write down the corresponding wave functions and exactly soluble Hamiltonians. However, the explicit Hamiltonian is very complicated. We wonder if there exist a better basis, in which the Hamiltonian will have a simpler form.

VIII. SUMMARY

It was shown in Ref. 50 that generalized topological non-linear σ-models with symmetry can be constructed from group cohomology theory of the symmetry group. This leads to a systematic construction of bosonic SPT phases in any dimensions and for any symmetry groups. This result allows us to construct new topological insulators (with symmetry group $U(1)\times Z_2^T$) and new topological superconductors (with symmetry group Z_2^T) for interacting boson systems (or qubit systems). It also leads to a complete classification of all gapped phases in 1D interacting boson/fermion systems.

In this paper, we introduce a group super-cohomology theory which is a generalization of the standard group cohomology theory. Using the group super-cohomology theory, we can construct new discrete fermionic topological non-linear σ-models with symmetry. This leads to a systematic construction of fermionic SPT phases in any dimensions and for certain symmetry groups G_f where the fermions form an 1D representation. The discrete fermionic topological non-linear σ-model, when defined on a space-time with boundary, can be viewed as a "non-local" boundary effective Lagrangian, which is a fermionic and discrete generalization of the bosonic continuous Wess-Zumino-Witten term. Thus the boundary excitations of a non-trivial SPT phase is described by a "non-local" boundary effective Lagrangian, which, we believe, implies that the boundary excitations are gapless if the symmetry is not broken.

As an application of this general result, we constructed three non-trivial SPT phases in 3D, for interacting fermionic superconductors with coplanar spin order. Such superconductors have time-reversal Z_2^T and fermion-number-parity Z_2^f symmetries described by the full symmetry group $G_f=Z_2^T\times Z_2^f$.[44] The three non-trivial SPT phases should have gapless excitations on the 2D surface if the time-reversal symmetry is not broken. It is known that non-trivial 3D gapped topological phases do not exist for such superconductors if the fermions are non-interacting. So the constructed non-trivial fermionic SPT phases are new.

We also constructed three non-trivial SPT phases in 2D, for interacting fermionic systems with the full symmetry group $G_f=Z_2\times Z_2^f$. We show that the three non-trivial SPT phases indeed have gapless excitations on the 1D edge described by central-charge $c=1$ conformal field theory, if the Z_2 symmetry is not broken.

Clearly, more work is needed to develop a group super-cohomology theory for the cases when the fermions do not form an 1D representation. That will allow us to construct more general interacting fermionic SPT phases.

IX. ACKNOWLEDGEMENTS

We would like to thank Michael Freedman, Michael Levin, and Zhenghan Wang for many helpful discussions. Z.C.G. is supported by NSF Grant No. PHY05-51164. X.G.W. is supported by NSF Grant No. DMR-1005541 and NSFC 11074140.

Appendix A: Topological invariance of the partition amplitudes

In this section, we will prove the topological invariance of the partition amplitudes Eq. (25) in 1+1D and 2+1D under their corresponding fermionic group cocycle condition. The 3 + 1D case is much more complicated but can still be checked by computer.

1. 1 + 1D

Similar as we prove the topological invariance of the partition amplitudes that describe bosonic SPT phases[50], we can check all the admissible branching 2 ↔ 2 and 1 ↔ 3 moves for the Grassmann graded 2 cocycle function:

$$\mathcal{V}_2^{\pm}(g_i,g_j,g_k)=\nu_2^{\pm}(g_i,g_j,g_k)\theta_{(i,j)}^{n_1(g_i,g_j)}\theta_{(j,k)}^{n_1(g_j,g_k)}\bar{\theta}_{(i,k)}^{n_1(g_i,g_k)}. \tag{A1}$$



FIG. 12: The admissible branching 2 ↔ 2 moves. (a)Branching moves that can be induced by a global time ordering. (b)Branching moves that can not be induced by a global time ordering.



FIG. 13: The admissible branching 1 ↔ 3 moves. (a)Branching moves that can be induced by a global time ordering. (b)Branching moves that can not be induced by a global time ordering.

For admissible branching 2 ↔ 2 moves, we can have three different equations(up to the orientation conjugate + → −; − → +):

$$\int \mathrm{d}\theta_{(12)}^{n_1(g_1,g_2)}\mathrm{d}\bar{\theta}_{(12)}^{n_1(g_1,g_2)}\mathcal{V}_2^{+}(g_1,g_2,g_3)\mathcal{V}_2^{-}(g_0,g_1,g_2)$$
$$=\int \mathrm{d}\theta_{(03)}^{n_1(g_0,g_3)}\mathrm{d}\bar{\theta}_{(03)}^{n_1(g_0,g_3)}\mathcal{V}_2^{+}(g_0,g_2,g_3)\mathcal{V}_2^{-}(g_0,g_1,g_3) \tag{A2}$$

$$\int \mathrm{d}\theta_{(13)}^{n_1(g_1,g_3)}\mathrm{d}\bar{\theta}_{(13)}^{n_1(g_1,g_3)}\mathcal{V}_2^{+}(g_1,g_2,g_3)\mathcal{V}_2^{+}(g_0,g_1,g_3)$$
$$=\int \mathrm{d}\theta_{(02)}^{n_1(g_0,g_2)}\mathrm{d}\bar{\theta}_{(02)}^{n_1(g_0,g_2)}\mathcal{V}_2^{+}(g_0,g_2,g_3)\mathcal{V}_2^{+}(g_0,g_1,g_2), \tag{A3}$$

and

$$\int \mathrm{d}\theta_{(23)}^{n_1(g_2,g_3)}\mathrm{d}\bar{\theta}_{(23)}^{n_1(g_2,g_3)}\mathcal{V}_2^{-}(g_1,g_2,g_3)\mathcal{V}_2^{+}(g_0,g_2,g_3)$$
$$=\int \mathrm{d}\theta_{(01)}^{n_1(g_0,g_1)}\mathrm{d}\bar{\theta}_{(01)}^{n_1(g_0,g_1)}\mathcal{V}_2^{+}(g_0,g_1,g_3)\mathcal{V}_2^{-}(g_0,g_1,g_2). \tag{A4}$$

For admissible branching 1 ↔ 3 moves, we can have four different equations(up to the orientation conjugate $+ \to -; - \to +$):

$$\begin{aligned}
&\mathcal{V}_2^+(g_0,g_1,g_3)\\
&=\int \mathrm{d}\theta_{(02)}^{n_1(g_0,g_2)}\mathrm{d}\bar\theta_{(02)}^{n_1(g_0,g_2)}\mathrm{d}\theta_{(12)}^{n_1(g_1,g_2)}\mathrm{d}\bar\theta_{(12)}^{n_1(g_1,g_2)}\\
&\quad\times \mathrm{d}\theta_{(23)}^{n_1(g_2,g_3)}\mathrm{d}\bar\theta_{(23)}^{n_1(g_2,g_3)}(-)^{m_0(g_2)}\\
&\quad\times \mathcal{V}_2^-(g_1,g_2,g_3)\mathcal{V}_2^+(g_0,g_2,g_3)\mathcal{V}_2^+(g_0,g_1,g_2) \qquad \text{(A5)}\\
&\mathcal{V}_2^+(g_0,g_2,g_3)\\
&=\int \mathrm{d}\theta_{(01)}^{n_1(g_0,g_1)}\mathrm{d}\bar\theta_{(01)}^{n_1(g_0,g_1)}\mathrm{d}\theta_{(12)}^{n_1(g_1,g_2)}\mathrm{d}\bar\theta_{(12)}^{n_1(g_1,g_2)}\\
&\quad\times \mathrm{d}\theta_{(13)}^{n_1(g_1,g_3)}\mathrm{d}\bar\theta_{(13)}^{n_1(g_1,g_3)}(-)^{m_0(g_1)}\\
&\quad\times \mathcal{V}_2^+(g_1,g_2,g_3)\mathcal{V}_2^+(g_0,g_1,g_3)\mathcal{V}_2^-(g_0,g_1,g_2), \qquad \text{(A6)}
\end{aligned}$$

and

$$\begin{aligned}
&\mathcal{V}_2^+(g_1,g_2,g_3)\\
&=\int \mathrm{d}\theta_{(01)}^{n_1(g_0,g_1)}\mathrm{d}\bar\theta_{(01)}^{n_1(g_0,g_1)}\mathrm{d}\theta_{(02)}^{n_1(g_0,g_2)}\mathrm{d}\bar\theta_{(02)}^{n_1(g_0,g_2)}\\
&\quad\times \mathrm{d}\theta_{(03)}^{n_1(g_0,g_3)}\mathrm{d}\bar\theta_{(03)}^{n_1(g_0,g_3)}(-)^{m_0(g_0)}\\
&\quad\times \mathcal{V}_2^+(g_0,g_2,g_3)\mathcal{V}_2^-(g_0,g_1,g_3)\mathcal{V}_2^+(g_0,g_1,g_2) \qquad \text{(A7)}\\
&\mathcal{V}_2^+(g_0,g_1,g_2)\\
&=\int \mathrm{d}\theta_{(03)}^{n_1(g_0,g_3)}\mathrm{d}\bar\theta_{(03)}^{n_1(g_0,g_3)}\mathrm{d}\theta_{(13)}^{n_1(g_1,g_3)}\mathrm{d}\bar\theta_{(13)}^{n_1(g_1,g_3)}\\
&\quad\times \mathrm{d}\theta_{(23)}^{n_1(g_2,g_3)}\mathrm{d}\bar\theta_{(23)}^{n_1(g_2,g_3)}(-)^{m_0(g_3)}\\
&\quad\times \mathcal{V}_2^+(g_1,g_2,g_3)\mathcal{V}_2^-(g_0,g_2,g_3)\mathcal{V}_2^+(g_0,g_1,g_3). \qquad \text{(A8)}
\end{aligned}$$

Note that the first two equations of 2 ↔ 2 and 1 ↔ 3 moves can be induced by a global time ordering(see in Fig. 12 (a) and Fig. 13 (a)) while the rest can not(see in Fig. 12 (b) and Fig. 13 (b)). Here g^0 is defined on the vertex with no incoming edge, g^1 with one incoming edge, etc.

Let us use the definition of $\mathcal{V}_2^\pm$ and integral out the Grassmann variables. For the branching 2 ↔ 2 moves, we have:

$$\nu_2^+(g_1,g_2,g_3)\nu_2^-(g_0,g_1,g_2) =\nu_2^+(g_0,g_2,g_3)\nu_2^-(g_0,g_1,g_3) \qquad \text{(A9)}$$

$$\nu_2^+(g_1,g_2,g_3)\nu_2^+(g_0,g_1,g_3) =\nu_2^+(g_0,g_2,g_3)\nu_2^+(g_0,g_1,g_2), \qquad \text{(A10)}$$

and

$$\begin{aligned}
&\nu_2^-(g_1,g_2,g_3)\nu_2^+(g_0,g_2,g_3)\\
&=(-)^{n_1(g_1,g_2)}\nu_2^+(g_0,g_1,g_3)\nu_2^-(g_0,g_1,g_2). \qquad \text{(A11)}
\end{aligned}$$

Similarly, for the branching 1 ↔ 3 moves, we have:

$$\begin{aligned}
&\nu_2^+(g_0,g_1,g_3) \qquad \text{(A12)}\\
&=(-)^{m_0(g_2)}\nu_2^-(g_1,g_2,g_3)\nu_2^+(g_0,g_2,g_3)\nu_2^+(g_0,g_1,g_2)\\
&\nu_2^+(g_0,g_2,g_3) \qquad \text{(A13)}\\
&=(-)^{m_0(g_1)}\nu_2^+(g_1,g_2,g_3)\nu_2^+(g_0,g_1,g_3)\nu_2^-(g_0,g_1,g_2),
\end{aligned}$$

and

$$\begin{aligned}
&\nu_2^+(g_1,g_2,g_3) \qquad \text{(A14)}\\
&=(-)^{m_0(g_1)}\nu_2^+(g_0,g_2,g_3)\nu_2^-(g_0,g_1,g_3)\nu_2^+(g_0,g_1,g_2)\\
&\nu_2^+(g_0,g_1,g_2) \qquad \text{(A15)}\\
&=(-)^{m_0(g_2)}\nu_2^+(g_1,g_2,g_3)\nu_2^-(g_0,g_2,g_3)\nu_2^+(g_0,g_1,g_3).
\end{aligned}$$

If we use the definition of $\nu_2^\pm$:

$$\begin{aligned}
&\nu_2^+(g_0,g_1,g_2) = \nu_2(g_0,g_1,g_2),\\
&\nu_2^-(g_0,g_1,g_2) = (-)^{m_0(g_1)}/\nu_2(g_0,g_1,g_2), \qquad \text{(A16)}
\end{aligned}$$

All the above admissible branching moves will be equivalent to a single fermionic 2 cocycle equation of ν_2

$$\nu_2(g_0,g_1,g_3)\nu_2(g_1,g_2,g_3) = \nu_2(g_0,g_1,g_2)\nu_2(g_0,g_2,g_3). \qquad \text{(A17)}$$

We note that this equation is the same as the 2 cocycle equation in bosonic systems.

2. 2 + 1D

In 2 + 1D, there are in total 10 admissible branching 2 ↔ 3 moves and 5 admissible branching 1 ↔ 4 moves. Let us show all theses moves will lead to the same fermionic 3 cocycle condition.

For example, the admissible 2 ↔ 3 in Fig. 6 represents the following equation for $\mathcal{V}_3^\pm$:

$$\begin{aligned}
&\int \mathrm{d}\theta_{(123)}^{n(g_1,g_2,g_3)}\mathrm{d}\bar\theta_{(123)}^{n(g_1,g_2,g_3)}\mathcal{V}_3^+(g_0,g_1,g_2,g_3)\mathcal{V}_3^+(g_1,g_2,g_3,g_4)\\
&=\int \mathrm{d}\theta_{(014)}^{n(g_0,g_1,g_4)}\mathrm{d}\bar\theta_{(014)}^{n(g_0,g_1,g_4)}\mathrm{d}\theta_{(024)}^{n(g_0,g_2,g_4)}\mathrm{d}\bar\theta_{(024)}^{n(g_0,g_2,g_4)}\mathrm{d}\theta_{(034)}^{n(g_0,g_3,g_4)}\mathrm{d}\bar\theta_{(034)}^{n(g_0,g_3,g_4)}(-)^{m_1(g_0,g_4)}\\
&\times\mathcal{V}_3^+(g_0,g_1,g_2,g_4)\mathcal{V}_3^+(g_0,g_2,g_3,g_4)\mathcal{V}_3^-(g_0,g_1,g_3,g_4) \qquad \text{(A18)}
\end{aligned}$$

Note that in the above expression, we put the sign factor $(-)^{m_1(g_0,g_4)}$ on the interior link and integral out the Grassmann variables on the interior faces. To simplify the representation, we can formally rewrite the above equation as (see eqn. (18)):

$$\begin{aligned}&\int \mathcal{V}_3^{+}(g_0,g_1,g_2,g_3)\mathcal{V}_3^{+}(g_1,g_2,g_3,g_4) \\ =&\int \mathcal{V}_3^{+}(g_0,g_1,g_2,g_4)\mathcal{V}_3^{+}(g_0,g_2,g_3,g_4)\mathcal{V}_3^{-}(g_0,g_1,g_3,g_4)\end{aligned} \tag{A19}$$

In such a way, we can formally write down all admissible branching 2 ↔ 3 moves in terms of $\mathcal{V}_3^{\pm}$(up to the orientation conjugate $+\to -; -\to +$):

$$\begin{aligned}&\int \mathcal{V}_3^{-}(g_0,g_1,g_2,g_4)\mathcal{V}_3^{-}(g_0,g_2,g_3,g_4) \\ =&\int \mathcal{V}_3^{-}(g_0,g_1,g_2,g_3)\mathcal{V}_3^{-}(g_0,g_1,g_3,g_4)\mathcal{V}_3^{-}(g_1,g_2,g_3,g_4)\end{aligned} \tag{A20}$$

$$\begin{aligned}&\int \mathcal{V}_3^{-}(g_0,g_1,g_2,g_4)\mathcal{V}_3^{+}(g_1,g_2,g_3,g_4) \\ =&\int \mathcal{V}_3^{-}(g_0,g_1,g_2,g_3)\mathcal{V}_3^{-}(g_0,g_1,g_3,g_4)\mathcal{V}_3^{+}(g_0,g_2,g_3,g_4)\end{aligned} \tag{A21}$$

$$\begin{aligned}&\int \mathcal{V}_3^{-}(g_0,g_1,g_2,g_3)\mathcal{V}_3^{+}(g_0,g_2,g_3,g_4) \\ =&\int \mathcal{V}_3^{-}(g_0,g_1,g_2,g_4)\mathcal{V}_3^{+}(g_0,g_1,g_3,g_4)\mathcal{V}_3^{+}(g_1,g_2,g_3,g_4)\end{aligned} \tag{A22}$$

$$\begin{aligned}&\int \mathcal{V}_3^{+}(g_0,g_1,g_3,g_4)\mathcal{V}_3^{+}(g_1,g_2,g_3,g_4) \\ =&\int \mathcal{V}_3^{-}(g_0,g_1,g_2,g_3)\mathcal{V}_3^{+}(g_0,g_1,g_2,g_4)\mathcal{V}_3^{+}(g_0,g_2,g_3,g_4)\end{aligned} \tag{A23}$$

$$\begin{aligned}&\int \mathcal{V}_3^{+}(g_0,g_1,g_2,g_3)\mathcal{V}_3^{+}(g_0,g_1,g_3,g_4) \\ =&\int \mathcal{V}_3^{+}(g_0,g_1,g_2,g_4)\mathcal{V}_3^{+}(g_0,g_2,g_3,g_4)\mathcal{V}_3^{-}(g_1,g_2,g_3,g_4)\end{aligned} \tag{A24}$$

$$\begin{aligned}&\int \mathcal{V}_3^{-}(g_0,g_1,g_2,g_4)\mathcal{V}_3^{+}(g_0,g_1,g_3,g_4) \\ =&\int \mathcal{V}_3^{-}(g_0,g_1,g_2,g_3)\mathcal{V}_3^{+}(g_0,g_2,g_3,g_4)\mathcal{V}_3^{-}(g_1,g_2,g_3,g_4)\end{aligned} \tag{A25}$$

$$\begin{aligned}&\int \mathcal{V}_3^{-}(g_0,g_1,g_3,g_4)\mathcal{V}_3^{+}(g_0,g_2,g_3,g_4) \\ =&\int \mathcal{V}_3^{+}(g_0,g_1,g_2,g_3)\mathcal{V}_3^{-}(g_0,g_1,g_2,g_4)\mathcal{V}_3^{+}(g_1,g_2,g_3,g_4)\end{aligned} \tag{A26}$$

$$\begin{aligned}&\int \mathcal{V}_3^{+}(g_0,g_1,g_2,g_3)\mathcal{V}_3^{+}(g_1,g_2,g_3,g_4) \\ =&\int \mathcal{V}_3^{+}(g_0,g_1,g_2,g_4)\mathcal{V}_3^{+}(g_0,g_2,g_3,g_4)\mathcal{V}_3^{-}(g_0,g_1,g_3,g_4)\end{aligned} \tag{A27}$$

and

$$\begin{aligned}&\int \mathcal{V}_3^{+}(g_0,g_2,g_3,g_4)\mathcal{V}_3^{-}(g_1,g_2,g_3,g_4) \\ =&\int \mathcal{V}_3^{+}(g_0,g_1,g_2,g_3)\mathcal{V}_3^{-}(g_0,g_1,g_2,g_4))\mathcal{V}_3^{+}(g_0,g_1,g_3,g_4)\end{aligned} \tag{A28}$$

$$\begin{aligned}&\int \mathcal{V}_3^{-}(g_0,g_1,g_2,g_3)\mathcal{V}_3^{+}(g_0,g_1,g_2,g_4) \\ =&\int \mathcal{V}_3^{-}(g_0,g_2,g_3,g_4)\mathcal{V}_3^{+}(g_0,g_1,g_3,g_4)\mathcal{V}_3^{+}(g_1,g_2,g_3,g_4)\end{aligned} \tag{A29}$$

Similarly, we can formally write down all the admissible branching 1 ↔ 4 moves in terms of $\mathcal{V}_3^{\pm}$(up to the orientation conjugate $+\to -; -\to +$):

$$\begin{aligned}\mathcal{V}_3^{+}(g_0,g_1,g_2,g_4) =& \int \mathcal{V}_3^{+}(g_0,g_1,g_2,g_3)\mathcal{V}_3^{+}(g_0,g_1,g_3,g_4) \\ &\times \mathcal{V}_3^{-}(g_0,g_2,g_3,g_4)\mathcal{V}_3^{+}(g_1,g_2,g_3,g_4)\end{aligned} \tag{A30}$$

$$\begin{aligned}\mathcal{V}_3^{+}(g_0,g_2,g_3,g_4) =& \int \mathcal{V}_3^{+}(g_0,g_1,g_2,g_3)\mathcal{V}_3^{-}(g_0,g_1,g_2,g_4) \\ &\times \mathcal{V}_3^{+}(g_0,g_1,g_3,g_4)\mathcal{V}_3^{+}(g_1,g_2,g_3,g_4)\end{aligned} \tag{A31}$$

$$\begin{aligned}\mathcal{V}_3^{+}(g_0,g_1,g_3,g_4) =& \int \mathcal{V}_3^{-}(g_0,g_1,g_2,g_3)\mathcal{V}_3^{+}(g_0,g_1,g_2,g_4) \\ &\times \mathcal{V}_3^{+}(g_0,g_2,g_3,g_4)\mathcal{V}_3^{-}(g_1,g_2,g_3,g_4)\end{aligned} \tag{A32}$$

and

$$\begin{aligned}\mathcal{V}_3^{+}(g_1,g_2,g_3,g_4) =& \int \mathcal{V}_3^{-}(g_0,g_1,g_2,g_3)\mathcal{V}_3^{+}(g_0,g_1,g_2,g_4) \\ &\times \mathcal{V}_3^{-}(g_0,g_1,g_3,g_4)\mathcal{V}_3^{+}(g_0,g_2,g_3,g_4)\end{aligned} \tag{A33}$$

$$\begin{aligned}\mathcal{V}_3^{+}(g_0,g_1,g_2,g_3) =& \int \mathcal{V}_3^{+}(g_0,g_1,g_2,g_4)\mathcal{V}_3^{-}(g_0,g_1,g_3,g_4) \\ &\times \mathcal{V}_3^{+}(g_0,g_2,g_3,g_4)\mathcal{V}_3^{-}(g_1,g_2,g_3,g_4)\end{aligned} \tag{A34}$$

Here the symbol $\int$ means that we put the sign factors $(-)^{m_1(g_i,g_j)}$ on all four interior links and integrate over the Grassmann variables on all six interior faces. Note that the first 8 2 ↔ 3 moves(Fig. 14) and first 3 1 ↔ 4 can be induced by a global time ordering while the rest can not. Again here g^0 is defined on the vertex with no incoming edge, g^1 with one incoming edge, etc.

Expressing $\mathcal{V}_3^{\pm}$ in terms of $\nu_3^{\pm}$ and integrating out all the Grassmann variables, the admissible branching 2 ↔ 3 moves lead to the following equations:

$$\begin{aligned}&\nu_3^{-}(g_0,g_1,g_2,g_4)\nu_3^{-}(g_0,g_2,g_3,g_4) \\ =&(-)^{n_2(g_0,g_1,g_2)n_2(g_2,g_3,g_4)+m_1(g_1,g_3)} \\ &\times\nu_3^{-}(g_0,g_1,g_2,g_3)\nu_3^{-}(g_0,g_1,g_3,g_4)\nu_3^{-}(g_1,g_2,g_3,g_4)\end{aligned} \tag{A35}$$



FIG. 14: (Color online) The admissible branching 2 ↔ 3 moves that can be induced by a global time ordering.

$$\nu_3^-(g_0,g_1,g_2,g_4)\nu_3^+(g_1,g_2,g_3,g_4) \tag{A36}$$
$$=(-)^{n_2(g_0,g_1,g_2)n_2(g_2,g_3,g_4)+m_1(g_0,g_3)}$$
$$\times\nu_3^-(g_0,g_1,g_2,g_3)\nu_3^-(g_0,g_1,g_3,g_4)\nu_3^+(g_0,g_2,g_3,g_4)$$



FIG. 15: (Color online) The admissible branching 2 ↔ 3 moves that can not be induced by a global time ordering.



FIG. 16: (Color online) The admissible branching 1 ↔ 4 moves that (a) can be induced by a global time ordering and (b) can not be induced by a global time ordering

$$\nu_3^-(g_0,g_1,g_2,g_3)\nu_3^+(g_0,g_2,g_3,g_4) \tag{A37}$$
$$=(-)^{n_2(g_0,g_1,g_2)n_2(g_2,g_3,g_4)+m_1(g_1,g_4)}$$
$$\times\nu_3^-(g_0,g_1,g_2,g_4)\nu_3^+(g_0,g_1,g_3,g_4)\nu_3^+(g_1,g_2,g_3,g_4)$$

$$\begin{aligned}&\nu_3^+(g_0,g_1,g_3,g_4)\nu_3^+(g_1,g_2,g_3,g_4)\\ =&(-)^{n_2(g_0,g_1,g_2)n_2(g_2,g_3,g_4)+m_1(g_0,g_2)}\\ &\times\nu_3^-(g_0,g_1,g_2,g_3)\nu_3^+(g_0,g_1,g_2,g_4)\nu_3^+(g_0,g_2,g_3,g_4)\end{aligned}\tag{A38}$$

$$\begin{aligned}&\nu_3^+(g_0,g_1,g_2,g_3)\nu_3^+(g_0,g_1,g_3,g_4)\\ =&(-)^{n_2(g_0,g_1,g_2)n_2(g_2,g_3,g_4)+m_1(g_2,g_4)}\\ &\times\nu_3^+(g_0,g_1,g_2,g_4)\nu_3^+(g_0,g_2,g_3,g_4)\nu_3^-(g_1,g_2,g_3,g_4)\end{aligned}\tag{A39}$$

$$\begin{aligned}&\nu_3^-(g_0,g_1,g_2,g_4)\nu_3^+(g_0,g_1,g_3,g_4)\\ =&(-)^{n_2(g_0,g_1,g_2)n_2(g_2,g_3,g_4)+n_2(g_1,g_2,g_3)+n_2(g_1,g_2,g_4)+m_1(g_2,g_3)}\\ &\times\nu_3^-(g_0,g_1,g_2,g_3)\nu_3^+(g_0,g_2,g_3,g_4)\nu_3^-(g_1,g_2,g_3,g_4)\end{aligned}\tag{A40}$$

$$\begin{aligned}&\nu_3^-(g_0,g_1,g_3,g_4)\nu_3^+(g_0,g_2,g_3,g_4)\\ =&(-)^{n_2(g_0,g_1,g_2)n_2(g_2,g_3,g_4)+n_2(g_0,g_1,g_2)+n_2(g_0,g_1,g_3)+m_1(g_1,g_2)}\\ &\times\nu_3^+(g_0,g_1,g_2,g_3)\nu_3^-(g_0,g_1,g_2,g_4)\nu_3^+(g_1,g_2,g_3,g_4)\end{aligned}\tag{A41}$$

$$\begin{aligned}&\nu_3^+(g_0,g_1,g_2,g_3)\nu_3^+(g_1,g_2,g_3,g_4)\\ =&(-)^{n_2(g_0,g_1,g_2)n_2(g_2,g_3,g_4)+n_2(g_0,g_3,g_4)+n_2(g_1,g_3,g_4)+m_1(g_0,g_4)}\\ &\times\nu_3^+(g_0,g_1,g_2,g_4)\nu_3^+(g_0,g_2,g_3,g_4)\nu_3^-(g_0,g_1,g_3,g_4)\end{aligned}\tag{A42}$$

and

$$\begin{aligned}&\nu_3^+(g_0,g_2,g_3,g_4)\nu_3^-(g_1,g_2,g_3,g_4)\\ =&(-)^{n_2(g_0,g_1,g_2)n_2(g_2,g_3,g_4)+n_2(g_0,g_1,g_2)+n_2(g_1,g_2,g_4)+m_1(g_0,g_1)}\\ &\times\nu_3^+(g_0,g_1,g_2,g_3)\nu_3^-(g_0,g_1,g_2,g_4))\nu_3^+(g_0,g_1,g_3,g_4)\end{aligned}\tag{A43}$$

$$\begin{aligned}&\nu_3^-(g_0,g_1,g_2,g_3)\nu_3^+(g_0,g_1,g_2,g_4)\\ =&(-)^{n_2(g_0,g_1,g_2)n_2(g_2,g_3,g_4)+n_2(g_0,g_2,g_4)+n_2(g_0,g_3,g_4)+m_1(g_3,g_4)}\\ &\times\nu_3^-(g_0,g_2,g_3,g_4)\nu_3^+(g_0,g_1,g_3,g_4)\nu_3^+(g_1,g_2,g_3,g_4)\end{aligned}\tag{A44}$$

We note that the sign factor $(-)^{m_1(g_i,g_j)}$ comes from the definition of the fermionic path integral. Later we will see such a factor is very important to make all the branching moves to be self consistent.

Similarly, all the admissible branching $1 \leftrightarrow 4$ moves lead to the following equations for $\nu_3^\pm$:

$$\begin{aligned}&\nu_3^+(g_0,g_1,g_2,g_4)\\ =&(-)^{n_2(g_0,g_1,g_2)n_2(g_2,g_3,g_4)+n_2(g_2,g_3,g_4)}\\ &\times(-)^{m_1(g_0,g_3)+m_1(g_1,g_3)+m_1(g_2,g_3)+m_1(g_3,g_4)}\nu_3^+(g_0,g_1,g_2,g_3)\\ &\times\nu_3^+(g_0,g_1,g_3,g_4)\nu_3^-(g_0,g_2,g_3,g_4)\nu_3^+(g_1,g_2,g_3,g_4)\end{aligned}\tag{A45}$$

$$\begin{aligned}&\nu_3^+(g_0,g_2,g_3,g_4)\\ =&(-)^{n_2(g_0,g_1,g_2)n_2(g_2,g_3,g_4)+n_2(g_0,g_1,g_2)}\\ &\times(-)^{m_1(g_0,g_1)+m_1(g_1,g_2)+m_1(g_1,g_3)+m_1(g_1,g_4)}\nu_3^+(g_0,g_1,g_2,g_3)\\ &\times\nu_3^-(g_0,g_1,g_2,g_4)\nu_3^+(g_0,g_1,g_3,g_4)\nu_3^+(g_1,g_2,g_3,g_4)\end{aligned}\tag{A46}$$

$$\begin{aligned}&\nu_3^+(g_0,g_1,g_3,g_4)\\ =&(-)^{n_2(g_0,g_1,g_2)n_2(g_2,g_3,g_4)+n_2(g_1,g_2,g_3)}\\ &\times(-)^{m_1(g_0,g_2)+m_1(g_1,g_2)+m_1(g_2,g_3)+m_1(g_2,g_4)}\nu_3^-(g_0,g_1,g_2,g_3)\\ &\times\nu_3^+(g_0,g_1,g_2,g_4)\nu_3^+(g_0,g_2,g_3,g_4)\nu_3^-(g_1,g_2,g_3,g_4)\end{aligned}\tag{A47}$$

and

$$\begin{aligned}&\nu_3^+(g_1,g_2,g_3,g_4)\\ =&(-)^{n_2(g_0,g_1,g_2)n_2(g_2,g_3,g_4)+n_2(g_0,g_1,g_4)}\\ &\times(-)^{m_1(g_0,g_1)+m_1(g_0,g_2)+m_1(g_0,g_3)+m_1(g_0,g_4)}\nu_3^-(g_0,g_1,g_2,g_3)\\ &\times\nu_3^+(g_0,g_1,g_2,g_4)\nu_3^-(g_0,g_1,g_3,g_4)\nu_3^+(g_0,g_2,g_3,g_4)\end{aligned}\tag{A48}$$

$$\begin{aligned}&\nu_3^+(g_0,g_1,g_2,g_3)\\ =&(-)^{n_2(g_0,g_1,g_2)n_2(g_2,g_3,g_4)+n_2(g_0,g_3,g_4)}\\ &\times(-)^{m_1(g_0,g_4)+m_1(g_1,g_4)+m_1(g_2,g_4)+m_1(g_3,g_4)}\nu_3^+(g_0,g_1,g_2,g_4)\\ &\times\nu_3^-(g_0,g_1,g_3,g_4)\nu_3^+(g_0,g_2,g_3,g_4)\nu_3^-(g_1,g_2,g_3,g_4)\end{aligned}\tag{A49}$$

Amazingly, if we define:

$$\begin{aligned}\nu_3^+(g_0,g_1,g_2,g_3)&=(-)^{m_1(g_0,g_2)}\nu_3(g_0,g_1,g_2,g_3),\\ \nu_3^-(g_0,g_1,g_2,g_3)&=(-)^{m_1(g_1,g_3)}/\nu_3(g_0,g_1,g_2,g_3)\end{aligned}\tag{A50}$$

we find all the above equations are equivalent to the following single equation, which is the fermionic 3 cocycle condition of ν_3.

$$\begin{aligned}&\nu_3(g_1,g_2,g_3,g_4)\nu_3(g_0,g_1,g_3,g_4)\nu_3(g_0,g_1,g_2,g_3)=\\ &(-)^{n_2(g_0,g_1,g_2)n_2(g_2,g_3,g_4)}\nu_3(g_0,g_2,g_3,g_4)\nu_3(g_0,g_1,g_2,g_4)\end{aligned}\tag{A51}$$

3. 3 + 1D

The admissible branching moves for 3+1D case is much more complicated. There are in total 10 $3 \leftrightarrow 3$ moves, 15 $2 \leftrightarrow 4$ moves and 6 $1 \leftrightarrow 5$ moves.

For example, Fig. 8 represents one admissible branching $2 \leftrightarrow 4$ move, which leads to the following equation for $\mathcal{V}_4^\pm$:

$$\begin{aligned}
&\int \mathrm{d}\theta_{(1234)}^{n(g_1,g_2,g_3,g_4)} \mathrm{d}\bar{\theta}_{(1234)}^{n(g_1,g_2,g_3,g_4)} \mathcal{V}_4^+(g_1,g_2,g_3,g_4,g_5)\mathcal{V}_4^-(g_0,g_1,g_2,g_3,g_4)\\
=&\int \mathrm{d}\theta_{(0125)}^{n(g_0,g_1,g_2,g_5)} \mathrm{d}\bar{\theta}_{(0125)}^{n(g_0,g_1,g_2,g_5)} \mathrm{d}\theta_{(0135)}^{n(g_0,g_1,g_3,g_5)} \mathrm{d}\bar{\theta}_{(0135)}^{n(g_0,g_1,g_3,g_5)} \mathrm{d}\theta_{(0145)}^{n(g_0,g_1,g_4,g_5)} \mathrm{d}\bar{\theta}_{(0145)}^{n(g_0,g_1,g_4,g_5)} \mathrm{d}\theta_{(0235)}^{n(g_0,g_2,g_3,g_5)} \mathrm{d}\bar{\theta}_{(0235)}^{n(g_0,g_2,g_3,g_5)}\\
&\times \mathrm{d}\theta_{(0245)}^{n(g_0,g_2,g_4,g_5)} \mathrm{d}\bar{\theta}_{(0245)}^{n(g_0,g_2,g_4,g_5)} \mathrm{d}\theta_{(0345)}^{n(g_0,g_3,g_4,g_5)} \mathrm{d}\bar{\theta}_{(0345)}^{n(g_0,g_3,g_4,g_5)} (-)^{m_2(g_0,g_1,g_5)+m_2(g_0,g_2,g_5)+m_2(g_0,g_3,g_5)+m_2(g_0,g_4,g_5)}\\
&\times \mathcal{V}_4^+(g_0,g_2,g_3,g_4,g_5)\mathcal{V}_4^+(g_0,g_1,g_2,g_4,g_5)\mathcal{V}_4^-(g_0,g_1,g_3,g_4,g_5)\mathcal{V}_4^-(g_0,g_1,g_2,g_3,g_5)
\end{aligned} \tag{A52}$$

Here we integrate out the Grassmann variables on the interior tetrahedra and put the sign factor on the interior surfaces. We can formally denote the above equation as (see eqn. (18)):

$$\begin{aligned}
&\int \mathcal{V}_4^+(g_1,g_2,g_3,g_4,g_5)\mathcal{V}_4^-(g_0,g_1,g_2,g_3,g_4)\\
=&\int \mathcal{V}_4^+(g_0,g_2,g_3,g_4,g_5)\mathcal{V}_4^+(g_0,g_1,g_2,g_4,g_5)\mathcal{V}_4^-(g_0,g_1,g_3,g_4,g_5)\mathcal{V}_4^-(g_0,g_1,g_2,g_3,g_5)
\end{aligned} \tag{A53}$$

Similarly, Fig. 9 represents one admissible branching 1 ↔ 5 move and can be formally written down in terms of $\mathcal{V}_4^\pm$ as:

$$\begin{aligned}
&\mathcal{V}_4^-(g_0,g_1,g_2,g_3,g_4)\\
=&\int \mathcal{V}_4^+(g_0,g_2,g_3,g_4,g_5)\mathcal{V}_4^+(g_0,g_1,g_2,g_4,g_5)\mathcal{V}_4^-(g_0,g_1,g_3,g_4,g_5)\mathcal{V}_4^-(g_0,g_1,g_2,g_3,g_5)\mathcal{V}_4^-(g_1,g_2,g_3,g_4,g_5)
\end{aligned} \tag{A54}$$

We note that here the symbol $\int$ means integrating over all Grassmann variables on 10 interior tetrahedra and put the sign factor $(-)^{m_2(g_i,g_j,g_k)}$ on 10 interior surfaces.

In addition, there are 3 ↔ 3 moves. For example, one of such moves gives rise the following formal equation for $\mathcal{V}_4^\pm$:

$$\begin{aligned}
&\int \mathcal{V}_4^+(g_1,g_2,g_3,g_4,g_5)\mathcal{V}_4^+(g_0,g_1,g_3,g_4,g_5)\mathcal{V}_4^+(g_0,g_1,g_2,g_3,g_5)\\
=&\int \mathcal{V}_4^+(g_0,g_1,g_2,g_3,g_4)\mathcal{V}_4^+(g_0,g_2,g_3,g_4,g_5)\mathcal{V}_4^+(g_0,g_1,g_2,g_4,g_5)
\end{aligned} \tag{A55}$$

In this case, the symbol $\int$ on both sides mean integrating over all Grassmann variables on 3 interior tetrahedra and put the sign factor $(-)^{m_2(g_i,g_j,g_k)}$ on 1 interior surface.

After integral over all the Grassmann variables, we can express the above three equations in terms of $\nu_4^\pm$:

$$\begin{aligned}
&\nu_4^+(g_1,g_2,g_3,g_4,g_5)\nu_4^-(g_0,g_1,g_2,g_3,g_4)\\
=&(-)^{n_3(g_0,g_1,g_2,g_3)n_3(g_0,g_3,g_4,g_5)+n_3(g_1,g_2,g_3,g_4)n_3(g_0,g_1,g_4,g_5)+n_3(g_2,g_3,g_4,g_5)n_3(g_0,g_1,g_2,g_5)+n_3(g_0,g_1,g_4,g_5)}\\
&\quad\times (-)^{m_2(g_0,g_1,g_5)+m_2(g_0,g_2,g_5)+m_2(g_0,g_3,g_5)+m_2(g_0,g_4,g_5)}\\
&\quad\times \nu_4^+(g_0,g_2,g_3,g_4,g_5)\nu_4^+(g_0,g_1,g_2,g_4,g_5)\nu_4^-(g_0,g_1,g_3,g_4,g_5)\nu_4^-(g_0,g_1,g_2,g_3,g_5),
\end{aligned} \tag{A56}$$

$$\begin{aligned}
&\nu_4^-(g_0,g_1,g_2,g_3,g_4)\\
=&(-)^{n_3(g_0,g_1,g_2,g_3)n_3(g_0,g_3,g_4,g_5)+n_3(g_1,g_2,g_3,g_4)n_3(g_0,g_1,g_4,g_5)+n_3(g_2,g_3,g_4,g_5)n_3(g_0,g_1,g_2,g_5)+n_3(g_0,g_1,g_4,g_5)+n_3(g_1,g_2,g_4,g_5)+n_3(g_2,g_3,g_4,g_5)}\\
\times&(-)^{m_2(g_0,g_1,g_5)+m_2(g_0,g_2,g_5)+m_2(g_0,g_3,g_5)+m_2(g_0,g_4,g_5)+m_2(g_1,g_2,g_5)+m_2(g_1,g_3,g_5)+m_2(g_1,g_4,g_5)+m_2(g_2,g_3,g_5)+m_2(g_2,g_4,g_5)+m_2(g_3,g_4,g_5)}\\
&\quad\times \nu_4^+(g_0,g_2,g_3,g_4,g_5)\nu_4^+(g_0,g_1,g_2,g_4,g_5)\nu_4^-(g_0,g_1,g_3,g_4,g_5)\nu_4^-(g_0,g_1,g_2,g_3,g_5)\nu_4^-(g_1,g_2,g_3,g_4,g_5),
\end{aligned} \tag{A57}$$

$$\begin{aligned}
&\nu_4^+(g_1,g_2,g_3,g_4,g_5)\nu_4^+(g_0,g_1,g_3,g_4,g_5)\nu_4^+(g_0,g_1,g_2,g_3,g_5)(-)^{m_2(g_1,g_3,g_5)}\\
=&(-)^{n_3(g_0,g_1,g_2,g_3)n_3(g_0,g_3,g_4,g_5)+n_3(g_1,g_2,g_3,g_4)n_3(g_0,g_1,g_4,g_5)+n_3(g_2,g_3,g_4,g_5)n_3(g_0,g_1,g_2,g_5)+m_2(g_0,g_2,g_4)}\\
&\quad\times \nu_4^+(g_0,g_1,g_2,g_3,g_4)\nu_4^+(g_0,g_2,g_3,g_4,g_5)\nu_4^+(g_0,g_1,g_2,g_4,g_5).
\end{aligned} \tag{A58}$$

Surprisingly, if we define $\nu_4^\pm$ as:

$$\begin{aligned}
\nu_4^+(g_0,g_1,g_2,g_3,g_4) =&(-)^{m_2(g_0,g_1,g_3)+m_2(g_1,g_3,g_4)+m_2(g_1,g_2,g_3)}\nu_4(g_0,g_1,g_2,g_3,g_4),\\
\nu_4^-(g_0,g_1,g_2,g_3,g_4) =&(-)^{m_2(g_0,g_2,g_4)}/\nu_4(g_0,g_1,g_2,g_3,g_4),
\end{aligned} \tag{A59}$$

we find all the above three equations will be equivalent to the following single equation of ν_4, which is the fermionic 4 cocycle condition:

$$\begin{aligned}
&\nu_4(g_1,g_2,g_3,g_4,g_5)\nu_4(g_0,g_1,g_3,g_4,g_5)\nu_4(g_0,g_1,g_2,g_3,g_5)\\
=&(-)^{n_3(g_0,g_1,g_2,g_3)n_3(g_0,g_3,g_4,g_5)+n_3(g_1,g_2,g_3,g_4)n_3(g_0,g_1,g_4,g_5)+n_3(g_2,g_3,g_4,g_5)n_3(g_0,g_1,g_2,g_5)}\times\\
&\nu_4(g_0,g_2,g_3,g_4,g_5)\nu_4(g_0,g_1,g_2,g_4,g_5)\nu_4(g_0,g_1,g_2,g_3,g_4).
\end{aligned} \tag{A60}$$

Indeed, it can be verified by computer that all other admissible 3 ↔ 3, 2 ↔ 4 and 1 ↔ 5 branching moves will give rise to the same fermionic 4 cocycle condition! Again, the phase factor $(-)^{m_2(g_i,g_j,g_k)}$ in the fermionic path integral is crucial for the self consistency of all admissible branching moves.

Appendix B: The fixed-point action on a closed complex and the unitary condition

In the following we will show that $\int \mathcal{V}_d^+\mathcal{V}_d^-$ is always equal to 1.

In $1+1$D, we have:

$$\begin{aligned}
&\int \mathcal{V}_2^+(g_0,g_1,g_2)\mathcal{V}_2^-(g_0,g_1,g_2)\\
=&\int \mathrm{d}\theta_{(01)}^{n_1(g_0,g_1)}\mathrm{d}\bar\theta_{(01)}^{n_1(g_0,g_1)}\mathrm{d}\theta_{(02)}^{n_1(g_0,g_2)}\mathrm{d}\bar\theta_{(02)}^{n_1(g_0,g_2)}\mathrm{d}\theta_{(12)}^{n_1(g_1,g_2)}\mathrm{d}\bar\theta_{(12)}^{n_1(g_1,g_2)}(-)^{m_0(g_0)+m_0(g_1)+m_0(g_2)}\\
&\times \nu_2^+(g_0,g_1,g_2)\theta_{(12)}^{n_1(g_1,g_2)}\theta_{(01)}^{n_1(g_0,g_1)}\bar\theta_{(02)}^{n_1(g_0,g_2)}\nu_2^-(g_0,g_1,g_2)\theta_{(02)}^{n_1(g_0,g_2)}\bar\theta_{(01)}^{n_1(g_0,g_1)}\bar\theta_{(12)}^{n_1(g_1,g_2)}\\
=&\int \mathrm{d}\theta_{(01)}^{n_1(g_0,g_1)}\mathrm{d}\bar\theta_{(01)}^{n_1(g_0,g_1)}\mathrm{d}\theta_{(02)}^{n_1(g_0,g_2)}\mathrm{d}\bar\theta_{(02)}^{n_1(g_0,g_2)}\mathrm{d}\theta_{(12)}^{n_1(g_1,g_2)}\mathrm{d}\bar\theta_{(12)}^{n_1(g_1,g_2)}(-)^{n_1(g_0,g_2)}\\
&\times \theta_{(12)}^{n_1(g_1,g_2)}\theta_{(01)}^{n_1(g_0,g_1)}\bar\theta_{(02)}^{n_1(g_0,g_2)}\theta_{(02)}^{n_1(g_0,g_2)}\bar\theta_{(01)}^{n_1(g_0,g_1)}\bar\theta_{(12)}^{n_1(g_1,g_2)}\\
=&1
\end{aligned} \tag{B1}$$

In the above calculation, we can move the pair $\mathrm{d}\theta_{(01)}^{n_1(g_0,g_1)}\mathrm{d}\bar\theta_{(01)}^{n_1(g_0,g_1)}$ to the front of $\bar\theta_{(02)}^{n_1(g_0,g_2)}\theta_{(02)}^{n_1(g_0,g_2)}$ without generating any signs. We then can evaluate $\int \mathrm{d}\theta_{(02)}^{n_1(g_0,g_1)}\mathrm{d}\bar\theta_{(02)}^{n_1(g_0,g_1)}\bar\theta_{(02)}^{n_1(g_0,g_2)}\theta_{(02)}^{n_1(g_0,g_2)} = 1$. We next move $\mathrm{d}\theta_{(01)}^{n_1(g_0,g_1)}\mathrm{d}\bar\theta_{(01)}^{n_1(g_0,g_1)}$ to the front of $\theta_{(01)}^{n_1(g_0,g_1)}\bar\theta_{(01)}^{n_1(g_0,g_1)}$ without generating any signs. We then evaluate $\int \mathrm{d}\theta_{(01)}^{n_1(g_0,g_1)}\mathrm{d}\bar\theta_{(01)}^{n_1(g_0,g_1)}\theta_{(01)}^{n_1(g_0,g_1)}\bar\theta_{(01)}^{n_1(g_0,g_1)} = (-)^{n_1(g_0,g_1)}$, which has a sign factor. The Grassmann integral $\int \mathrm{d}\theta_{(12)}^{n_1(g_1,g_2)}\mathrm{d}\bar\theta_{(12)}^{n_1(g_1,g_2)}\theta_{(12)}^{n_1(g_1,g_2)}\bar\theta_{(12)}^{n_1(g_1,g_2)} = (-)^{n_1(g_1,g_2)}$ also generate a sign factor. Such two sign factors $(-)^{n_1(g_0,g_1)}(-)^{n_1(g_1,g_2)}$ cancel the sign factor $(-)^{n_1(g_0,g_2)}$ due to the condition eqn. (10) on n_d.

Similarly, in $(2+1)$D and in $(3+1)$D, we find that the sign factors in the integration measure (18) and in the $\nu_d^\pm$ expressions (35) or (35), when combined with those from exchanging the Grassmann numbers, just cancel each other:

$$\begin{aligned}
&\int \mathcal{V}_3^+(g_0,g_1,g_2,g_3)\mathcal{V}_3^-(g_0,g_1,g_2,g_3)\\
=&\int \mathrm{d}\theta_{(012)}^{n_2(g_0,g_1,g_2)}\mathrm{d}\bar\theta_{(012)}^{n_2(g_0,g_1,g_2)}\mathrm{d}\theta_{(013)}^{n_2(g_0,g_1,g_3)}\mathrm{d}\bar\theta_{(013)}^{n_2(g_0,g_1,g_3)}\mathrm{d}\theta_{(023)}^{n_2(g_0,g_2,g_3)}\mathrm{d}\bar\theta_{(023)}^{n_2(g_0,g_2,g_3)}\mathrm{d}\theta_{(123)}^{n_2(g_1,g_2,g_3)}\mathrm{d}\bar\theta_{(123)}^{n_2(g_1,g_2,g_3)}\\
&\times (-)^{m_1(g_0,g_1)+m_1(g_0,g_2)+m_1(g_0,g_3)+m_1(g_1,g_2)+m_1(g_1,g_3)+m_1(g_2,g_3)}\\
&\times \nu_3^+(g_0,g_1,g_2,g_3)\theta_{(123)}^{n_2(g_1,g_2,g_3)}\theta_{(013)}^{n_2(g_0,g_1,g_3)}\bar\theta_{(023)}^{n_2(g_0,g_2,g_3)}\bar\theta_{(012)}^{n_2(g_0,g_1,g_2)}\\
&\times \nu_3^-(g_0,g_1,g_2,g_3)\theta_{(012)}^{n_2(g_0,g_1,g_2)}\theta_{(023)}^{n_2(g_0,g_2,g_3)}\bar\theta_{(013)}^{n_2(g_0,g_1,g_3)}\bar\theta_{(123)}^{n_2(g_1,g_2,g_3)}\\
=&\int \mathrm{d}\theta_{(012)}^{n_2(g_0,g_1,g_2)}\mathrm{d}\bar\theta_{(012)}^{n_2(g_0,g_1,g_2)}\mathrm{d}\theta_{(013)}^{n_2(g_0,g_1,g_3)}\mathrm{d}\bar\theta_{(013)}^{n_2(g_0,g_1,g_3)}\mathrm{d}\theta_{(023)}^{n_2(g_0,g_2,g_3)}\mathrm{d}\bar\theta_{(023)}^{n_2(g_0,g_2,g_3)}\mathrm{d}\theta_{(123)}^{n_2(g_1,g_2,g_3)}\mathrm{d}\bar\theta_{(123)}^{n_2(g_1,g_2,g_3)}\\
&\times (-)^{n_2(g_0,g_1,g_3)+n_2(g_1,g_2,g_3)}\theta_{(123)}^{n_2(g_1,g_2,g_3)}\theta_{(013)}^{n_2(g_0,g_1,g_3)}\bar\theta_{(023)}^{n_2(g_0,g_2,g_3)}\bar\theta_{(012)}^{n_2(g_0,g_1,g_2)}\theta_{(012)}^{n_2(g_0,g_1,g_2)}\theta_{(023)}^{n_2(g_0,g_2,g_3)}\bar\theta_{(013)}^{n_2(g_0,g_1,g_3)}\bar\theta_{(123)}^{n_2(g_1,g_2,g_3)}\\
=&1;
\end{aligned} \tag{B2}$$

$$
\begin{aligned}
&\int \mathcal{V}_4^+(g_0,g_1,g_2,g_3,g_4)\mathcal{V}_4^-(g_0,g_1,g_2,g_3,g_4)\\
=&\int \mathrm{d}\theta_{(0123)}^{n_3(g_0,g_1,g_2,g_3)}\mathrm{d}\bar\theta_{(0123)}^{n_3(g_0,g_1,g_2,g_3)}\mathrm{d}\theta_{(0124)}^{n_3(g_0,g_1,g_2,g_4)}\mathrm{d}\bar\theta_{(0124)}^{n_3(g_0,g_1,g_2,g_4)}\mathrm{d}\theta_{(0134)}^{n_3(g_0,g_1,g_3,g_4)}\mathrm{d}\bar\theta_{(0134)}^{n_3(g_0,g_1,g_3,g_4)}\\
&\times \mathrm{d}\theta_{(0234)}^{n_3(g_0,g_2,g_3,g_4)}\mathrm{d}\bar\theta_{(0234)}^{n_3(g_0,g_2,g_3,g_4)}\mathrm{d}\theta_{(1234)}^{n_3(g_1,g_2,g_3,g_4)}\mathrm{d}\bar\theta_{(1234)}^{n_3(g_1,g_2,g_3,g_4)}(-)^{m_2(g_0,g_1,g_2)+m_2(g_0,g_1,g_3)}\\
&\times (-)^{m_2(g_0,g_1,g_4)+m_2(g_0,g_2,g_3)+m_2(g_0,g_2,g_4)+m_2(g_0,g_3,g_4)+m_2(g_1,g_2,g_3)+m_2(g_1,g_2,g_4)+m_2(g_1,g_3,g_4)+m_2(g_2,g_3,g_4)}\\
&\times \nu_4^+(g_0,g_1,g_2,g_3,g_4)\theta_{(1234)}^{n_3(g_1,g_2,g_3,g_4)}\theta_{(0134)}^{n_3(g_0,g_1,g_3,g_4)}\theta_{(0123)}^{n_3(g_0,g_1,g_2,g_3)}\bar\theta_{(0234)}^{n_3(g_0,g_2,g_3,g_4)}\bar\theta_{(0124)}^{n_3(g_0,g_1,g_2,g_4)}\\
&\times \nu_4^-(g_0,g_1,g_2,g_3,g_4)\theta_{(0124)}^{n_3(g_0,g_1,g_2,g_4)}\theta_{(0234)}^{n_3(g_0,g_2,g_3,g_4)}\bar\theta_{(0123)}^{n_3(g_0,g_1,g_2,g_3)}\bar\theta_{(0134)}^{n_3(g_0,g_1,g_3,g_4)}\bar\theta_{(1234)}^{n_3(g_1,g_2,g_3,g_4)}\\
=&\int \mathrm{d}\theta_{(0123)}^{n_3(g_0,g_1,g_2,g_3)}\mathrm{d}\bar\theta_{(0123)}^{n_3(g_0,g_1,g_2,g_3)}\mathrm{d}\theta_{(0124)}^{n_3(g_0,g_1,g_2,g_4)}\mathrm{d}\bar\theta_{(0124)}^{n_3(g_0,g_1,g_2,g_4)}\mathrm{d}\theta_{(0134)}^{n_3(g_0,g_1,g_3,g_4)}\mathrm{d}\bar\theta_{(0134)}^{n_3(g_0,g_1,g_3,g_4)}\\
&\times \mathrm{d}\theta_{(0234)}^{n_3(g_0,g_2,g_3,g_4)}\mathrm{d}\bar\theta_{(0234)}^{n_3(g_0,g_2,g_3,g_4)}\mathrm{d}\theta_{(1234)}^{n_3(g_1,g_2,g_3,g_4)}\mathrm{d}\bar\theta_{(1234)}^{n_3(g_1,g_2,g_3,g_4)}(-)^{n_3(g_0,g_1,g_2,g_4)+n_3(g_0,g_2,g_3,g_4)}\\
&\times \theta_{(1234)}^{n_3(g_1,g_2,g_3,g_4)}\theta_{(0134)}^{n_3(g_0,g_1,g_3,g_4)}\theta_{(0123)}^{n_3(g_0,g_1,g_2,g_3)}\bar\theta_{(0234)}^{n_3(g_0,g_2,g_3,g_4)}\bar\theta_{(0124)}^{n_3(g_0,g_1,g_2,g_4)}\\
&\times \theta_{(0124)}^{n_3(g_0,g_1,g_2,g_4)}\theta_{(0234)}^{n_3(g_0,g_2,g_3,g_4)}\bar\theta_{(0123)}^{n_3(g_0,g_1,g_2,g_3)}\bar\theta_{(0134)}^{n_3(g_0,g_1,g_3,g_4)}\bar\theta_{(1234)}^{n_3(g_1,g_2,g_3,g_4)}\\
=&1.
\end{aligned}
\tag{B3}
$$

Appendix C: Group super-cohomology

In Section V, we studied the condition on $\nu_d(g_0,g_1,...)$ so that the fermionic path integral constructed from $\nu_d(g_0,g_1,...)$ correspond to a fermionic topological non-linear σ-model which is a fixed point theory under the RG flow. Those conditions on $\nu_d(g_0,g_1,...)$ actually define a generalization of group cohomology.[50,51] We will call such a generalization group super-cohomology. In this section, we will study the group super-cohomology in detail.

A d-cohomology class of group super-cohomology is a set that depend on a full symmetry group G_f and a fermionic G-module M. We will denote the fermionic d-cohomology class as $\mathscr{H}^d(G_f,M)$.

We note that a fermion system always has a Z_2 symmetry which corresponds to the conservation of fermion-number parity. Such a symmetry group is denoted as Z_2^f which is generated by $P_f=(-)^{N_f}$ where N_f is the fermion number. The full symmetry group G_f always contain Z_2^f as a subgroup. We will define $G_b=G_f/Z_2^f$.

1. Graded structure of a group G_b

First, let us introduce the graded structure of a group G_b. A dD-graded structure of a group G_b is an integer function $n_{d-1}(g_1,...,g_d)$ of d variables, whose values are $0,1$. Here we choose g_i to be an element of G_b, rather then an element of the full symmetry group G_f. This is because g_i correspond to the *bosonic* fields in the path integral, and the bosonic field is invariant under Z_2^f.

The function $n_{d-1}(g_1,...,g_d)$ satisfies

$$
\begin{aligned}
&n_{d-1}(gg_1,...,gg_d)=n_{d-1}(g_1,...,g_d), \quad \forall g\in G_b,\\
&\sum_{i=0}^{d} n_{d-1}(g_0,..,g_{i-1},g_{i+1},..,g_d)= \text{ even}, \quad \forall g_0,..,g_d\in G.
\end{aligned}
\tag{C1}
$$

We see that a dD-graded structure of a group G_b is a $(d-1)$-cocycle $n_{d-1}\in\mathcal{Z}^{d-1}(G_b,\mathbb{Z}_2)$. If a function n_{d-1} only satisfies the first condition in eqn. (C1), then it will be called a $(d-1)$-cochain. The space of all $(d-1)$-cochains is denoted as $\mathcal{C}^{d-1}(G_b,\mathbb{Z}_2)$.

The coboundary is given by

$$
\begin{aligned}
\mathcal{B}^{d-1}(G_b,\mathbb{Z}_2)=\Big\{ & n'_{d-1}(g_1,...,g_d)=\sum_i m'_{d-2}(g_1,...,g_{i-1},g_{i+1},...,g_d) \text{ mod } 2 \Big|\\
& m'_{d-2}(gg_1,...,gg_{d-1})=m'_{d-2}(g_1,...,g_{d-1}), \quad \forall g,g_1,...,g_{d-1}\in G_b\Big\}.
\end{aligned}
$$

We say that two graded structures differ by an above coboundary are equivalent. Thus different classes of dD-graded structures are given by $\mathcal{H}^{d-1}(G_b,\mathbb{Z}_2) = \mathcal{Z}^{d-1}(G_b,\mathbb{Z}_2)/\mathcal{B}^{d-1}(G_b,\mathbb{Z}_2)$.

2. G-module

For a group G_f, let M be a G_f-module, which is an Abelian group (with multiplication operation) on which G_f acts compatibly with the multiplication operation (*ie* the Abelian group structure):

$$g\cdot(ab) = (g\cdot a)(g\cdot b), \quad g\in G_f, \quad a,b\in M. \tag{C2}$$

For the most cases studied in this paper, M is simply the $U(1)$ group and a an $U(1)$ phase. The multiplication operation ab is the usual multiplication of the $U(1)$ phases. The group action is trivial: $g\cdot a = a$, $g\in G_f$, $a\in U(1)$. We will denote such a trivial G_f-module as $M = U(1)$.

For a group G_f that contain time-reversal operation, we can define a non-trivial G_f-module which is denoted as $U_T(1)$. $U_T(1)$ is also a $U(1)$ group whose elements are the $U(1)$ phases. The multiplication operation ab, $a,b\in U_T(1)$, is still the usual multiplication of the $U(1)$ phases. However, the group action is non-trivial now: $g\cdot a = a^{s(g)}$, $g\in G_f$, $a\in U_T(1)$. Here $s(g)=1$ if the number of time-reversal operations in the group operation g is even and $s(g)=-1$ if the number of time-reversal operations in g is odd.

3. Fermionic d-cochain

A fermionic d-cochain is described by a set of three functions $\nu_d(g_0,...,g_d)$, $n_{d-1}(g_1,...,g_d)$, and $u^g_{d-1}(g_1,...,g_d)$. $n_{d-1}(g_1,...,g_d)$ is $(d-1)$-cocycle in $\mathcal{Z}^{d-1}(G_b,\mathbb{Z}_2)$ which has been discussed above. $\nu_d(g_0,...,g_d)$ is a function of $1+d$ variables whose value is in a G-module M, $\nu_d : G_b^{1+d}\to M$. Again, note that g_i is an elements of G_b rather then the full symmetry group G_f. For cases studied in this paper, M is always a $U(1)$ group (*ie* $\nu_d(g_0,...,g_d)$ is a complex phase).

Since the action amplitude is invariant under the symmetry transformation in G_f, $\nu_d(g_0,...,g_d)$ must satisfy certain conditions. Note that a transformation g in G_f will generate a transformation in G_b, $gG_b\to G_b$, since $G_b = G_f/Z_2^f$. So $\nu_d(g_0,...,g_d)$ transform as $\nu_d(g_0,...,g_d)\to\nu_d^{s(g)}(gg_0,...,gg_d)$.

The invariance under the symmetry transformation in G_f is discussed in Sections V E and VI A. This motivates us to require that the fermionic d-cochain $\nu_d(g_0,...,g_d)$ to satisfy the following symmetry condition

$$\begin{aligned}&\nu_d^{s(g)}(gg_0,...,gg_d)\\ &=\nu_d(g_0,...,g_d)\prod_i[u^g_{d-1}(g_0,..,\hat{g}_i,..,g_d)]^{(-)^i}, \qquad \text{(C3)}\\ &n_{d-1}(gg_0,...,gg_{d-1}) = n_{d-1}(g_0,...,g_{d-1}), \quad g\in G_f,\end{aligned}$$

where the sequence $g_0,..,\hat{g}_i,..,g_d$ is the sequence $g_0,...,g_d$ with g_i removed, and $u^g_{d-1}(g_i,g_j,...,g_k)$ is an 1D representation of G_f

$$\begin{aligned}&u^{g_a}_{d-1}(g_i,...,g_k)[u^{g_b}_{d-1}(g_i,...,g_k)]^{s(g_a)} = u^{g_ag_b}_{d-1}(g_i,...,g_k),\\ &u^{g_a}_{d-1}(gg_i,gg_j,...,gg_k) = [u^{g_a}_{d-1}(g_i,g_j,...,g_k)]^{s(g)}, \qquad \text{(C4)}\end{aligned}$$

such that $u^{P_f}_{d-1}(g_i,g_j,...,g_k) = (-)^{n_{d-1}(g_i,g_j,...,g_k)}$. The triple functions $(\nu_d, n_{d-1}, u^g_{d-1})$ that satisfy eqn. (C3) and eqn. (C4) are called fermionic cochains. We will denote the collection of all those fermionic cochains $(\nu_d, n_{d-1}, u^g_{d-1})$ as $\mathscr{C}^d[G_f,U_T(1)]$.

4. Fermionic d-cocycle

With the above setup, now we are ready to define fermionic d-cocycle. The fermionic d-cocycles are fermionic d-cochains $[\nu_d(g_0,..,g_d), n_{d-1}(g_1,..,g_d), u^g_{d-1}(g_1,..,g_d)]$ in $\mathscr{C}^d[G_f,U_T(1)]$ that satisfy some additional conditions. Fermionic 1-cocycles are fermionic 1-cochains $[\nu_1(g_0,g_1), n_0(g_1), u^g_0(g_1)]$ that satisfy

$$\nu_1(g_1,g_2)\nu_1^{-1}(g_0,g_2)\nu_1(g_0,g_1) = 1 \tag{C5}$$

Note that $n_0(g_1)=0$ always. Fermionic 2-cocycles are fermionic 2-cochains $[\nu_2(g_0,g_1,g_2), n_1(g_1,g_2), u^g_1(g_1,g_2)]$ that satisfy

$$\nu_2(g_1,g_2,g_3)\nu_2^{-1}(g_0,g_2,g_3)\nu_2(g_0,g_1,g_3)\nu_2^{-1}(g_0,g_1,g_2) = 1 \tag{C6}$$

Fermionic 3-cocycles are fermionic 3-cochains $[\nu_3(g_0,g_1,g_2,g_3), n_2(g_1,g_2,g_3), u^g_2(g_1,g_2,g_3)]$ that satisfy

$$\begin{aligned}&(-)^{n_2(g_0,g_1,g_2)n_2(g_2,g_3,g_4)}\nu_3(g_1,g_2,g_3,g_4)\nu_3^{-1}(g_0,g_2,g_3,g_4)\times\\ &\nu_3(g_0,g_1,g_3,g_4)\nu_3^{-1}(g_0,g_1,g_2,g_4)\nu_3(g_0,g_1,g_2,g_3) = 1\end{aligned} \tag{C7}$$

Fermionic 4-cocycles are fermionic 4-cochains $[\nu_4(g_0,g_1,g_2,g_3,g_4), n_3(g_1,g_2,g_3,g_4), u^g_3(g_1,g_2,g_3,g_4)]$ that satisfy

$$(-)^{n_3(g_0,g_1,g_2,g_3)n_3(g_0,g_3,g_4,g_5)+n_3(g_1,g_2,g_3,g_4)n_3(g_0,g_1,g_4,g_5)+n_3(g_2,g_3,g_4,g_5)n_3(g_0,g_1,g_2,g_5)}\nu_4(g_1,g_2,g_3,g_4,g_5)\times$$
$$\nu_4^{-1}(g_0,g_2,g_3,g_4,g_5)\nu_4(g_0,g_1,g_3,g_4,g_5)\nu_4^{-1}(g_0,g_1,g_2,g_4,g_5)\nu_4(g_0,g_1,g_2,g_3,g_5)\nu_4^{-1}(g_0,g_1,g_2,g_3,g_4)=1 \quad \text{(C8)}$$

We will denote the collections of all d-cocycles $[\nu_d(g_0,...,g_d),n_{d-1}(g_1,...,g_d),u^g_{d-1}(g_1,...,g_d)]$ that satisfy the above conditions as $\mathscr{Z}^d[G_f,U_T(1)]$.

5. An "linearized" representation of group super-cohomology

Since the coefficient of the group super-cohomology is always $U(1)$, we can map the fermionic cochains $[\nu_d(g_0,...,g_d),n_{d-1}(g_1,...,g_d),u^g_{d-1}(g_1,...,g_d)]$ into two real vectors and an integer vector $(\boldsymbol{v}_d,\boldsymbol{n}_{d-1},\boldsymbol{h}^g_{d-1})$, where the components of $\boldsymbol{v}_d$ and $\boldsymbol{h}^g_{d-1}$ are in $[0,1)$ and the components of $\boldsymbol{n}_{d-1}$ are $0,1$. Then, we can rewrite the cochain and the cocycle conditions on $[\nu_d(g_0,...,g_d),n_{d-1}(g_1,...,g_d),u^g_{d-1}(g_1,...,g_d)]$ as "linear equations" on $(\boldsymbol{v}_d,\boldsymbol{n}_{d-1},\boldsymbol{h}^g_{d-1})$.

To map $\nu_d(g_0,...,g_d)$ into a real vector $\boldsymbol{v}_d$, we can introduce the log of $\nu_d(g_0,...,g_d)$:

$$v_d(g_0,g_1,...,g_d)=\frac{1}{2\pi \mathrm{i}}\log\nu_d(g_0,g_1,...,g_d) \text{ mod } 1, \quad \text{(C9)}$$

So we define $\boldsymbol{v}_d$ as a $|G_b|^{d+1}$ dimensional vector, whose components are given by

$$(\boldsymbol{v}_d)_{(g_0,...,g_d)}=v_d(1,g_0,...,g_d). \quad \text{(C10)}$$

Similarly, $\boldsymbol{n}_{d-1}$ is a $|G_b|^d$ dimensional integer vector

$$(\boldsymbol{n}_{d-1})_{(g_0,...,g_{d-1})}=n_d(g_0,...,g_{d-1}), \quad \text{(C11)}$$

and $\boldsymbol{h}^g_{d-1}$ is a $|G_b|^d$ dimensional vector

$$\begin{aligned}(\boldsymbol{h}^g_{d-1})_{(g_0,...,g_{d-1})}&=h^g_{d-1}(g_0,...,g_{d-1}), \quad \text{(C12)}\\ h^g_{d-1}(g_0,...,g_{d-1})&=\frac{1}{2\pi \mathrm{i}}\ln u^g_{d-1}(g_0,...,g_{d-1}) \text{ mod } 1.\end{aligned}$$

Now, the cochain conditions (*ie* the symmetry condition), eqn. (C3), eqn. (C1), and eqn. (C4), can be written in a "linearized" form

$$\begin{aligned}s(g)S^g_d\boldsymbol{v}_d&=\boldsymbol{v}_d+D_{d-1}\boldsymbol{h}^g_{d-1} \text{ mod } 1,\\ S^g_{d-1}\boldsymbol{n}_{d-1}&=\boldsymbol{n}_{d-1},\\ D_{d-1}\boldsymbol{n}_{d-1}&=0 \text{ mod } 2\\ s(g)S^g_{d-1}\boldsymbol{h}^{g_1}_{d-1}&=\boldsymbol{h}^{g_1}_{d-1},\\ \boldsymbol{h}^{g_1g_2}_{d-1}&=\boldsymbol{h}^{g_1}_{d-1}+s(g_1)\boldsymbol{h}^{g_2}_{d-1} \text{ mod } 1,\\ \boldsymbol{h}^{P_f}_{d-1}&=\frac{1}{2}\boldsymbol{n}_{d-1} \text{ mod } 1,\\ g,g_1,g_2&\in G_f, \quad \text{(C13)}\end{aligned}$$

where S^g_d is a $|G_b|^{d+1}\times|G_b|^{d+1}$ dimensional matrix, whose elements are

$$\begin{aligned}&(S^g_d)_{(g_0,...,g_d),(g'_0,...,g'_d)} \quad \text{(C14)}\\ &=\delta_{(g'_0,...,g'_d),(gg_0,...,gg_{d+1})}.\end{aligned}$$

Also D_d is a $|G_b|^{d+2}\times|G_b|^{d+1}$ dimensional integer matrix, whose elements are all 0 or ±1:

$$\begin{aligned}&(D_d)_{(g_0,...,g_{d+1}),(g'_0,...,g'_d)} \quad \text{(C15)}\\ &=(-)^i\delta_{(g'_0,...,g'_d),(g_0,...,\hat{g}_i,...,g_{d+1})},\end{aligned}$$

where $(g_0,...,\hat{g}_i,...,g_{d+1})$ is the sequence obtained from the sequence $(g_0,...,g_{d+1})$ with the element g_i removed. The matrices D_d satisfy

$$D_{d+1}D_d=0. \quad \text{(C16)}$$

A triple $(\boldsymbol{v}_d,\boldsymbol{n}_{d-1},\boldsymbol{h}^g_{d-1})$ that satisfies eqn. (C13) corresponds to a fermionic d-cochain in $\mathscr{C}^d[G_f,U_T(1)]$.

We can also rewrite the cocycle conditions eqn. (50) as a "linear" equation:

$$D_d\boldsymbol{v}_d+\frac{1}{2}\boldsymbol{f}_{d+1}=0 \text{ mod } 1. \quad \text{(C17)}$$

Here $\boldsymbol{f}_d$ is a $|G_b|^{d+1}$ dimensional integer vector whose components are given by

$$(\boldsymbol{f}_d)_{(g_0,...,g_d)}=f_d(g_0,...,g_d) \text{ mod } 2. \quad \text{(C18)}$$

where $f_d(g_0,...,g_d)$ is obtained from $n_{d-2}(g_0,...,g_{d-2})$ according to the relation eqn. (48). A triple $(\boldsymbol{v}_d,\boldsymbol{n}_{d-1},\boldsymbol{h}^g_{d-1})$ that satisfies eqn. (C13) and eqn. (C17) corresponds to a fermionic d-cocycke in $\mathscr{Z}^d[G_f,U_T(1)]$.

6. Fermionic d-cohomology class

As discussed in section V, each $(d+1)$-cocycle will define a fermionic topological non-linear σ model in d spatial dimensions which is a fixed point of the RG flow. Those $(d+1)$-cocycles give rise to new fermionic SPT phases. However, different $(d+1)$-cocycles may give rise to the same fermionic SPT phase. Those $(d+1)$-cocycles are said to be equivalent. To find (or guess) the equivalence relation between cocycles, in this section, we will discuss some natural choices of possible equivalence relations. A more physical discussion of this issue is given in Section G 1.

First, we would like to show that (for $d\leq4$)

$$D_d\boldsymbol{f}_d=0 \text{ mod } 2, \quad \text{(C19)}$$

where $\boldsymbol{f}_d$ is determined from $n_{d-1}(g_1,...,g_d)$ in $\mathcal{Z}^{d-1}(G_f,\mathbb{Z}_2)$ (see eqn. (48)). For $1 \leq d \leq 3$, the above is trivially satisfied since the corresponding $f_d = 0$. To show the above for $d = 4$, we note that $n_{d-2}(g_0,..,g_{d-2})$ (which give rise to $\boldsymbol{f}_d$) is a $(d-2)$-cocycle in $\mathcal{H}^{d-2}(G_b,\mathbb{Z}_2)$. Let us assume that a 2-cocycle $n_2(g_0,g_1,g_2)$ gives rise to $\boldsymbol{f}_4$. The expression, $f_4(g_0,g_1,g_2,g_3,g_4) = n_2(g_0,g_1,g_2)n_2(g_2,g_3,g_4)$ implies that the 4-cochain $f_4(g_0,g_1,g_2,g_3,g_4)$ is the cup product of two 2-cocycles $n_2(g_0,g_1,g_2)$ and $n_2(g_0,g_1,g_2)$ in $\mathcal{Z}^2(G_f,\mathbb{Z}_2)$. So $f_4(g_0,g_1,g_2,g_3,g_4)$ is also a cocycle in $\mathcal{Z}^4(G_f,\mathbb{Z}_2)$. Therefore $D_4\boldsymbol{f}_4 = 0 \mod 2$. We can also show that $D_5\boldsymbol{f}_5 = 0 \mod 2$ by an explicit calculation using Mathematica. Therefore, for $d \leq 5$,

$$D_d[D_{d-1}\boldsymbol{v}_{d-1} + \frac{1}{2}\boldsymbol{f}_d] = 0 \text{ mod } 1. \tag{C20}$$

for any fermionic $(d-1)$-cochains $(\boldsymbol{v}_{d-1},\boldsymbol{n}_{d-1},\boldsymbol{h}^g_{d-1})$.

There is another important relation. Let us assume that a d-cocycle $n_d(g_0,g_1,...,g_d)$ gives rise to $\boldsymbol{f}_{d+2}$, and another d-cocycle $\tilde{n}_2(g_0,g_1,...,g_d)$ gives rise to $\tilde{\boldsymbol{f}}_{d+2}$ (assuming $d \leq 3$). If $n_d(g_0,g_1,...,g_d)$ and $\tilde{n}_d(g_0,g_1,...,g_d)$ are related by a coboundary in $\mathcal{B}^d(G_b,\mathbb{Z}_2)$

$$\begin{aligned}
&n_d(g_0,g_1,...,g_d) + \sum_i m'_{d-1}(g_0,...,\hat{g}_i,...,g_d)\\
&= n_d(g_0,g_1,...,g_d) + (\delta m'_{d-1})(g_0,...,g_d)\\
&= \tilde{n}_d(g_0,g_1,...,g_d) \text{ mod } 2,
\end{aligned} \tag{C21}$$

then what is the relation between $\boldsymbol{f}_{d+2}$ and $\tilde{\boldsymbol{f}}_{d+2}$? In the Appendix D, we will show that $\boldsymbol{f}_{d+2}$ and $\tilde{\boldsymbol{f}}_{d+2}$ also differ by a coboundary $D_{d+1}\boldsymbol{g}_{d+1}$ in $\mathcal{B}^{d+2}(G_b,\mathbb{Z}_2)$:

$$\tilde{\boldsymbol{f}}_{d+2} = \boldsymbol{f}_{d+2} + D_{d+1}\boldsymbol{g}_{d+1} \text{ mod } 2, \tag{C22}$$

where $\boldsymbol{g}_{d+1}$ is determined from $\boldsymbol{n}_d$ and $\boldsymbol{m}_{d-1}$ (see Appendix D).

Using the above two relations, we can define the first equivalence relation between two fermionic d-cocycles. Starting from a fermionic d-cocycle, $(\boldsymbol{v}_d,\boldsymbol{n}_{d+1},\boldsymbol{h}^g_{d-1})$, we can use a fermionic $(d-1)$-cochain $(\boldsymbol{v}_{d-1},\boldsymbol{n}_{d-2},\boldsymbol{h}^g_{d-2})$ to transform the above d-cocycle to an equivalent d-cocycle $(\tilde{\boldsymbol{v}}_d,\tilde{\boldsymbol{n}}_{d-1},\tilde{\boldsymbol{h}}^g_{d-1})$:

$$\begin{aligned}
\tilde{\boldsymbol{v}}_d =& \boldsymbol{v}_d - \frac{1}{2}\boldsymbol{g}_d + D_{d-1}\boldsymbol{v}_{d-1},\\
\tilde{\boldsymbol{n}}_{d-1} =& \boldsymbol{n}_{d-1} + D_{d-2}\boldsymbol{n}_{d-2}\\
\tilde{\boldsymbol{h}}^g_{d-1} =& \boldsymbol{h}^g_{d-1} + D_{d-2}\boldsymbol{h}^g_{d-2},
\end{aligned} \tag{C23}$$

where $\boldsymbol{g}_d$ is determined from $\boldsymbol{n}_{d-1}$ and $\boldsymbol{n}_{d-2}$ (see Appendix D). We note that $\boldsymbol{g}_d = 0$ if $\boldsymbol{n}_{d-2} = 0$.

First, we would like to show that if $(\boldsymbol{v}_d,\boldsymbol{n}_{d-1},\boldsymbol{h}^g_{d-1})$ satisfies the cochain condition eqn. (C13), then $(\tilde{\boldsymbol{v}}_d,\tilde{\boldsymbol{n}}_{d+1},\tilde{\boldsymbol{h}}^g_{d-1})$ obtained above also satisfies the same cochain condition. Let us first consider the first equation in eqn. (C13):

$$\begin{aligned}
&s(g)S^g_d\tilde{\boldsymbol{v}}_d = s(g)S^g_d\boldsymbol{v}_d - \frac{1}{2}\boldsymbol{g}_d + s(g)S^g_d D_{d-1}\boldsymbol{v}_{d-1}\\
=&\boldsymbol{v}_d + D_{d-1}\boldsymbol{h}^g_{d-1} - \frac{1}{2}\boldsymbol{g}_d + D_{d-1}s(g)S^g_{d-1}\boldsymbol{v}_{d-1} \text{ mod } 1\\
=&\tilde{\boldsymbol{v}}_d + D_{d-1}(\boldsymbol{h}^g_{d-1} + D_{d-2}\boldsymbol{h}^g_{d-2}) \text{ mod } 1,\\
=&\tilde{\boldsymbol{v}}_d + D_{d-1}\tilde{\boldsymbol{h}}^g_{d-1} \text{ mod } 1,
\end{aligned} \tag{C24}$$

where we have used

$$S^g_d\boldsymbol{g}_d = \boldsymbol{g}_d, \tag{C25}$$

and the relation

$$S^g_d D_{d-1} = D_{d-1}S^g_{d-1}. \tag{C26}$$

It is easy to show that $(\tilde{\boldsymbol{v}}_d,\tilde{\boldsymbol{n}}_{d+1},\tilde{\boldsymbol{h}}^g_{d-1})$ also satisfies other equations in eqn. (C13).

Now we like to show that if $(\boldsymbol{v}_d,\boldsymbol{n}_{d-1},\boldsymbol{h}^g_{d-1})$ satisfies the cocycle condition eqn. (C17), then $(\tilde{\boldsymbol{v}}_d,\tilde{\boldsymbol{n}}_{d+1},\tilde{\boldsymbol{h}}^g_{d-1})$ obtained above also satisfies the same cocycle condition:

$$\begin{aligned}
&D_d\tilde{\boldsymbol{v}}_d + \frac{1}{2}\tilde{\boldsymbol{f}}_{d+1}\\
&= D_d\boldsymbol{v}_d - \frac{1}{2}D_d\boldsymbol{g}_d + \frac{1}{2}(\boldsymbol{f}_{d+1} + D_d\boldsymbol{g}_d)\\
&= 0 \text{ mod } 1.
\end{aligned} \tag{C27}$$

We can also define the second equivalence relation between two fermionic d-cocycles. Let $\bar{h}^g_{d-1}(g_0,...,g_{d-1})$ be representations of G_b that satisfy

$$\begin{aligned}
\bar{h}^{g_a}_{d-1}(g_0,g_1,..)+s(g_a)\bar{h}^{g_b}_{d-1}(g_0,g_1,..) &= \bar{h}^{g_ag_b}_{d-1}(g_0,g_1,..),\\
s(g)\bar{h}^{g_a}_{d-1}(gg_0,...,gg_{d-1}) &= \bar{h}^{g_a}_{d-1}(g_0,...,g_{d-1}),\\
g,g_a,g_b \in G_b.&
\end{aligned} \tag{C28}$$

Then, starting from a fermionic d-cocycle, $(\boldsymbol{v}_d,\boldsymbol{n}_{d+1},\boldsymbol{h}^g_{d-1})$, we can use such a $\bar{h}^g_{d-1}(g_0,...,g_{d-1})$ to transform the above d-cocycle to an equivalent d-cocycle $(\tilde{\boldsymbol{v}}_d,\tilde{\boldsymbol{n}}_{d-1},\tilde{\boldsymbol{h}}^g_{d-1})$:

$$\begin{aligned}
&\tilde{v}_d(g_0,g_1,...,g_d) = v_d(g_0,g_1,...,g_d)\\
&\quad + \bar{h}^{g_1}_{d-1}(g_1,...,g_d) + \sum_{i=1}^d(-)^i\bar{h}^{g_0}_{d-1}(g_0,..,\hat{g}_i,..,g_d),\\
&\tilde{n}_{d-1}(g_0,g_1,...,g_{d-1}) = n_{d-1}(g_0,g_1,...,g_{d-1}), \qquad \text{(C29)}\\
&\tilde{h}^g_{d-1}(g_0,..,g_{d-1}) = h^g_{d-1}(g_0,..,g_{d-1}) + \bar{h}^g_{d-1}(g_0,..,g_{d-1}).
\end{aligned}$$

The above can also be written in a more compact form:

$$\begin{aligned}
\tilde{\boldsymbol{v}}_d &= \boldsymbol{v}_d + D_{d-1}\bar{\boldsymbol{h}}_{d-1},\\
\tilde{\boldsymbol{n}}_{d-1} &= \boldsymbol{n}_{d-1},\\
\tilde{\boldsymbol{h}}^g_{d-1} &= \boldsymbol{h}^g_{d-1} + \bar{\boldsymbol{h}}^g_{d-1},
\end{aligned} \tag{C30}$$

where $\bar{\boldsymbol{h}}_{d-1}$ and $\bar{\boldsymbol{h}}^g_{d-1}$ the $|G_b|^d$ dimensional vectors:

$$\begin{aligned}
(\bar{\boldsymbol{h}}_{d-1})_{(g_0,...,g_{d-1})} &= h^{g_0}_{d-1}(g_0,...,g_{d-1}),\\
(\bar{\boldsymbol{h}}^g_{d-1})_{(g_0,...,g_{d-1})} &= h^g_{d-1}(g_0,...,g_{d-1}).
\end{aligned} \tag{C31}$$

Let us first show that $(\tilde{\boldsymbol{v}}_d, \tilde{\boldsymbol{n}}_{d-1}, \tilde{\boldsymbol{h}}^g_{d-1})$ is a d-cochain:

$$\begin{aligned}
&s(g)\tilde{v}_d(gg_0, gg_1, ..., gg_d) = s(g)v_d(gg_0, gg_1, ..., gg_d) \\
&+ s(g)\bar{h}^{gg_1}_{d-1}(gg_1, .., gg_d) + s(g)\sum_{i=1}^{d}(-)^i\bar{h}^{gg_0}_{d-1}(gg_0, .., g\hat{g}_i, .., gg_d) \\
&= v_d(g_0, g_1, ..., g_d) + \sum_{i=0}^{d}(-)^i h^g_{d-1}(g_0, .., \hat{g}_i, .., g_d) \quad \text{(C32)} \\
&\quad + \bar{h}^{gg_1}_{d-1}(g_1, .., g_d) + \sum_{i=0}^{d}(-)^i\bar{h}^{gg_0}_{d-1}(g_0, .., \hat{g}_i, .., g_d) \\
&= \tilde{v}_d(g_0, g_1, ..., g_d) + \sum_{i=0}^{d}(-)^i\tilde{h}^g_{d-1}(g_0, .., \hat{g}_i, .., g_d).
\end{aligned}$$

Second, since $\tilde{\boldsymbol{v}}_d = \boldsymbol{v}_d + D_{d-1}\bar{\boldsymbol{h}}_{d-1}$, it also clear that $(\tilde{\boldsymbol{v}}_d, \tilde{\boldsymbol{n}}_{d-1}, \tilde{\boldsymbol{h}}^g_{d-1})$ is a d-cocycle.

The equivalence classes of fermionic d-cocycles obtained from the above two kinds of equivalence relations, eqn. (C24) and eqn. (C29), form the group d-super-cohomology $\mathscr{H}^d[G_f, U_T(1)]$.

We note that, when $\boldsymbol{n}_{d-1} = 0$, the representations of G_f, $\boldsymbol{h}^g_{d-1}$, are also representations of G_b. So the second equivalence relation implies that a d-cocycle $(\boldsymbol{v}_d, \boldsymbol{n}_{d-1} = 0, \boldsymbol{h}^g_{d-1})$ is always equivalent to a simpler d-cocycle $(\tilde{\boldsymbol{v}}_d, \boldsymbol{n}_{d-1} = 0, \boldsymbol{h}^g_{d-1} = 0)$. Such a kind of d-cocycles are described by group cohomology and form group cohomology class $\mathcal{H}^d[G_b, U_T(1)]$.

We also note that, due to the requirement $\boldsymbol{h}^{P_f}_{d-1} = \frac{1}{2}\boldsymbol{n}_{d-1}$ mod 1, for each fixed $\boldsymbol{n}_{d-1}$, all allowed choices of $\boldsymbol{h}^g_{d-1}$ belong to one class under the second equivalence relation. So we just need to choose one allowed $\boldsymbol{h}^g_{d-1}$ for each $\boldsymbol{n}_{d-1}$.

7. Group super-cohomology classes and fermionic SPT phases

Each element in group super-cohomology class $\mathscr{H}^d[G_f, U_T(1)]$ correspond to a fermionic topological non-linear σ-model defined via the fermion path integral in imaginary time (see section IV). The fermion path integral over a $(d_{sp}+1)$-dimensional complex Σ will give rise to a ground state on its d_{sp} boundary. Such a ground state represents the corresponding fermionic SPT state. So each element in group super-cohomology class $\mathscr{H}^d[G_f, U_T(1)]$ corresponds to a fermionic SPT state.

Note that, in $d_{sp} = 0$ spatial dimension, the elements in $\mathscr{H}^1[G_f, U_T(1)]$ can have the trivial graded structure $n_0(g_0) = 0$, or the non-trivial graded structure $n_0(g_0) = 1$. The corresponding fermionic gapped states can have even or odd numbers of fermions. So we can have two different fermionic SPT phases in $d_{sp} = 0$ spatial dimension even without symmetry.

8. A simplified "linear" equations for group super-cohomology

The cochain condition (or the symmetry condition) eqn. (C13) can be solved, which implies that only the following components $v_d(1, g_1, ..., g_d)$, $n_{d-1}(1, g_1, ..., g_{d-1})$, and $h^g_{d-1}(1, g_1, ..., g_{d-1})$, are independent. Other components with $g_0 \neq 1$ can be expressed from those independent components:

$$\begin{aligned}
n_{d-1}(g_0, g_1, ..g_{d-1}) &= n_{d-1}(1, g_0^{-1}g_1, ..g_0^{-1}g_{d-1}), \quad \text{(C33)} \\
s(g_0)h^g_{d-1}(g_0, g_1, .., g_{d-1}) &= h^g_{d-1}(1, g_0^{-1}g_1, .., g_0^{-1}g_{d-1}),
\end{aligned}$$

and

$$\begin{aligned}
v_d(g_0, g_1, ..., g_d) = s(g_0)\Big(& v_d(1, g_0^{-1}g_1, ..., g_0^{-1}g_d) \\
&+ \sum_{i=0}^{d}(-)^i h^{g_0^{-1}}_d(g_0^{-1}g_0, ..., g_0^{-1}\hat{g}_i, ..., g_0^{-1}g_d)\Big). \quad \text{(C34)}
\end{aligned}$$

Using the independent components, we can rewrite the cocycle condition as

$$v_d(g_1, g_2, ..., g_{d+1}) - v_d(1, g_2, ..., g_{d+1}) + v_d(1, g_1, ..., g_{d+1}) \text{ ... ... } + f_{d+1}(1, g_1, g_2, ..., g_{d+1}) = 0 \text{ mod } 1, \qquad \text{(C35)}$$

which can be rewritten as

$$\begin{aligned}
& s(g_1)\Big(v_d(1, g_1^{-1}g_2, ..., g_1^{-1}g_{d+1}) + s(g_1^{-1}g_2)h^{g_1^{-1}}_{d-1}(1, g_2^{-1}g_3, ..., g_2^{-1}g_{d+1}) - h^{g_1^{-1}}_{d-1}(1, g_1^{-1}g_3, ..., g_1^{-1}g_{d+1}) \\
&\quad + h^{g_1^{-1}}_{d-1}(1, g_1^{-1}g_2, ..., g_1^{-1}g_{d+1}) \text{ ... ...}\Big) - v_d(1, g_2, ..., g_{d+1}) + v_d(1, g_1, ..., g_{d+1}) \text{ ... ... } + f_{d+1}(1, g_1, g_2, ..., g_{d+1}) \\
&= s(g_1)v_d(1, g_1^{-1}g_2, ..., g_1^{-1}g_{d+1}) - v_d(1, g_2, ..., g_{d+1}) + v_d(1, g_1, ..., g_{d+1}) \text{ ... ... } + f_{d+1}(1, g_1, g_2, ..., g_{d+1}) \\
&\quad + s(g_2)h^{g_1^{-1}}_{d-1}(1, g_2^{-1}g_3, ..., g_2^{-1}g_{d+1}) - s(g_1)\sum_{i=2}^{d+1}(-)^i h^{g_1^{-1}}_{d-1}(1, g_1^{-1}g_2, ...g_1^{-1}\hat{g}_i, ..., g_1^{-1}g_{d+1}) = 0 \text{ mod } 1 \qquad \text{(C36)}
\end{aligned}$$

These are the simplified "linear" equations for group super-cohomology. They include both the cochain (C13) and cocycle (C17) conditions.

Appendix D: The mapping $n_d \to f_{d+2}$ induces a mapping $\mathcal{H}^d(G_b,\mathbb{Z}_2) \to \mathcal{H}^{d+2}(G_b,\mathbb{Z}_2)$

We note that eqn. (48) defines a mapping from a d-cochain $n_d \in \mathcal{C}^d(G_b,\mathbb{Z}_2)$ to a $(d+2)$-cochain $f_d \in \mathcal{C}^{d+2}(G_b,\mathbb{Z}_2)$. We can show that, if n_d is a cocycle $n_d \in \mathcal{Z}^d(G_b,\mathbb{Z}_2)$, then the corresponding f_{d+2} is also a cocycle $f_d \in \mathcal{Z}^{d+2}(G_b,\mathbb{Z}_2)$. Thus eqn. (48) defines a mapping from a d-cocycle $n_d \in \mathcal{Z}^d(G_b,\mathbb{Z}_2)$ to a $(d+2)$-cocycle $f_d \in \mathcal{Z}^{d+2}(G_b,\mathbb{Z}_2)$. In this section, we are going to show that eqn. (48) actually define a mapping from d-cohomology classes to $(d+2)$-cohomology classes: $\mathcal{H}^d(G_b,\mathbb{Z}_2) \to \mathcal{H}^{d+2}(G_b,\mathbb{Z}_2)$. This is because if n_d and $\tilde n_d$ differ by a coboundary, then the corresponding f_{d+2} and $\tilde f_{d+2}$ also differ by a coboundary.

To show this, let us first assume $d = 2$. Since the 4-cochain $f_4(g_0,g_1,g_2,g_3,g_4)$ in $\mathcal{C}^4(G_b,\mathbb{Z}_2)$ is the cup product to two 2-cocycles $n_2(g_0,g_1,g_2)$ and $n_2(g_0,g_1,g_2)$ in $\mathcal{Z}^2(G_b,\mathbb{Z}_2)$, so if we change the two cocycles by a coboundary, the 4-cochain f_4 will also change by a coboundary. Thus there exist a vector $\boldsymbol{g}_3$ such that

$$\tilde{\boldsymbol{f}}_4 = \boldsymbol{f}_4 + D_3\boldsymbol{g}_3 \text{ mod } 2. \tag{D1}$$

and eqn. (C22) is valid for $d = 2$.

To show eqn. (C22) to be valid for $d = 3$, we need to show when $n_3(g_0,g_1,g_2,g_3)$ and $\tilde n_3(g_0,g_1,g_2,g_3)$ are related by a coboundary in $\mathcal{B}^3(G_b,\mathbb{Z}_2)$:

$$\begin{aligned}\tilde n_3(g_0,g_1,g_2,g_3) &= n_3(g_0,g_1,g_2,g_3) + m_2'(g_1,g_2,g_3)\\ &+ m_2'(g_0,g_2,g_3) + m_2'(g_0,g_1,g_3) + m_2'(g_0,g_1,g_2) \text{ mod } 2,\end{aligned} \tag{D2}$$

the corresponding $\boldsymbol{f}_5$ and $\tilde{\boldsymbol{f}}_5$ are also related by a coboundary in $\mathcal{B}^5(G_b,\mathbb{Z}_2)$:

$$\boldsymbol{f}_5 = \tilde{\boldsymbol{f}}_5 + D_4\boldsymbol{g}_4 \text{ mod } 2. \tag{D3}$$

Let us introduce

$$\begin{aligned}n_3'(g_0,g_1,g_2,g_3) &= m_2'(g_1,g_2,g_3) + m_2'(g_0,g_2,g_3)\\ &+ m_2'(g_0,g_1,g_3) + m_2'(g_0,g_1,g_2).\end{aligned} \tag{D4}$$

Then we can express $\tilde f_5$ as $\tilde f_5 = f_5 + f_5^I + f_5^{II} + f_5^{III}$, with:

$$\begin{aligned}
&f_5^I(g_0,g_1,g_2,g_3,g_4,g_5)\\
=&n_3(g_0,g_1,g_2,g_3)n_3'(g_0,g_3,g_4,g_5) + n_3(g_1,g_2,g_3,g_4)n_3'(g_0,g_1,g_4,g_5) + n_3(g_2,g_3,g_4,g_5)n_3'(g_0,g_1,g_2,g_5),\\
&f_5^{II}(g_0,g_1,g_2,g_3,g_4,g_5)\\
=&n_3'(g_0,g_1,g_2,g_3)n_3(g_0,g_3,g_4,g_5) + n_3'(g_1,g_2,g_3,g_4)n_3(g_0,g_1,g_4,g_5) + n_3'(g_2,g_3,g_4,g_5)n_3(g_0,g_1,g_2,g_5),\\
&f_5^{III}(g_0,g_1,g_2,g_3,g_4,g_5)\\
=&n_3'(g_0,g_1,g_2,g_3)n_3'(g_0,g_3,g_4,g_5) + n_3'(g_1,g_2,g_3,g_4)n_3'(g_0,g_1,g_4,g_5) + n_3'(g_2,g_3,g_4,g_5)n_3'(g_0,g_1,g_2,g_5).
\end{aligned} \tag{D5}$$

In the following, we would like to show g_4 can be constructed as $g_4 = g_4^I + g_4^{II} + g_4^{III}$, with:

$$\begin{aligned}
g_4^I(g_0,g_1,g_2,g_3,g_4) &= n_3(g_0,g_1,g_2,g_3)m_2'(g_0,g_3,g_4) + n_3(g_1,g_2,g_3,g_4)m_2'(g_0,g_1,g_4),\\
g_4^{II}(g_0,g_1,g_2,g_3,g_4) &= n_3(g_0,g_2,g_3,g_4)m_2'(g_0,g_1,g_2) + n_3(g_0,g_1,g_2,g_4)m_2'(g_2,g_3,g_4) + n_3(g_0,g_1,g_3,g_4)m_2'(g_1,g_2,g_3),\\
g_4^{III}(g_0,g_1,g_2,g_3,g_4) &= n_3'(g_0,g_1,g_2,g_3)m_2'(g_0,g_3,g_4) + n_3'(g_1,g_2,g_3,g_4)m_2'(g_0,g_1,g_4) + m_2'(g_0,g_1,g_2)m_2'(g_2,g_3,g_4).
\end{aligned} \tag{D6}$$

Let us prove the above statement in three steps:
First step: After plug-in, we find that $\boldsymbol{f}_5^I - D_4\boldsymbol{g}_4^I$ is given by

$$\begin{aligned}
&f_5^I(g_0,g_1,g_2,g_3,g_4,g_5) - g_4^I(g_0,g_1,g_2,g_3,g_4) - g_4^I(g_0,g_1,g_2,g_3,g_5) - g_4^I(g_0,g_1,g_2,g_4,g_5)\\
&\qquad - g_4^I(g_0,g_1,g_3,g_4,g_5) - g_4^I(g_0,g_2,g_3,g_4,g_5) - g_4^I(g_1,g_2,g_3,g_4,g_5)\\
=\ & n_3(g_0,g_1,g_2,g_3)\,[m_2'(g_0,g_3,g_4) + m_2'(g_0,g_3,g_5) + m_2'(g_0,g_4,g_5) + m_2'(g_3,g_4,g_5)]\\
&+ n_3(g_1,g_2,g_3,g_4)\,[m_2'(g_0,g_1,g_4) + m_2'(g_0,g_1,g_5) + m_2'(g_0,g_4,g_5) + m_2'(g_1,g_4,g_5)]\\
&+ n_3(g_2,g_3,g_4,g_5)\,[m_2'(g_0,g_1,g_2) + m_2'(g_0,g_1,g_5) + m_2'(g_0,g_2,g_5) + m_2'(g_1,g_2,g_5)]\\
&- n_3(g_0,g_1,g_2,g_3)m_2'(g_0,g_3,g_4) - n_3(g_1,g_2,g_3,g_4)m_2'(g_0,g_1,g_4) - n_3(g_0,g_1,g_2,g_3)m_2'(g_0,g_3,g_5)\\
&- n_3(g_1,g_2,g_3,g_5)m_2'(g_0,g_1,g_5) - n_3(g_0,g_1,g_2,g_4)m_2'(g_0,g_4,g_5) - n_3(g_1,g_2,g_4,g_5)m_2'(g_0,g_1,g_5)\\
&- n_3(g_0,g_1,g_3,g_4)m_2'(g_0,g_4,g_5) - n_3(g_1,g_3,g_4,g_5)m_2'(g_0,g_1,g_5) - n_3(g_0,g_2,g_3,g_4)m_2'(g_0,g_4,g_5)\\
&- n_3(g_2,g_3,g_4,g_5)m_2'(g_0,g_2,g_5) - n_3(g_1,g_2,g_3,g_4)m_2'(g_1,g_4,g_5) - n_3(g_2,g_3,g_4,g_5)m_2'(g_1,g_2,g_5)
\end{aligned}$$

We note that there are 5 terms containing $m'_2(g_0,g_4,g_5)$ and 5 terms containing $m'_2(g_0,g_1,g_5)$. Using the condition that $n_3(g_0,g_1,g_2,g_3)$ is a 3 cocycle:

$$n_3(g_0,g_1,g_2,g_3)+n_3(g_0,g_1,g_2,g_3)+n_3(g_0,g_1,g_2,g_3)+n_3(g_0,g_1,g_2,g_3)+n_3(g_0,g_1,g_2,g_3)=0 \text{ mode } 2, \tag{D7}$$

we find that those terms cancel out. Also, there are two terms containing $m'_2(g_0,g_3,g_4)$, two terms containing $m'_2(g_0,g_4,g_5)$, ..., etc. Those terms also cancel out under the mod 2 calculation. But the $m'_2(g_0,g_1,g_2)$ term and the $m'_2(g_3,g_4,g_5)$ term appear only once. Thus we simplify the above to

$$\begin{aligned}
&f^I_5(g_0,g_1,g_2,g_3,g_4,g_5)-g^I_4(g_0,g_1,g_2,g_3,g_4)-g^I_4(g_0,g_1,g_2,g_3,g_5)-g^I_4(g_0,g_1,g_2,g_4,g_5)\\
&\qquad -g^I_4(g_0,g_1,g_3,g_4,g_5)-g^I_4(g_0,g_2,g_3,g_4,g_5)-g^I_4(g_1,g_2,g_3,g_4,g_5)\\
=&n_3(g_0,g_1,g_2,g_3)m'_2(g_3,g_4,g_5)+n_3(g_2,g_3,g_4,g_5)m'_2(g_0,g_1,g_2) \bmod 2
\end{aligned} \tag{D8}$$

Second step: Similarly, we have

$$\begin{aligned}
&f^{II}_5(g_0,g_1,g_2,g_3,g_4,g_5)-g^{II}_4(g_0,g_1,g_2,g_3,g_4)-g^{II}_4(g_0,g_1,g_2,g_3,g_5)-g^{II}_4(g_0,g_1,g_2,g_4,g_5)\\
&\qquad -g^{II}_4(g_0,g_1,g_3,g_4,g_5)-g^{II}_4(g_0,g_2,g_3,g_4,g_5)-g^{II}_4(g_1,g_2,g_3,g_4,g_5)\\
=&-n_3(g_0,g_1,g_2,g_3)m'_2(g_3,g_4,g_5)-n_3(g_2,g_3,g_4,g_5)m'_2(g_0,g_1,g_2) \bmod 2
\end{aligned} \tag{D9}$$

Third step: Finally, by using the results derived in the first step, we have

$$\begin{aligned}
&f^{III}_5(g_0,g_1,g_2,g_3,g_4,g_5)-g^{III}_4(g_0,g_1,g_2,g_3,g_4)-g^{III}_4(g_0,g_1,g_2,g_3,g_5)-g^{III}_4(g_0,g_1,g_2,g_4,g_5)\\
&\qquad -g^{III}_4(g_0,g_1,g_3,g_4,g_5)-g^{III}_4(g_0,g_2,g_3,g_4,g_5)-g^{III}_4(g_1,g_2,g_3,g_4,g_5)\\
=&n'_3(g_0,g_1,g_2,g_3)m'_2(g_3,g_4,g_5)+n'_3(g_2,g_3,g_4,g_5)m'_2(g_0,g_1,g_2)\\
&\quad +m'_2(g_0,g_1,g_2)m'_2(g_2,g_3,g_4)+m'_2(g_0,g_1,g_2)m'_2(g_2,g_3,g_5)+m'_2(g_0,g_1,g_2)m'_2(g_2,g_4,g_5)\\
&\quad +m'_2(g_0,g_1,g_3)m'_2(g_3,g_4,g_5)+m'_2(g_0,g_2,g_3)m'_2(g_3,g_4,g_5)+m'_2(g_1,g_2,g_3)m'_2(g_3,g_4,g_5) \bmod 2\\
=&0 \bmod 2
\end{aligned} \tag{D10}$$

Combine the results from the above three steps, we can show:

$$D_4\boldsymbol{g}_4=\boldsymbol{f}^I_5+\boldsymbol{f}^{II}_5+\boldsymbol{f}^{III}_5 \bmod 2 \tag{D11}$$

Thus, we prove eqn. (C22) for $d=3$.

We like to mention that the induced mapping $\mathcal{H}^d(G_b,\mathbb{Z}_2)\to\mathcal{H}^{d+2}(G_b,\mathbb{Z}_2)$ by the $n_d\to f_{d+2}$ mapping (48), appears to be the Steenrod square Sq^2.[64,65] This realization will allow us to generalize eqn. (48) to higher dimensions.

Appendix E: Calculate $\mathcal{H}^d(G_b,\mathbb{Z}_2)$ – the graded structure

1. Calculate $\mathcal{H}^d(Z_n,\mathbb{Z}_2)$

The cohomology group $\mathcal{H}^d(Z_n,M)$ has a very simple form. To describe the simple form in a more general setting, let us define Tate cohomology groups $\hat{\mathcal{H}}^d(G,M)$.

For d to be 0 or -1, we have

$$\begin{aligned}
\hat{\mathcal{H}}^0(G,M)&=M^G/\mathrm{Img}(N_G,M),\\
\hat{\mathcal{H}}^{-1}(G,M)&=\mathrm{Ker}(N_G,M)/I_GM.
\end{aligned} \tag{E1}$$

Here M^G, $\mathrm{Img}(N_G,M)$, $\mathrm{Ker}(N_G,M)$, and I_GM are submodule of M. M^G is the maximal submodule that is invariant under the group action. Let us define a map $N_G: M\to M$ as

$$a\to\sum_{g\in G}g\cdot a,\quad a\in M. \tag{E2}$$

$\mathrm{Img}(N_G,M)$ is the image of the map and $\mathrm{Ker}(N_G,M)$ is the kernel of the map. The submodule I_GM is given by

$$I_GM=\{\sum_{g\in G}n_g g\cdot a|\sum_{g\in G}n_g=0,\ a\in M\} \tag{E3}$$

In other words, I_GM is generated by $g\cdot a-a$, $\forall\ g\in G$, $a\in M$.

For d other then 0 and -1, Tate cohomology groups $\hat{\mathcal{H}}^d(G,M)$ is given by

$$\begin{aligned}
\hat{\mathcal{H}}^d(G,M)&=\mathcal{H}^d(G,M),\quad \text{for } d>0\\
\hat{\mathcal{H}}^d[G,M]&=\mathcal{H}_{-d-1}[G,M],\quad \text{for } d<-1.
\end{aligned} \tag{E4}$$

For cyclic group Z_n, its (Tate) group cohomology over a generic Z_n-module M is given by[51,66]

$$\hat{\mathcal{H}}^d(Z_n,M)=\begin{cases}\hat{\mathcal{H}}^0(Z_n,M) & \text{if } d=0 \bmod 2,\\ \hat{\mathcal{H}}^{-1}(Z_n,M) & \text{if } d=1 \bmod 2.\end{cases} \tag{E5}$$

where

$$\begin{aligned}\hat{\mathcal{H}}^0(Z_n,M)&=M^{Z_n}/\mathrm{Img}(N_{Z_n},M),\\ \hat{\mathcal{H}}^{-1}(Z_n,M)&=\mathrm{Ker}(N_{Z_n},M)/I_{Z_n}M.\end{aligned}\tag{E6}$$

For example, when the group action is trivial, we have $M^{Z_n}=M$ and $I_{Z_n}M=\mathbb{Z}_1$. The map N_{Z_n} becomes $N_{Z_n}:a\to na$. For $M=\mathbb{Z}_2$, we have $\mathrm{Img}(N_{Z_n},\mathbb{Z}_2)=n\mathbb{Z}_2$ and $\mathrm{Ker}(N_{Z_n},\mathbb{Z}_2)=\mathbb{Z}_{(n,2)}$ (where (n,m) is the greatest common divisor of n and m). So we have

$$\mathcal{H}^d(Z_n,\mathbb{Z}_2)=\begin{cases}\mathbb{Z}_2 & \text{if } d=0,\\ \mathbb{Z}_{(n,2)} & \text{if } d>0.\end{cases}\tag{E7}$$

What is the non-trivial cocycle in $\mathcal{H}^d[Z_2,\mathbb{Z}_2]$? In Ref. 50, it is shown that a d-cocycle can be chosen to satisfy

$$\begin{aligned}&\nu_d(\mathbf{g},\mathbf{g},g_2,g_3,...,g_{d-2},g_{d-1},g_d)=1\\ &\nu_d(g_0,\mathbf{g},\mathbf{g},g_3,...,g_{d-2},g_{d-1},g_d)=1\\ &\qquad ...\quad ...\quad ...\\ &\nu_d(g_0,g_1,g_2,g_3,...,g_{d-2},\mathbf{g},\mathbf{g})=1.\end{aligned}\tag{E8}$$

by adding a proper coboundary.

Let us denote the two elements in Z_2 as $\{E,\sigma\}$ with $\sigma^2=E$. If we choose the cocycle $\nu_d(g_0,g_1,g_2,...)$ in $\mathcal{H}^d(Z_2,\mathbb{Z}_2)$ to satisfy the above condition, then only $\nu_d(E,\sigma,E,...)=\nu_d(\sigma,E,\sigma,...)$ can be non zero. Since $\mathcal{H}^d(Z_2,\mathbb{Z}_2)=\mathbb{Z}_2$, So $\nu_d(E,\sigma,E,...)=\nu_d(\sigma,E,\sigma,...)=0$ must correspond to the trivial class and $\nu_d(E,\sigma,E,...)=\nu_d(\sigma,E,\sigma,...)=1$ must correspond to the non-trivial class in $\mathcal{H}^d(Z_2,\mathbb{Z}_2)$.

2. Calculate $\mathcal{H}^d(G,\mathbb{Z}_2)$ from $\mathcal{H}^d(G,\mathbb{Z})$ using Künneth formula for group cohomology

We can also calculate $\mathcal{H}^d(G,\mathbb{Z}_2)$ from $\mathcal{H}^d(G,\mathbb{Z})$ using the Künneth formula for group cohomology. Let M (resp. M') be an arbitrary G-module (resp. G'-module) over a principal ideal domain R. We also assume that either M or M' is R-free. Then we have a Künneth formula for group cohomology[67,68]

$$\begin{aligned}\mathbb{Z}_1&\to\prod_{p=0}^{d}\mathcal{H}^p(G,M)\otimes_R\mathcal{H}^{d-p}(G',M')\\ &\to\mathcal{H}^d(G\times G',M\otimes_R M')\\ &\to\prod_{p=0}^{d+1}\mathrm{Tor}_1^R(\mathcal{H}^p(G,M),\mathcal{H}^{d-p+1}(G',M'))\to\mathbb{Z}_1\end{aligned}\tag{E9}$$

If both M and M' are R-free, then the sequence splits and we have

$$\begin{aligned}&\mathcal{H}^d(G\times G',M\otimes_R M')\\ &=\Big[\prod_{p=0}^{d}\mathcal{H}^p(G,M)\otimes_R\mathcal{H}^{d-p}(G',M')\Big]\times\\ &\quad\Big[\prod_{p=0}^{d+1}\mathrm{Tor}_1^R(\mathcal{H}^p(G,M),\mathcal{H}^{d-p+1}(G',M'))\Big].\end{aligned}\tag{E10}$$

If R is a field K, we have

$$\begin{aligned}&\mathcal{H}^d(G\times G',M\otimes_K M')\\ &=\prod_{p=0}^{d}\Big[\mathcal{H}^p(G,M)\otimes_K\mathcal{H}^{d-p}(G',M')\Big].\end{aligned}\tag{E11}$$

Let us choose $R=\mathbb{Z}$, $M=\mathbb{Z}$ and $M'=\mathbb{Z}_2$. Note that $M=\mathbb{Z}$ is a free module over $R=\mathbb{Z}$ but $M'=\mathbb{Z}_2$ is not a free module over $R=\mathbb{Z}$. We also note that $M\otimes_\mathbb{Z}M'=\mathbb{Z}\otimes_\mathbb{Z}\mathbb{Z}_2=\mathbb{Z}_2$. Let us choose $G'=Z_1$ the trivial group with only identity. We have

$$\mathcal{H}^d(Z_1,\mathbb{Z}_2)=\begin{cases}\mathbb{Z}_2 & \text{if } d=0,\\ \mathbb{Z}_1 & \text{if } d>0.\end{cases}\tag{E12}$$

Thus we obtain

$$\begin{aligned}&\mathbb{Z}_1\to\mathcal{H}^d(G,\mathbb{Z})\otimes_\mathbb{Z}\mathbb{Z}_2\to\mathcal{H}^d(G,\mathbb{Z}_2)\\ &\quad\to\mathrm{Tor}_1^\mathbb{Z}(\mathcal{H}^{d+1}(G,\mathbb{Z}),\mathbb{Z}_2)\to\mathbb{Z}_1.\end{aligned}\tag{E13}$$

The above can be calculated using the following simple properties of the tensor product $\otimes_\mathbb{Z}$ and $\mathrm{Tor}_1^\mathbb{Z}$ functor:

$$\begin{aligned}&\mathbb{Z}\otimes_\mathbb{Z}M=M\otimes_\mathbb{Z}\mathbb{Z}=M,\\ &\mathbb{Z}_n\otimes_\mathbb{Z}M=M\otimes_\mathbb{Z}\mathbb{Z}_n=M/nM,\\ &\mathbb{Z}_m\otimes_\mathbb{Z}\mathbb{Z}_n=\mathbb{Z}_{(m,n)},\\ &(A\times B)\otimes_R M=(A\otimes_R M)\times(B\otimes_R M),\\ &M\otimes_R(A\times B)=(M\otimes_R A)\times(M\otimes_R B);\end{aligned}\tag{E14}$$

$$\begin{aligned}&\mathrm{Tor}_1^\mathbb{Z}(\mathbb{Z},M)=\mathrm{Tor}_1^\mathbb{Z}(M,\mathbb{Z})=\mathbb{Z}_1,\\ &\mathrm{Tor}_1^\mathbb{Z}(\mathbb{Z}_n,M)=\mathrm{Tor}_1^\mathbb{Z}(M,\mathbb{Z}_n)=M/nM,\\ &\mathrm{Tor}_1^\mathbb{Z}(\mathbb{Z}_m,\mathbb{Z}_n)=\mathbb{Z}_{(m,n)},\\ &\mathrm{Tor}_1^R(A\times B,M)=\mathrm{Tor}_1^R(A,M)\times\mathrm{Tor}_1^R(B,M),\\ &\mathrm{Tor}_1^R(M,A\times B)=\mathrm{Tor}_1^R(M,A)\times\mathrm{Tor}_1^R(M,B).\end{aligned}\tag{E15}$$

Here (m,n) is the greatest common divisor of m and n.

Using

$$\mathcal{H}^d(Z_n,\mathbb{Z})=\begin{cases}\mathbb{Z} & \text{if } d=0,\\ \mathbb{Z}_n & \text{if } d=0 \bmod 2,\\ \mathbb{Z}_1 & \text{if } d=1 \bmod 2,\end{cases}\tag{E16}$$

we find that for $d=$ even

$$\begin{aligned}&\mathbb{Z}_1 \to \mathbb{Z}_n \otimes_{\mathbb{Z}} \mathbb{Z}_2 \to \mathcal{H}^d(\mathbb{Z}_n, \mathbb{Z}_2) \to \mathrm{Tor}_1^{\mathbb{Z}}(\mathbb{Z}_1, \mathbb{Z}_2) \to \mathbb{Z}_1,\\ &\text{or } \mathbb{Z}_1 \to \mathbb{Z}_{(n,2)} \to \mathcal{H}^d(\mathbb{Z}_n, \mathbb{Z}_2) \to \mathbb{Z}_1 \to \mathbb{Z}_1; \qquad \text{(E17)}\end{aligned}$$

and for $d=$odd

$$\begin{aligned}&\mathbb{Z}_1 \to \mathbb{Z}_1 \otimes_{\mathbb{Z}} \mathbb{Z}_2 \to \mathcal{H}^d(\mathbb{Z}_n, \mathbb{Z}_2) \to \mathrm{Tor}_1^{\mathbb{Z}}(\mathbb{Z}_n, \mathbb{Z}_2) \to \mathbb{Z}_1,\\ &\text{or } \mathbb{Z}_1 \to \mathbb{Z}_1 \to \mathcal{H}^d(\mathbb{Z}_n, \mathbb{Z}_2) \to \mathbb{Z}_{(n,2)} \to \mathbb{Z}_1. \qquad \text{(E18)}\end{aligned}$$

This allows us to obtain eqn. (E7) using a different approach.

Using

$$\mathcal{H}^d[Z_n \rtimes Z_2, \mathbb{Z}] \leq \begin{cases} \mathbb{Z}, & d=0,\\ \mathbb{Z}_2 \times \mathbb{Z}_n \times \mathbb{Z}_{(2,n)}^{\frac{d-2}{2}}, & d=0 \text{ mod } 4,\\ \mathbb{Z}_{(2,n)}^{\frac{d-1}{2}}, & d=1 \text{ mod } 2,\\ \mathbb{Z}_2 \times \mathbb{Z}_{(2,n)}^{\frac{d}{2}}, & d=2 \text{ mod } 4, \end{cases} \qquad \text{(E19)}$$

we can show that for $d=0$ mod 4, $d>0$,

$$\begin{aligned}&\mathbb{Z}_1 \to (\mathbb{Z}_2 \times \mathbb{Z}_n \times \mathbb{Z}_{(2,n)}^{\frac{d-2}{2}}) \otimes_{\mathbb{Z}} \mathbb{Z}_2 \to \mathcal{H}^d[Z_n \rtimes Z_2, \mathbb{Z}_2]\\ &\quad \to \mathrm{Tor}_1^{\mathbb{Z}}(\mathbb{Z}_{(2,n)}^{\frac{d}{2}}, \mathbb{Z}_2) \to \mathbb{Z}_1, \qquad \text{(E20)}\\ &\text{or } \mathbb{Z}_1 \to \mathbb{Z}_2 \times \mathbb{Z}_{(2,n)}^{\frac{d}{2}} \to \mathcal{H}^d[Z_n \rtimes Z_2, \mathbb{Z}_2] \to \mathbb{Z}_{(2,n)}^{\frac{d}{2}} \to \mathbb{Z}_1;\end{aligned}$$

for $d=1$ mod 4, $d>0$,

$$\begin{aligned}&\mathbb{Z}_1 \to \mathbb{Z}_{(2,n)}^{\frac{d-1}{2}} \otimes_{\mathbb{Z}} \mathbb{Z}_2 \to \mathcal{H}^d[Z_n \rtimes Z_2, \mathbb{Z}_2]\\ &\quad \to \mathrm{Tor}_1^{\mathbb{Z}}(\mathbb{Z}_2 \times \mathbb{Z}_{(2,n)}^{\frac{d+1}{2}}, \mathbb{Z}_2) \to \mathbb{Z}_1, \qquad \text{(E21)}\\ &\text{or } \mathbb{Z}_1 \to \mathbb{Z}_{(2,n)}^{\frac{d-1}{2}} \to \mathcal{H}^d[Z_n \rtimes Z_2, \mathbb{Z}_2] \to \mathbb{Z}_2 \times \mathbb{Z}_{(2,n)}^{\frac{d+1}{2}} \to \mathbb{Z}_1;\end{aligned}$$

for $d=2$ mod 4, $d>0$,

$$\begin{aligned}&\mathbb{Z}_1 \to (\mathbb{Z}_2 \times \mathbb{Z}_{(2,n)}^{\frac{d}{2}}) \otimes_{\mathbb{Z}} \mathbb{Z}_2 \to \mathcal{H}^d[Z_n \rtimes Z_2, \mathbb{Z}_2]\\ &\quad \to \mathrm{Tor}_1^{\mathbb{Z}}(\mathbb{Z}_{(2,n)}^{\frac{d}{2}}, \mathbb{Z}_2) \to \mathbb{Z}_1, \qquad \text{(E22)}\\ &\text{or } \mathbb{Z}_1 \to \mathbb{Z}_2 \times \mathbb{Z}_{(2,n)}^{\frac{d}{2}} \to \mathcal{H}^d[Z_n \rtimes Z_2, \mathbb{Z}_2] \to \mathbb{Z}_{(2,n)}^{\frac{d}{2}} \to \mathbb{Z}_1;\end{aligned}$$

for $d=3$ mod 4, $d>0$,

$$\begin{aligned}&\mathbb{Z}_1 \to \mathbb{Z}_{(2,n)}^{\frac{d-1}{2}} \otimes_{\mathbb{Z}} \mathbb{Z}_2 \to \mathcal{H}^d[Z_n \rtimes Z_2, \mathbb{Z}_2]\\ &\quad \to \mathrm{Tor}_1^{\mathbb{Z}}(\mathbb{Z}_2 \times \mathbb{Z}_n \times \mathbb{Z}_{(2,n)}^{\frac{d-1}{2}}, \mathbb{Z}_2) \to \mathbb{Z}_1, \qquad \text{(E23)}\\ &\text{or } \mathbb{Z}_1 \to \mathbb{Z}_{(2,n)}^{\frac{d-1}{2}} \to \mathcal{H}^d[Z_n \rtimes Z_2, \mathbb{Z}_2] \to \mathbb{Z}_2 \times \mathbb{Z}_{(2,n)}^{\frac{d+1}{2}} \to \mathbb{Z}_1.\end{aligned}$$

Combining the above results, we find that, for $d>0$,

$$\mathcal{H}^d[Z_n \rtimes Z_2, \mathbb{Z}_2] = \mathbb{Z}_2 \times \mathbb{Z}_{(2,n)}^d. \qquad \text{(E24)}$$

Using

$$\mathcal{H}^d[U(1), \mathbb{Z}] = \begin{cases} \mathbb{Z} & \text{if } d=0 \text{ mod } 2,\\ \mathbb{Z}_1 & \text{if } d=1 \text{ mod } 2, \end{cases} \qquad \text{(E25)}$$

we find that for $d=$ even

$$\begin{aligned}&\mathbb{Z}_1 \to \mathbb{Z} \otimes_{\mathbb{Z}} \mathbb{Z}_2 \to \mathcal{H}^d[U(1), \mathbb{Z}_2] \to \mathrm{Tor}_1^{\mathbb{Z}}(\mathbb{Z}_1, \mathbb{Z}_2) \to \mathbb{Z}_1,\\ &\text{or } \mathbb{Z}_1 \to \mathbb{Z}_2 \to \mathcal{H}^d[U(1), \mathbb{Z}_2] \to \mathbb{Z}_1 \to \mathbb{Z}_1; \qquad \text{(E26)}\end{aligned}$$

and for $d=$odd

$$\begin{aligned}&\mathbb{Z}_1 \to \mathbb{Z}_1 \otimes_{\mathbb{Z}} \mathbb{Z}_2 \to \mathcal{H}^d[U(1), \mathbb{Z}_2] \to \mathrm{Tor}_1^{\mathbb{Z}}(\mathbb{Z}, \mathbb{Z}_2) \to \mathbb{Z}_1,\\ &\text{or } \mathbb{Z}_1 \to \mathbb{Z}_1 \to \mathcal{H}^d(U(1), \mathbb{Z}_2) \to \mathbb{Z}_1 \to \mathbb{Z}_1. \qquad \text{(E27)}\end{aligned}$$

This allows us to obtain

$$\mathcal{H}^d[U(1), \mathbb{Z}_2] = \begin{cases} \mathbb{Z}_2 & \text{if } d=0 \text{ mod } 2,\\ \mathbb{Z}_1 & \text{if } d=1 \text{ mod } 2, \end{cases} \qquad \text{(E28)}$$

Note that in $\mathcal{H}^d[U(1), \mathbb{Z}]$ and $\mathcal{H}^d[U(1), \mathbb{Z}_2]$, the cocycles $\nu_d : [U(1)]^{d+1} \to \mathbb{Z}$ or $\nu_d : [U(1)]^{d+1} \to \mathbb{Z}_2$ are Borel measurable functions.

Let us choose $R = M = M' = \mathbb{Z}_2$. Since R is a field and $\mathbb{Z}_2 \otimes_{\mathbb{Z}_2} \mathbb{Z}_2$, we have

$$\mathcal{H}^d(G \times G', \mathbb{Z}_2) = \prod_{p=0}^{d} \Big[\mathcal{H}^p(G, \mathbb{Z}_2) \otimes_{\mathbb{Z}_2} \mathcal{H}^{d-p}(G', \mathbb{Z}_2)\Big]. \qquad \text{(E29)}$$

Note that

$$\mathbb{Z}_2 \otimes_{\mathbb{Z}_2} \mathbb{Z}_2 = \mathbb{Z}_2, \quad \mathbb{Z}_2 \otimes_{\mathbb{Z}_2} \mathbb{Z}_1 = \mathbb{Z}_1. \qquad \text{(E30)}$$

We find

$$\mathcal{H}^d[Z_2 \times Z_2, \mathbb{Z}_2] = \mathbb{Z}_2^{d+1}. \qquad \text{(E31)}$$

Appendix F: Calculate the group super-cohomology classes $\mathscr{H}^d[G_f, U_T(1)]$

1. A general discussion

From the discussions in Appendix C, we see that to calculate $\mathscr{H}^d[G_f, U_T(1)]$ (which is formed by the equivalent classes of the fermionic cocycles $[\nu_d(g_0,...,g_d), n_{d-1}(g_0,...,g_{d-1})]$), we need to go through the following steps.

First step: Calculate $\mathcal{H}^{d-1}(G_b, \mathbb{Z}_2)$ which give us different classes of dD-graded structure $n_{d-1}(g_0,...,g_{d-1})$. For each $n_{d-1}(g_0,...,g_{d-1})$, find an allowed $u^g_{d-1}(g_0,...,g_{d-1})$.

Second step: Next we need to find out $B\mathcal{H}^{d-1}(G_b, \mathbb{Z}_2)$ from $\mathcal{H}^{d-1}(G_b, \mathbb{Z}_2)$. We note that for each graded structure $n_{d-1}(g_0,...,g_{d-1})$ in $\mathcal{H}^{d-1}(G_b, \mathbb{Z}_2)$, the corresponding $\frac{1}{2}\boldsymbol{f}_{d+1}$ is a $(d+1)$ is a cochain in $\mathcal{C}^{d+1}[G_b, U_T(1)]$. Then, $B\mathcal{H}^{d-1}(G_b, \mathbb{Z}_2)$ denote the subset of $\mathcal{H}^{d-1}(G_b, \mathbb{Z}_2)$, such that for each graded structure $n_{d-1}(g_0,...,g_{d-1})$ in $B\mathcal{H}^{d-1}(G_b, \mathbb{Z}_2)$, the corresponding $\frac{1}{2}\boldsymbol{f}_{d+1}$ is a $(d+1)$-coboundary in $\mathcal{B}^{d+1}[G_b, U_T(1)]$. Thus we have

$$\begin{aligned}&\frac{1}{2}\boldsymbol{f}_{d+1} = D_d \boldsymbol{g}_d \text{ mod } 1,\\ &\text{for all } n_{d-1}(g_0,...,g_{d-1}) \in B\mathcal{H}^{d-1}(G_b, \mathbb{Z}_2), \qquad \text{(F1)}\end{aligned}$$

where $\boldsymbol{f}_{d+1}$ is obtained from $n_{d-1}(g_0,...,g_{d-1})$ through eqn. (48). We note that for $n_{d-1}(g_0,...,g_{d-1}) \notin B\mathcal{H}^{d-1}(G_b,\mathbb{Z}_2)$, the corresponding $\boldsymbol{f}_d$ will not be a coboundary in $\mathcal{B}^{d+1}[G_b,U_T(1)]$ and eqn. (C17) will not have any solution.

Third step: We need to calculate $\nu_d(g_0,...,g_d)$ from eqn. (C17). For each fixed $n_{d-1}(g_0,...,g_{d-1}) \in B\mathcal{H}^{d-1}(G_b,\mathbb{Z}_2)$, eqn. (C17) is reduced to

$$D_d(\boldsymbol{v}_d+\boldsymbol{g}_d)=0 \text{ mod } 1. \tag{F2}$$

The solutions $\boldsymbol{v}_d$ of eqn. (F2) have a one-to-one correspondence with the cocycles in $\mathcal{Z}^d[G_b,U_T(1)]$, provided that $u^g_{d-1}(g_0,...,g_{d-1})=1$ for $g\in G_b$.

Fourth step: From the equivalence relation $\boldsymbol{v}_d \sim \boldsymbol{v}_d + D_{d-1}\boldsymbol{v}_{d-1}+\frac{1}{2}\boldsymbol{f}_d$, we see that the equivalent classes of the solutions can be labeled by $\mathcal{H}^d[G_b,U_T(1)]$. However, the labeling is not one-to-one, since in obtaining $\mathcal{H}^d[G_b,U_T(1)]$, we only used the equivalence relation $\boldsymbol{v}_d \sim \boldsymbol{v}_d + D_{d-1}\boldsymbol{v}_{d-1}$. The different classes in $\mathcal{H}^d[G_b,U_T(1)]$ differ by $\frac{1}{2}\boldsymbol{f}_d$ are regarded as equivalent under the group super-cohomology. The labeling is one-to-one only when $\frac{1}{2}\boldsymbol{f}_d$ happen to be coboundaries in $\mathcal{B}^d[G_b,U_T(1)]$.

Through the above four steps, and combining with results from the standard group cohomology, we can calculate the group super-cohomology classes $\mathscr{H}^d[G_f,U_T(1)]$. We like to point out that a subset of $\mathscr{H}^d[G_f,U_T(1)]$ that has trivial dD-graded structure $n_{d-1}(g_0,...,g_{d-1})=0$ is given by $\mathcal{H}^d[G_f,U_T(1)]$. Therefore, the fermionic SPT phases contain the bosonic SPT phases with symmetry G_b as special cases. In the following, we will discuss some simple examples.

2. Calculate $\mathscr{H}^d[Z_1\times Z_2^f,U(1)]$

Here we choose $G_b=Z_1$ (the trivial group) which correspond to fermion systems with no symmetry. Since $\mathcal{H}^{d-1}(Z_1,\mathbb{Z}_2)=\mathbb{Z}_2$ for $d=1$ and $\mathcal{H}^{d-1}(Z_1,\mathbb{Z}_2)=\mathbb{Z}_1=0$ for $d>1$. We find that there are two 1D-graded structures and only one trivial dD-graded structure for $d>1$. For each dD-graded structure, we can choose a $u^g_{d-1}(g_0,...,g_{d-1})$ that satisfies $u^g_{d-1}(g_0,...,g_{d-1})=1$ for $g\in G_b$.

For $d>1$ and for the only trivial graded structure, the equivalent classes of $\boldsymbol{v}_d$ are described by $\mathcal{H}^d[Z_1,U(1)]=\mathbb{Z}_1$. Therefore $\mathscr{H}^d[Z_1\times Z_2^f,U(1)]=\mathbb{Z}_1$ for $d>1$.

For $d=1$, $\mathcal{H}^0(Z_1,\mathbb{Z}_2)=\mathbb{Z}_2$. The trivial graded structure is given by $n^{(0)}(E)=0$ and the other non-trivial graded structure is given by $n^{(1)}(E)=1$. The corresponding $f_2^{(a)}(g_0,...,g_d)$ are given by $f_2^{(a)}=0$, $a=0,1$. Thus for the each graded structure, the equivalent classes of $\boldsymbol{v}_d$ are described by $\mathcal{H}^1[Z_1,U(1)]=\mathbb{Z}_1$. Therefore $\mathscr{H}^1[Z_1\times Z_2^f,U(1)]=\mathbb{Z}_2$. To summarize

$$\mathscr{H}^d[Z_1\times Z_2^f,U(1)]=\begin{cases}\mathbb{Z}_2, & d=1,\\ \mathbb{Z}_1, & d>1.\end{cases} \tag{F3}$$

$\mathscr{H}^1[Z_1\times Z_2^f,U(1)]=\mathbb{Z}_2$ implies that there are two possible fermionic SPT phases in $d_{sp}=0$ spatial dimension if there is no symmetry. One has even number of fermions and the other has odd number of fermions.

$\mathscr{H}^2[Z_1\times Z_2^f,U(1)]=\mathbb{Z}_1$ implies that there is only one trivial gapped fermionic SPT phases in $d_{sp}=1$ spatial dimension if there is no symmetry. It is well known that there are two gapped fermionic phases in $d_{sp}=1$ spatial dimension if there is no symmetry, the trivial phase and the phase with boundary Majorana zero mode.[57] One way to see this result is to use the Jordan-Wigner transformation to map the 1D fermion system with no symmetry to an 1D boson system with Z_2^f symmetry. 1D boson system with Z_2^f symmetry can have only two gapped phase: the symmetry unbroken one (which corresponds to the trivial fermionic phase) and the symmetry breaking one (which corresponds to the Majorana chain).[48] Both gapped phases can be realized by non-interacting fermions (see Table II). However, the non-trivial gapped phase (the bosonic Z_2^f symmetry breaking phase) has non-trivial intrinsic topological orders. So the non-trivial gapped phase is not a fermionic SPT phase. This is why we only has one trivial fermionic SPT phase in 1D if there is no symmetry.

$\mathscr{H}^3[Z_1\times Z_2^f,U(1)]=\mathbb{Z}_1$ implies that there is only one trivial gapped fermionic SPT phases in $d_{sp}=2$ spatial dimension if there is no symmetry. It is well known that even non-interacting fermions have infinite gapped phases labeled by $\mathbb{Z}$ in $d_{sp}=2$ spatial dimensions if there is no symmetry (see table II). Those phases correspond to integer quantum Hall states and/or $p+\mathrm{i}p$, $d+\mathrm{i}d$ superconductors. Again, all those gapped phases have non-trivial intrinsic topological orders. So we only have one trivial fermionic SPT phase in $d_{sp}=2$ spatial dimensions if there is no symmetry.

3. Calculate $\mathscr{H}^d[Z_{2k+1}\times Z_2^f,U(1)]$

Next we choose $G_f=Z_{2k+1}\times Z_2^f$. Again, since $\mathcal{H}^{d-1}(Z_{2k+1},\mathbb{Z}_2)=\mathbb{Z}_2$ for $d=1$ and $\mathcal{H}^{d-1}(Z_{2k+1},\mathbb{Z}_2)=\mathbb{Z}_1$ for $d>1$, there are two 1D-graded structures and only one trivial dD-graded structure for $d>1$. For each dD-graded structure, we can choose a $u^g_{d-1}(g_0,...,g_{d-1})$ that satisfies $u^g_{d-1}(g_0,...,g_{d-1})=1$ for $g\in G_b$.

For $d>1$ and for the only trivial graded structure, the equivalent classes of $\boldsymbol{v}_d$ are described by $\mathcal{H}^d[Z_{2k+1},U(1)]=\mathbb{Z}_1$ for d=even and $\mathcal{H}^d[Z_{2k+1},U(1)]=\mathbb{Z}_{2k+1}$ for d=odd. Therefore $\mathscr{H}^d[Z_1\times Z_2^f,U(1)]=\mathbb{Z}_1$ for d=even and $\mathscr{H}^d[Z_1\times Z_2^f,U(1)]=\mathbb{Z}_{2k+1}$ for d=odd.

For $d=1$, $\mathcal{H}^0(Z_{2k+1},\mathbb{Z}_2)=\mathbb{Z}_2$. The trivial graded structure is given by $n^{(0)}(g_0)=0$ and the other non-trivial graded structure is given by $n^{(1)}(g_0)=1$. The corresponding $f_2^{(a)}(g_0,...,g_d)$ are given by $f_2^{(a)}=0$, $a=0,1$. Thus for the each graded structure, the equivalent classes of $\boldsymbol{v}_d$ are described by $\mathcal{H}^1[Z_{2k+1},U(1)]=\mathbb{Z}_{2k+1}$. Therefore $\mathscr{H}^1[Z_{2k+1}\times Z_2^f,U(1)]=\mathbb{Z}_2\times\mathbb{Z}_{2k+1}=\mathbb{Z}_{4k+2}$.

To summarize

$$\mathscr{H}^d[Z_{2k+1}\times Z_2^f, U(1)] = \begin{cases}\mathbb{Z}_{4k+2}, & d=1,\\ \mathbb{Z}_1, & d=\text{even}, d>0,\\ \mathbb{Z}_{2k+1}, & d=\text{odd}.\end{cases} \tag{F4}$$

$\mathscr{H}^1[Z_{2k+1}\times Z_2^f, U(1)] = \mathbb{Z}_{2k+1}\times\mathbb{Z}_2$ implies that there are $4k+2$ possible fermionic SPT phases in $d_{sp}=0$ spatial dimension if there is a Z_{2k+1} symmetry. $2k+1$ of them have even number of fermions and the other $2k+1$ of them have odd number of fermions. The $2k+1$ phases are separated by the $2k+1$ possible Z_{2k+1} quantum numbers.

$\mathscr{H}^2[Z_{2k+1}\times Z_2^f, U(1)] = \mathbb{Z}_1$ implies that there is no non-trivial gapped fermionic SPT phases in $d_{sp}=1$ spatial dimension if there is a Z_{2k+1} symmetry. We can use the Jordan-Wigner transformation to map 1D fermion systems with Z_{2k+1} symmetry to 1D boson systems with $Z_{2k+1}\times Z_2^f = Z_{4k+2}$ symmetry. Since $\mathcal{H}^2[Z_{4k+2}, U(1)] = \mathbb{Z}_1$, 1D boson systems have only two phases that do not break the Z_{2k+1} symmetry: one does not break the Z_2^f symmetry and one breaks the Z_2^f symmetry. The fermionic SPT phase has no intrinsic topological order and correspond to the bosonic phase that does not break the Z_2^f symmetry (and the Z_{2k+1} symmetry). Thus 1D fermion systems with Z_{2k+1} symmetry indeed have only one trivial fermionic SPT phase.

$\mathscr{H}^3[Z_{2k+1}\times Z_2^f, U(1)] = \mathbb{Z}_{2k+1}$ implies that our construction gives rise to $2k+1$ gapped fermionic SPT phases in $d_{sp}=2$ spatial dimension if there is a Z_{2k+1} symmetry. The constructed $2k+1$ gapped fermionic SPT phases actually correspond to the $2k+1$ gapped bosonic SPT phases with Z_{2k+1} symmetry [since $\mathcal{H}^2(Z_{2k+1}, \mathbb{Z}_2) = \mathbb{Z}_1$ and there is no non-trivial graded structure].

4. Calculate $\mathscr{H}^d[Z_2\times Z_2^f, U(1)]$

Now, let us choose $G_f = Z_2\times Z_2^f$. Following the first step in section F 1, we find that $\mathcal{H}^{d-1}(Z_2,\mathbb{Z}_2) = \mathbb{Z}_2$ and there are two graded structures in all dimensions. For each dD-graded structure, we can choose a $u^g_{d-1}(g_0,...,g_{d-1})$ that satisfies $u^g_{d-1}(g_0,...,g_{d-1}) = 1$ for $g\in G_b$.

Following the second step in section F 1, we note that the trivial graded structure is given by $n_{d-1}(g_0,...,g_{d-1}) = 0$. The non-trivial graded structure is given by

$$n_{d-1}(e,o,e,...) = n_{d-1}(o,e,o,...) = 1, \quad \text{others} = 0, \tag{F5}$$

where e represents the identity element in Z_2 and o represents the other element in Z_2.

For $d=1,2$, $\mathcal{H}^{d-1}(Z_2,\mathbb{Z}_2) = \mathbb{Z}_2$. For each graded structure $n_{d-1}(g_0,...,g_{d-1})$, the correspond $f_{d+1}(g_0,...,g_{d+1}) = 0$. Thus $B\mathcal{H}^{d-1}(Z_2,\mathbb{Z}_2) = \mathbb{Z}_2$.

For $d=3$, $\mathcal{H}^2(Z_2,\mathbb{Z}_2) = \mathbb{Z}_2$. For the non-trivial 3D-graded structure, $n_2(e,o,e) = n_2(o,e,o) = 1$ and others $n_2 = 0$, the corresponding f_4 has a form

$$f_4(e,o,e,o,e) = f_4(o,e,o,e,o) = 1, \quad \text{others} = 0. \tag{F6}$$

One can show that $(-)^{f_4(g_0,\cdots,g_4)}$ is a cocycle in $\mathcal{Z}^4[Z_2, U(1)]$. Since $\mathcal{H}^4[Z_2, U(1)] = \mathbb{Z}_1$, $(-)^{f_4(g_0,\cdots,g_4)}$ is also a coboundary in $\mathcal{B}^4[Z_2, U(1)]$. Therefore, $B\mathcal{H}^2(Z_2,\mathbb{Z}_2) = \mathbb{Z}_2$.

For $d=4$, $\mathcal{H}^3(Z_2,\mathbb{Z}_2) = \mathbb{Z}_2$. For the non-trivial 4D-graded structure, $n_3(e,o,e,o) = n_3(o,e,o,e) = 1$ and others $n_3 = 0$, the corresponding f_5 has a form

$$f_5(e,o,e,o,e,o) = f_5(o,e,o,e,o,e) = 1, \quad \text{others} = 0. \tag{F7}$$

One can show that $(-)^{f_5(g_0,\cdots,g_5)}$ is a cocycle in $\mathcal{Z}^5[Z_2, U(1)]$. Since $\mathcal{H}^5[Z_2, U(1)] = \mathbb{Z}_2$ and $(-)^{f_5(g_0,\cdots,g_5)}$ corresponds to a non-trivial cocycle in $\mathcal{H}^5[Z_2, U(1)] = \mathbb{Z}_2$, we find that $B\mathcal{H}^3(Z_2,\mathbb{Z}_2) = \mathbb{Z}_1$.

Following the third step in section F 1, we can show that the elements in $\mathscr{H}^d[Z_2\times Z_2^f, U(1)]$ can be labeled by $\mathcal{H}^d[Z_2\times Z_2^f, U(1)]\times B\mathcal{H}^{d-1}(Z_2,\mathbb{Z}_2)$.

Following the fourth step in section F 1, we like to show that the above labeling is one-to-one. For $d=1,2,3$, $f_d(g_0,...,g_d) = 0$. Thus the labeling is one-to-one. For $d=4$, a 3D-graded structure in $B\mathcal{H}^2(Z_2,\mathbb{Z}_2)$ gives rise to a $f_4(g_0,...,g_4)$. Since $(-)^{f_4(g_0,\cdots,g_4)}$ is a cocycle in $\mathcal{Z}^4[Z_2, U(1)]$ and since $\mathcal{H}^4[Z_2, U(1)] = \mathbb{Z}_1$, we find that $(-)^{f_4(g_0,\cdots,g_4)}$ is also a coboundary in $\mathcal{B}^4[Z_2, U(1)]$ and the labeling is one-to-one.

So, using

$$\mathcal{H}^d[Z_n, U(1)] = \begin{cases}\mathbb{Z}_1, & d = 0 \bmod 2,\ d>0,\\ \mathbb{Z}_n, & d = 1 \bmod 2,\end{cases} \tag{F8}$$

we find that[50]

$$\mathscr{H}^d[Z_2\times Z_2^f, U(1)] = \begin{cases}\mathbb{Z}_2^2, & d=1,\\ \mathbb{Z}_2, & d=2,\\ \mathbb{Z}_2^2, & d=3,\\ \mathbb{Z}_1, & d=4.\end{cases} \tag{F9}$$

$\mathscr{H}^1[Z_2\times Z_2^f, U(1)] = \mathbb{Z}_2^2$ implies that there are 4 possible fermionic SPT phases in $d_{sp}=0$ spatial dimension if there is a Z_2 symmetry. 2 of them have even number of fermions and the other 2 of them have odd number of fermions. The 2 phases with even number of fermions are separated by the 2 possible Z_2 quantum numbers. The 2 phases with odd number of fermions are also separated by the 2 possible Z_2 quantum numbers.

$\mathscr{H}^2[Z_2\times Z_2^f, U(1)] = \mathbb{Z}_2$ implies that there are two gapped fermionic SPT phases in $d_{sp}=1$ spatial dimension if there is a Z_2 symmetry. We can use the Jordan-Wigner transformation to map 1D fermion systems with Z_2 symmetry to 1D boson systems with $Z_2\times Z_2^f$ symmetry. Since $\mathcal{H}^2[Z_2\times Z_2, U(1)] = \mathbb{Z}_2$, 1D boson systems have two phases that do not break the $Z_2\times Z_2^f$ symmetry. 1D boson systems also have another phase that do not break the Z_2 symmetry: the phase that break the Z_2^f symmetry. The fermionic SPT phases have no intrinsic topological order and correspond to the bosonic phases that does not break the Z_2^f symmetry (and the Z_2 symmetry). Thus 1D fermion systems with Z_2 symmetry indeed have two fermionic SPT phases.

$\mathscr{H}^3[Z_2\times Z_2^f, U(1)] = \mathbb{Z}_2^2$ implies that our construction gives rise to 4 gapped fermionic SPT phases in $d_{sp}=2$ spatial dimension if there is a Z_2 symmetry.

5. Calculate $\mathscr{H}^d[Z_2^T\times Z_2^f, U_T(1)]$

Now, let us choose $G_f = Z_2^T\times Z_2^f$. Since $\mathcal{H}^{d-1}(Z_2^T,\mathbb{Z}_2)=\mathbb{Z}_2$, there are two graded structures in all dimensions. The trivial graded structure is given by $n_{d-1}(g_0,...,g_{d-1})=0$. The non-trivial graded structure is given by

$$n_{d-1}(e,o,e,...) = n_{d-1}(o,e,o,...) = 1, \quad \text{others} = 0, \tag{F10}$$

where e is the identity element in Z_2^T and o is the non-trivial element in Z_2^T. For each dD-graded structure, we can choose a $u^g_{d-1}(g_0,...,g_{d-1})$ that satisfies $u^g_{d-1}(g_0,...,g_{d-1})=1$ for $g\in G_b$.

For $d=1,2$, $\mathcal{H}^{d-1}(Z_2^T,\mathbb{Z}_2)=\mathbb{Z}_2$. For each graded structure $n_{d-1}(g_0,...,g_{d-1})$, the corresponding $f_{d+1}(g_0,...,g_{d+1})=0$, and the equivalent classes of $\boldsymbol{v}_d$ are described by $\mathcal{H}^d[Z_2^T,U_T(1)]$, Therefore $\mathscr{H}^1[Z_2^T\times Z_2^f,U_T(1)]=\mathbb{Z}_2$ and $\mathscr{H}^2[Z_2^T\times Z_2^f,U_T(1)]=\mathbb{Z}_2^2$.

For $d=3$, $\mathcal{H}^2(Z_2^T,\mathbb{Z}_2)=\mathbb{Z}_2$. For the non-trivial 3D-graded structure, $n_2(e,o,e)=n_2(o,e,o)=1$ and others $n_2=0$, the corresponding f_4 has a form

$$f_4(e,o,e,o,e) = f_4(o,e,o,e,o) = 1, \quad \text{others} = 0. \tag{F11}$$

One can show that $(-)^{f_4(g_0,...,g_4)}$ is a cocycle in $\mathcal{Z}^4[Z_2^T,U_T(1)]$. Since $\mathcal{H}^4[Z_2^T,U_T(1)]=\mathbb{Z}_2$ and $(-)^{f_4(g_0,...,g_4)}$ is a non-trivial cocycle in $\mathcal{H}^4[Z_2^T,U_T(1)]$, so for the non-trivial 3D-graded structure, eqn. (C17) for $\boldsymbol{v}_d$ has no solutions. For the trivial 3D-graded structure, the equivalent classes of $\boldsymbol{v}_d$ are described by $\mathcal{H}^3[Z_2^T,U_T(1)]$. Thus $\mathscr{H}^3[Z_2^T\times Z_2^f,U_T(1)]=\mathbb{Z}_1$.

For $d=4$, $\mathcal{H}^3(Z_2^T,\mathbb{Z}_2)=\mathbb{Z}_2$. For the non-trivial 4D-graded structure, $n_3(e,o,e,o)=n_3(o,e,o,e)=1$ and others $n_3=0$, the corresponding f_5 has a form

$$f_5(e,o,e,o,e,o) = f_5(o,e,o,e,o,e) = 1, \quad \text{others} = 0. \tag{F12}$$

One can show that $(-)^{f_5(g_0,...,g_5)}$ is a cocycle in $\mathcal{Z}^5[Z_2^T,U_T(1)]$. Since $\mathcal{H}^5[Z_2^T,U_T(1)]=\mathbb{Z}_1$, $(-)^{f_5(g_0,...,g_5)}$ is also a coboundary in $\mathcal{B}^5[Z_2^T,U_T(1)]$. So for the non-trivial 4D-graded structure, the equivalent classes of $\boldsymbol{v}_d$ are labeled by $\mathcal{H}^4[Z_2^T,U_T(1)]=\mathbb{Z}_2$. Since $(-)^{f_4(g_0,...,g_4)}$ is real and thus can be viewed as an elements in $\mathcal{H}^4[Z_2,U(1)]$. Since $\mathcal{H}^4[Z_2,U(1)]=\mathbb{Z}_1$, so $(-)^{f_4(g_0,...,g_4)}$ is always a coboundary in $\mathcal{B}^4[Z_2^T,U_T(1)]$, the labeling is one-to-one. For the trivial 4D-graded structure, the equivalent classes of $\boldsymbol{v}_d$ are labeled by $\mathcal{H}^4[Z_2^T,U_T(1)]=\mathbb{Z}_1$. Since $(-)^{f_4(g_0,...,g_4)}$ is always a coboundary in $\mathcal{B}^4[Z_2^T,U_T(1)]$, the labeling is one-to-one. Thus $\mathscr{H}^4[Z_2^T\times Z_2^f,U_T(1)]=\mathbb{Z}_2^2$. To summarize

$$\mathscr{H}^d[Z_2^T\times Z_2^f,U_T(1)] = \begin{cases} \mathbb{Z}_2, & d=1,\\ \mathbb{Z}_2^2, & d=2,\\ \mathbb{Z}_1, & d=3,\\ \mathbb{Z}_2^2, & d=4. \end{cases} \tag{F13}$$

$\mathcal{H}^1[Z_2^T\times Z_2^f,U_T(1)]=\mathbb{Z}_2$ implies that there are two possible fermionic SPT phases in $d_{sp}=0$ spatial dimension if there is a Z_2^T symmetry. One has even number of fermions and the other one has odd number of fermions.

$\mathcal{H}^2[Z_2^T\times Z_2^f,U_T(1)]=\mathbb{Z}_2^2$ implies that there is four gapped fermionic SPT phases in $d_{sp}=1$ spatial dimension if there is a Z_2^T symmetry. We can use the Jordan-Wigner transformation to map 1D fermion systems with Z_2^T symmetry to 1D boson systems with $Z_2^T\times Z_2^f$ symmetry. Since $\mathcal{H}^2[Z_2^T\times Z_2,U_T(1)]=\mathbb{Z}_2^2$, 1D boson systems have four phases that do not break the $Z_2^T\times Z_2^f$ symmetry. 1D boson systems also have four phases that do break the Z_2^f symmetry: two do not break the time-reversal T symmetry and two do not break the TP_f symmetry. The fermionic SPT phases have no intrinsic topological order and correspond to the bosonic phases that do not break the Z_2^f symmetry (and the Z_2^T symmetry). Thus 1D fermion systems with Z_2^T symmetry indeed have four fermionic SPT phases.

$\mathcal{H}^3[Z_2^T\times Z_2^f,U_T(1)]=\mathbb{Z}_1$ implies that our construction only gives rise to one trivial gapped fermionic SPT phases in $d_{sp}=2$ spatial dimension if there is a Z_2^T symmetry. $\mathcal{H}^4[Z_2^T\times Z_2^f,U_T(1)]=\mathbb{Z}_2^2$ implies that our construction gives rise to three non-trivial gapped fermionic SPT phases in $d_{sp}=3$ spatial dimension if there is a Z_2^T symmetry. One of the non-trivial SPT phases is actially the purely bosonic SPT phase. The other two are intrinsically fermionic. We note that, using non-interacting fermions, we cannot construct any non-trivial gapped phases with Z_2^T symmetry (the $T^2=1$ time-reversal symmetry).

Appendix G: Properties of the constructed SPT states

1. Equivalence between ground state wave functions

In Section VI, we have demonstrated how to construct an ideal ground state wave function $\Phi(\{g_i\},\{n_{ij...k}\})$ in d spatial dimension from a $(d+1)$-cocycle (ν_{d+1},n_d,u^g_d).

However, a natural question is: how do we know these wave functions describe the same fermionic SPT phases or not? Here we will address this important question based on the new mathematical structure - group super-cohomology class invented in Appendix C.

a. *A special equivalent relation*

Let us choose arbitrary $(\nu_d, n_{d-1}, u^g_{d-1})$ in one lower dimension. Here $(\nu_d, n_{d-1}, u^g_{d-1})$ is not a d-cocycle. They just satisfy the symmetry condition eqn. (46). In Appendix C, we have shown that $(\tilde{\nu}_{d+1}, \tilde{n}_d, \tilde{u}^g_d)$ obtained from (ν_{d+1}, n_d, u^g_d) through eqn. (C23) is also a $(d+1)$-cocycle. Here we will assume that $n_{d-1} = 0$ and $u^g_{d-1} = 1$, and the resulting

$$\tilde{\nu}_{d+1} = \nu_{d+1}\delta\nu_d, \quad \tilde{n}_d = n_d, \quad \tilde{u}^g_d = u^g_d, \tag{G1}$$

is a $(d{+}1)$-cocycle. The new $(d{+}1)$-cocycle $(\tilde{\nu}_{d+1}, \tilde{n}_d, \tilde{u}^g_d)$ gives rise to a new ideal wave function $\Phi'(\{g_i\}, \{n_{ij...k}\})$. Should we view the two wave functions $\Phi(\{g_i\}, \{n_{ij...k}\})$ and $\Phi'(\{g_i\}, \{n_{ij...k}\})$ as wave functions for different SPT phases?

From the definition of $\delta\nu_d$ (49), we can show that the two wave functions $\Phi(\{g_i\}, \{n_{ijk}\})$ and $\Phi'(\{g_i\}, \{n_{ijk}\})$ are related through

$$\begin{aligned}&\Phi'(\{g_i\}, \{n_{ijk}\}) = \Phi(\{g_i\}, \{n_{ijk}\})\times \\ &\prod_{(ij...k)} (-)^{f_{d+1}(g_0,g_i,g_j,...,g_k)} \nu_d^{s_{ij...k}}(g_i, g_j, ..., g_k),\end{aligned} \tag{G2}$$

where $\prod_{(ij...k)}$ is the product over all the d-simplices that form the d dimensional space [in our (2+1)D example, $\prod_{(ij...k)}$ is the product over all triangles]. Also $s_{ij...k} = \pm 1$ depending on the orientation of the d-simplex $(ij...k)$. Since $\nu_d^{s_{ij...k}}(g_i, g_j, ..., g_k)$ is a pure $U(1)$ phase that satisfies $\nu_d^{s_{ij...k}}(gg_i, gg_j, ..., gg_k) = \nu_d^{s_{ij...k}}(g_i, g_j, ..., g_k)$, it represents a bosonic *symmetric* local unitary (LU) transformation. Therefore the new wave function Φ' induced by $(\nu_d \neq 0, n_{d-1} = 0, u^g_{d-1} = 1)$ belong to the same SPT phase as the original wave function Φ.

Also since f_{d+1} for $d < 3$ is always zero. Therefore, below 3 spatial dimensions, new wave function Φ' belongs to the same SPT phase as the original wave function Φ.

In 3 spatial dimensions and above ($d \geq 3$), if we choose $(\nu_d = 0, n_{d-1} \neq 0)$, the new induced wave function Φ' will differ from the original wave function Φ by a phase factor $\prod_{(ij...k)}(-)^{f_{d+1}(g_0,g_i,g_j,...,g_k)}$. $(-)^{f_{d+1}(g_0,g_i,g_j,...,g_k)}$ represents a LU transformation, and the two wave functions, Φ' and Φ, have the same intrinsic topological order. But since $(-)^{f_{d+1}(g_0,gg_i,gg_j,...,gg_k)} \neq (-)^{f_{d+1}(g_0,g_i,g_j,...,g_k)}$, $(-)^{f_{d+1}(g_0,g_i,g_j,...,g_k)}$ does not represent a bosonic symmetric LU transformation. So we do not know whether Φ' and Φ belong to the same fermionic SPT phase or not.

We would like to point out that in order for Φ' and Φ to belong to the same *fermionic* SPT phase, the two wave functions can only differ by a *fermionic* symmetric LU transformation (which is defined in Ref. 25). The *bosonic* symmetric LU transformations are a subset of *fermionic* symmetric LU transformations. So even though $(-)^{f_{d+1}(g_0,g_i,g_j,...,g_k)}$ is not a *bosonic* symmetric LU transformation, we do not know whether it is a *fermionic* symmetric LU transformation or not. The structure of group super-cohomology theory developed in Appendix C shows that $f_{d+1}(g_0, g_i, g_j, ..., g_k)$ is a d+1-cocycle with $\mathbb{Z}_2$ coefficient, suggesting the sign factor $(-)^{f_{d+1}(g_0,g_i,g_j,...,g_k)}$ can be generated by *fermionic* symmetric LU transformation.

b. *Generic equivalent relations*

In the above, we discussed a special case of $(\tilde{\nu}_{d+1}, \tilde{n}_d, \tilde{u}^g_d)$ where $\tilde{n}_d$ and $\tilde{u}^g_d$ are the same as n_d and u^g_d. However, the equivalent class of group super-cohomology has a more complicated structure. In Appendix C, we have shown that $(\tilde{\nu}_{d+1}, \tilde{n}_d, \tilde{u}^g_d)$ obtained from:

$$\tilde{\nu}_{d+1} = \nu_{d+1}(-)^{g_d}, \ \tilde{n}_d = n_d + \delta m'_{d-1}, \ \tilde{u}^g_d = u^g_d, \tag{G3}$$

(with $g_d(g_0, \cdots, g_d) = (\boldsymbol{g}_d)_{(g_0,\cdots,g_d)}$ mod 2) also belongs to the same equivalent class of group super-cohomology. We note that here the graded structure n_d changes into $\tilde{n}_d$. It would be much harder to understand why the corresponding new wave function still describes the same fermionic SPT phase. Let us consider the $2+1$D wave function on a sphere as a simple example. Since the wave function derived from the fixed point amplitude Eq. (30) is a fixed point wave function with zero correlation length, it is enough to only consider a minimal wave function with 4 sites on a sphere(labeled as $1, 2, 3, 4$). Such a minimal wave functions will contain 4 triangles 123, 124, 134 and 234. From the definition of $\tilde{\nu}_d$, we can show that the two wave functions $\Phi(\{g_i\}, \{n_{ijk}\})$ and $\tilde{\Phi}(\{g_i\}, \{\tilde{n}_{ijk}\})$ are related through $\tilde{\Phi}(\{g_i\}, \{n'_{ijk}\}) = (-)^{g_4(g_1,g_2,g_3,g_4)}\Phi(\{g_i\}, \{n_{ijk}\})$. To see this explicitly, let us construct the ground state wave function by adding one more point 0 inside the sphere. In this way, the wave function can be formally expressed as:

$$\begin{aligned}&\Psi(g_1, g_2, g_3, g_4, \theta_{234}, \theta_{124}, \bar{\theta}_{134}, \bar{\theta}_{123}) \\ =&(-)^{m_1(g_1,g_3)} \int \mathcal{V}_3^-(g_0, g_1, g_2, g_3)\mathcal{V}_3^+(g_0, g_1, g_2, g_4) \\ &\times \mathcal{V}_3^-(g_0, g_1, g_3, g_4)\mathcal{V}_3^+(g_0, g_2, g_3, g_4) \\ =&(-)^{m_1(g_1,g_3)}\mathcal{V}_3^+(g_1, g_2, g_3, g_4) \\ =&\nu_3(g_1, g_2, g_3, g_4)\times \\ &\theta_{234}^{n_2(g_2,g_3,g_4)}\theta_{124}^{n_2(g_1,g_2,g_4)}\bar{\theta}_{234}^{n_2(g_2,g_3,g_4)}\bar{\theta}_{123}^{n_2(g_1,g_2,g_3)}.\end{aligned} \tag{G4}$$

Again, the symbol $\int$ is defined as integrating over the internal Grassmann variables with respecting to the weights m_1 on the internal links(see Eq.(18)). Similarly, the wave function corresponding to a different solution

$(\tilde{\nu}_{d+1}, \tilde{n}_d, \tilde{u}_d^g)$ takes a form:

$$\begin{aligned}&\tilde{\Psi}(g_1, g_2, g_3, g_4, \theta_{234}, \theta_{124}, \bar{\theta}_{134}, \bar{\theta}_{123})\\ =&\nu_3(g_1, g_2, g_3, g_4)(-)^{g_4(g_1,g_2,g_3,g_4)}\times\\ &\theta_{234}^{n_2'(g_2,g_3,g_4)}\theta_{124}^{n_2'(g_1,g_2,g_4)}\bar{\theta}_{234}^{n_2'(g_2,g_3,g_4)}\bar{\theta}_{123}^{n_2'(g_1,g_2,g_3)}. \end{aligned} \tag{G5}$$

We note that $g_4(gg_1, gg_2, gg_3, gg_4) = g_4(g_1, g_2, g_3, g_4)$ implies the phase factor $(-)^{g_4(g_1,g_2,g_3,g_4)}$ is a *symmetric* LU transformation. However, we still do not know whether the two wave functions with different patterns of fermion number $\{\tilde{n}_{ijk}\}$ and $\{n_{ijk}\}$ describe the same fermionic SPT phase or not.

Indeed, we find that up to some *symmetric* phase factors, the state $|\Phi\rangle$ and $|\tilde{\Phi}\rangle$ can be related through a *symmetric* LU transformation $|\tilde{\Phi}\rangle \sim \hat{U}'|\Phi\rangle$, where

$$\begin{aligned}\hat{U}' = &\prod_{(0ij)} c_{(0ij)}^{m_1'(g_i,g_j)}\bar{c}_{(0ij)}^{m_1'(g_i,g_j)}\times\\ &\prod_{\triangle} c_{(0ij)}^{\dagger m_1'(g_i,g_j)} c_{(0jk)}^{\dagger m_1'(g_j,g_k)} \bar{c}_{(0ik)}^{\dagger m_1'(g_i,g_k)} \left(c_{(ijk)}^{\dagger}\right)^{\tilde{n}_2(g_i,g_j,g_k)-n_2(g_i,g_j,g_k)}\times\\ &\prod_{\triangledown} \left(c_{(ijk)}^{\dagger}\right)^{\tilde{n}_2(g_i,g_j,g_k)-n_2(g_i,g_j,g_k)} c_{(0ik)}^{\dagger m_1'(g_i,g_k)} \bar{c}_{(0jk)}^{\dagger m_1'(g_j,g_k)} \bar{c}_{(0ij)}^{\dagger m_1'(g_i,g_j)},\end{aligned} \tag{G6}$$

with $\tilde{n}_2(g_i, g_j, g_k) = n_2(g_i, g_j, g_k) + m_1'(g_i, g_j) + m_1'(g_j, g_k) + m_1'(g_k, g_i)$. We note that both n_2 and m_1' are invariant under symmetry transformation. Here $\left(c_{(ijk)}^{\dagger}\right)^{-1}$ is defined as $\left(c_{(ijk)}^{\dagger}\right)^{-1} = c_{(ijk)}$. $\sim$ means equivalent up to some symmetric sign factors which will arise when we reorder the ferminon creation/anihilation operators in $\hat{U}'$. We also note that $\hat{U}'$ is a generalized LU transformation which only has non-zero action on the subspace with fixed fermion occupation pattern n_2. The above discussions can be generalized into any dimension. Thus we have shown the wave function constructed from the above $(\tilde{\nu}_{d+1}, \tilde{n}_d, \tilde{u}_2^g)$ describes the same fermionic SPT phase.

Finally, we can consider more general case where u_d^g also changes into $\tilde{u}_d^g$. It is very easy to see that such changes in the wave function can be realized by symmetric LU transformation since those u_d^g evolve g_0 are all canceled in the wave function.

In conclusion, we have shown that (ν_{d+1}, n_d, u_d^g) and $(\tilde{\nu}_{d+1}, \tilde{n}_d, \tilde{u}_d^g)$ belonging to the same group supercohomology class will give rise to fixed point wave functions describing the same fermionic SPT phase.

2. Symmetry transformations and generic SPT states

In the Section VI B, we have constructed an ideal fermion SPT state $|\Psi\rangle = \hat{U}|\Phi_0\rangle$ using the the fermionic LU transformation $\hat{U}$ constructed from a cocycle (ν_3, n_2). In this section, we are going to discuss how to construct a more generic SPT state $|\Psi'\rangle$ that is in the same phase as the ideal state $|\Psi\rangle$.

Clearly, $|\Psi'\rangle$ and $|\Psi\rangle$ has the same symmetry. So to construct $|\Psi'\rangle$, we need first discuss how symmetry transformation changes under the fermionic LU transformation $\hat{U}$.

The symmetry G_b that act on $|\Psi\rangle$ is represented by the following (anti) unitary operators

$$\begin{aligned}\hat{W}(g) = &\sum_{\{g_i\}} \prod_{\triangledown} [u_2^g(g_i, g_j, g_k)]^{-1} \prod_{\triangle} u_2^g(g_i, g_j, g_k)\times\\ &\prod_i |g_i\rangle\langle gg_i|\ K^{\frac{1-s(g)}{2}},\end{aligned} \tag{G7}$$

where $\hat{W}_i$ is the symmetry transformation acting on a single site i, and K is the anti unitary operator

$$K\,\mathrm{i}K^{-1} = -\mathrm{i}, \quad KcK^{-1} = c, \quad Kc^{\dagger}K^{-1} = c^{\dagger}. \tag{G8}$$

We find that

$$\hat{W}^0(g) = \hat{U}^\dagger \hat{W}(g) \hat{U} = \sum_{\{g_i\}} \prod_i |g_i\rangle\langle g g_i| \prod_{\bigtriangledown} [u_2^g(g_i,g_j,g_k)]^{-1} \prod_{\bigtriangleup} u_2^g(g_i,g_j,g_k) \times \tag{G9}$$
$$\prod_{\bigtriangledown} c_{(ijk)}^{n_2(g_i,g_j,g_k)} c_{(0ik)}^{n_2(g_0,g_i,g_k)} \bar{c}_{(0jk)}^{n_2(g_0,g_j,g_k)} \bar{c}_{(0ij)}^{n_2(g_0,g_i,g_j)} \prod_{\bigtriangleup} c_{(0ij)}^{n_2(g_0,g_i,g_j)} c_{(0jk)}^{n_2(g_0,g_j,g_k)} \bar{c}_{(0ik)}^{n_2(g_0,g_i,g_k)} \bar{c}_{(ijk)}^{n_2(g_i,g_j,g_k)} \times$$
$$\prod_{\bigtriangleup} \nu_3(g_0,g_i,g_j,g_k) \prod_{\bigtriangledown} \nu_3^{-1}(g_0,g_i,g_j,g_k) \prod_{(0ij)} c_{(0ij)}^{\dagger n_2(g_0,g_i,g_j)} \bar{c}_{(0ij)}^{\dagger n_2(g_0,g_i,g_j)} \times$$
$$\prod_{(0ij)} c_{(0ij)}^{n_2(g_0,gg_i,gg_j)} \bar{c}_{(0ij)}^{n_2(g_0,gg_i,gg_j)} \prod_{\bigtriangleup} [\nu_3^{-1}(g_0,gg_i,gg_j,gg_k)]^{s(g)} \prod_{\bigtriangledown} [\nu_3(g_0,gg_i,gg_j,gg_k)]^{s(g)} K^{\frac{1-s(g)}{2}} \times$$
$$\prod_{\bigtriangleup} c_{(0ij)}^{\dagger n_2(g_0,gg_i,gg_j)} c_{(0jk)}^{\dagger n_2(g_0,gg_j,gg_k)} \bar{c}_{(0ik)}^{\dagger n_2(g_0,gg_i,gg_k)} \bar{c}_{(ijk)}^{\dagger n_2(gg_i,gg_j,gg_k)} \prod_{\bigtriangledown} c_{(ijk)}^{\dagger n_2(gg_i,gg_j,gg_k)} c_{(0ik)}^{\dagger n_2(g_0,gg_i,gg_k)} \bar{c}_{(0jk)}^{\dagger n_2(g_0,gg_j,gg_k)} \bar{c}_{(0ij)}^{\dagger n_2(g_0,gg_i,gg_j)}$$

Using the fact that $\hat{U}$ is independent of g_0, we can change the g_0 in the second half of the right-hand-side of the above expression to gg_0. Then we can use the symmetry condition (46) on (ν_3, n_2, u_2^g) to show that the above is reduced to

$$\hat{W}^0(g) = \hat{U}^\dagger \hat{W}(g) \hat{U} = \sum_{\{g_i\}} \prod_i |g_i\rangle\langle g g_i| K^{\frac{1-s(g)}{2}}. \tag{G10}$$

Note that $\hat{W}^0(g)$ acts on $|\Phi_0\rangle$. If we choose a new $|\Psi_0'\rangle$ that is symmetric under the symmetry G_b: $\hat{W}^0(g)|\Psi_0'\rangle = |\Psi_0'\rangle$, then the resulting $|\Psi'\rangle = \hat{U}|\Psi_0'\rangle$ will be symmetric under $\hat{W}(g)$: $\hat{W}(g)|\Psi'\rangle = |\Psi'\rangle$. Since $\hat{W}^0(g) = \prod_i |g_i\rangle\langle gg_i| K^{\frac{1-s(g)}{2}}$ has a simple form, it is easy to construct the deformed $|\Phi_0\rangle$ that has the same symmetry under $\hat{W}^0(g) = \prod_i |g_i\rangle\langle gg_i|$. Then after the fermion LU transformation $\hat{U}$, we can obtain generic SPT states that are in the same phase as $|\Psi\rangle$.

Here we would like to stress that, only on a system without boundary, the total symmetry transformation $\hat{W}$ is mapped into a simple on-site symmetry $\hat{W}^0$ the LU transformation $\hat{U}$. For system with boundary, under the LU transformation $\hat{U}$, the total symmetry transformation $\hat{W}$ will be mapped into a complicated symmetry transformation which do not have an on-site form on the boundary.[49]

3. Entanglement density matrix

The non-trivialness of the SPT states is in their symmetry-protected gapless boundary excitations. To study the gapless boundary excitations, in this section, we are going to study entanglement density matrix ρ_E and its entanglement Hamiltonian H_E: $\rho_E = \mathrm{e}^{-H_E}$, from the above constructed ground state wave functions for SPT states. The entanglement Hamiltonian H_E can be view as the effective Hamiltonian for the gapless boundary excitations.

To calculate the entanglement density matrix, we first cut the system in to two halves along a horizontal line. To do cutting, we first split each site on the cutting line into two sites: $|g_i\rangle \to |g_i\rangle \otimes |g_{i'}\rangle$ (see Fig. G 4a). The ground state lives in the subspace where $g_i = g_{i'}$. We then cut between the splitted sites.

Because the entanglement spectrum do not change if we perform LU transformations within each half of the system, we can use the LU transformations to deform the lattice into a simpler one in Fig. G 4b, where the sites are labeled by 1, 2, .., L, $1'$, $2'$, .., L', along the edge which forms a ring, and the only two interior sites are labeled by a and a'.

On the deformed lattice, the ground state wave function can be written as $\hat{U}|\Phi_0\rangle$, where

$$|\Phi_0\rangle = |\phi_a\rangle\langle\phi_a| \otimes |\phi_{a'}\rangle\langle\phi_{a'}| \otimes_{i=1}^{L} |\phi_i\rangle\langle\phi_i| \otimes_{i'=1'}^{L'} |\phi_{i'}\rangle\langle\phi_{i'}| \tag{G11}$$



FIG. 17: (Color online) (a) We divide the system into two half alone a line by doubling the sites on the line $|g_i\rangle \to |g_i\rangle \otimes |g_i\rangle$ and then cut between the pairs. (b) The interior of each region is deformed into a simpler lattice without affecting the spectrum of the entanglement density matrix.

and

$$
\begin{aligned}
\hat{U} = {} & \prod_{\bigtriangleup} \nu_3^{-1}(g_0, g_{i'}, g_{j'}, g_{a'}) \prod_{\bigtriangledown} \nu_3(g_0, g_a, g_i, g_j) \times \quad \text{(G12)} \\
& \prod_{ij} \bar{c}_{(0ij)}^{n_2(g_0,g_i,g_j)} c_{(0i'j')}^{n_2(g_0,g_{i'},g_{j'})} \prod_{ai} \bar{c}_{(0ai)}^{n_2(g_0,g_a,g_i)} c_{(0ai)}^{n_2(g_0,g_a,g_i)} \times \\
& \prod_{i'a'} \bar{c}_{(0i'a')}^{n_2(g_0,g_{i'},g_{a'})} c_{(0i'a')}^{n_2(g_0,g_{i'},g_{a'})} \times \\
& \prod_{\bigtriangledown} c_{(aij)}^{\dagger n_2(g_a,g_i,g_j)} c_{(0aj)}^{\dagger n_2(g_0,g_a,g_j)} \bar{c}_{(0ij)}^{\dagger n_2(g_0,g_i,g_j)} \bar{c}_{(0ai)}^{\dagger n_2(g_0,g_a,g_i)} \times \\
& \prod_{\bigtriangleup} c_{(0i'j')}^{\dagger n_2(g_0,g_{i'},g_{j'})} c_{(0j'a')}^{\dagger n_2(g_0,g_{j'},g_{a'})} \bar{c}_{(0i'a')}^{\dagger n_2(g_0,g_{i'},g_{a'})} \bar{c}_{(i'j'a')}^{\dagger n_2(g_{i'},g_{j'},g_{a'})} .
\end{aligned}
$$

Now, let us rewrite $\hat{U}$ as $\hat{U} = \hat{U}_{\text{up}} \hat{U}_{\text{down}}$, where $\hat{U}_{\text{up}}$ acts on the upper half and $\hat{U}_{\text{down}}$ acts on the lower half of the system:

$$
\begin{aligned}
\hat{U} = {} & (-)^{f(g_{1'},\dots,g_{L'})} \times \quad \text{(G13)} \\
& \prod_{\bigtriangleup} \nu_3^{-1}(g_0, g_{i'}, g_{j'}, g_{a'}) \prod_{i'} c_{(0,i'+1',i')}^{n_2(g_0,g_{i'+1'},g_{i'})} \times \\
& \prod_{i'a'} \bar{c}_{(0i'a')}^{n_2(g_0,g_{i'},g_{a'})} c_{(0i'a')}^{n_2(g_0,g_{i'},g_{a'})} \times \\
& \prod_{\bigtriangleup} c_{(0i'j')}^{\dagger n_2(g_0,g_{i'},g_{j'})} c_{(0j'a')}^{\dagger n_2(g_0,g_{j'},g_{a'})} \bar{c}_{(0i'a')}^{\dagger n_2(g_0,g_{i'},g_{a'})} \bar{c}_{(i'j'a')}^{\dagger n_2(g_{i'},g_{j'},g_{a'})} \times \\
& \prod_{\bigtriangledown} \nu_3(g_0, g_a, g_i, g_j) \prod_{i} \bar{c}_{(0,i+1,i)}^{n_2(g_0,g_{i+1},g_i)} \times \\
& \prod_{ai} \bar{c}_{(0ai)}^{n_2(g_0,g_a,g_i)} c_{(0ai)}^{n_2(g_0,g_a,g_i)} \times \\
& \prod_{\bigtriangledown} c_{(aij)}^{\dagger n_2(g_a,g_i,g_j)} c_{(0aj)}^{\dagger n_2(g_0,g_a,g_j)} \bar{c}_{(0ij)}^{\dagger n_2(g_0,g_i,g_j)} \bar{c}_{(0ai)}^{\dagger n_2(g_0,g_a,g_i)} ,
\end{aligned}
$$

where the pure sign factor $(-)^{f(g_{1'},\dots,g_{L'})}$ arises from rewriting $\prod_{ij} \bar{c}_{(0ij)}^{n_2(g_0,g_i,g_j)} c_{(0i'j')}^{n_2(g_0,g_{i'},g_{j'})}$ as $\prod_{i'} c_{(0,i'+1',i')}^{n_2(g_0,g_{i'+1'},g_{i'})} \prod_{i} c_{(0,i+1,i)}^{n_2(g_0,g_{i+1},g_i)}$, and we have used the fact that $g_i = g_{i'}$ in the ground state subspace. Thus the express eqn. (G13) is valid only when $\hat{U}$ acts within the ground state subspace. So we find that

$$
\begin{aligned}
\hat{U}_{\text{up}} = {} & (-)^{f(g_{1'},\dots,g_{L'})} \prod_{\bigtriangleup} \nu_3^{-1}(g_0, g_{i'}, g_{j'}, g_{a'}) \times \quad \text{(G14)} \\
& \prod_{i'} c_{(0,i'+1',i')}^{n_2(g_0,g_{i'+1'},g_{i'})} \prod_{i'a'} \bar{c}_{(0i'a')}^{n_2(g_0,g_{i'},g_{a'})} c_{(0i'a')}^{n_2(g_0,g_{i'},g_{a'})} \times \\
& \prod_{\bigtriangleup} c_{(0i'j')}^{\dagger n_2(g_0,g_{i'},g_{j'})} c_{(0j'a')}^{\dagger n_2(g_0,g_{j'},g_{a'})} \bar{c}_{(0i'a')}^{\dagger n_2(g_0,g_{i'},g_{a'})} \bar{c}_{(i'j'a')}^{\dagger n_2(g_{i'},g_{j'},g_{a'})} ,
\end{aligned}
$$

$$
\begin{aligned}
\hat{U}_{\text{down}} = {} & \prod_{\bigtriangledown} \nu_3(g_0, g_a, g_i, g_j) \prod_{i} \bar{c}_{(0,i+1,i)}^{n_2(g_0,g_{i+1},g_i)} \times \quad \text{(G15)} \\
& \prod_{ai} \bar{c}_{(0ai)}^{n_2(g_0,g_a,g_i)} c_{(0ai)}^{n_2(g_0,g_a,g_i)} \times \\
& \prod_{\bigtriangledown} c_{(aij)}^{\dagger n_2(g_a,g_i,g_j)} c_{(0aj)}^{\dagger n_2(g_0,g_a,g_j)} \bar{c}_{(0ij)}^{\dagger n_2(g_0,g_i,g_j)} \bar{c}_{(0ai)}^{\dagger n_2(g_0,g_a,g_i)} .
\end{aligned}
$$

Now the entanglement density matrix ρ_E can be written as

$$
\begin{aligned}
\rho_E &= \text{Tr}_{\text{upper}} \hat{U} |\Phi_0\rangle\langle\Phi_0| \hat{U}^\dagger \\
&= \text{Tr}_{\text{upper}} \hat{U}_{\text{up}} \hat{U}_{\text{down}} |\Phi_0\rangle\langle\Phi_0| \hat{U}_{\text{down}}^\dagger \hat{U}_{\text{up}}^\dagger \\
&= \text{Tr}_{\text{upper}} \hat{U}_{\text{down}} |\Phi_0\rangle\langle\Phi_0| \hat{U}_{\text{down}}^\dagger , \qquad \text{(G16)}
\end{aligned}
$$

where Tr_{upper} is the trace over the degrees of freedom on the upper half of the system. Since $g_i = g_{i'}$ along the edge in the ground state, ρ_E (as an operator) does not change g_i along the edge (*ie* ρ_E is diagonal in the $|g_i\rangle$ basis). So it is sufficient to discuss ρ_E in the subspace of fixed $|g_i\rangle$'s:

$$
\begin{aligned}
& \rho_E(g_1, \dots, g_L) \\
\propto {} & \sum_{g_a, g'_a} \prod_{\bigtriangledown} \nu_3(g_0, g_a, g_i, g_j) \nu_3^*(g_0, g'_a, g_i, g_j) \times \\
& \prod_{i=1}^{L} \bar{c}_{(0,i+1,i)}^{n_2(g_0,g_{i+1},g_i)} \prod_{ai} \bar{c}_{(0ai)}^{n_2(g_0,g_a,g_i)} c_{(0ai)}^{n_2(g_0,g_a,g_i)} \times \quad \text{(G17)} \\
& \prod_{\bigtriangledown} c_{(aij)}^{\dagger n_2(g_a,g_i,g_j)} c_{(0aj)}^{\dagger n_2(g_0,g_a,g_j)} \bar{c}_{(0ij)}^{\dagger n_2(g_0,g_i,g_j)} \bar{c}_{(0ai)}^{\dagger n_2(g_0,g_a,g_i)} \times \\
& |g_a\rangle\langle g'_a| \prod_{\bigtriangledown} \bar{c}_{(0ai)}^{n_2(g_0,g'_a,g_i)} \bar{c}_{(0ij)}^{n_2(g_0,g_i,g_j)} c_{(0aj)}^{n_2(g_0,g'_a,g_j)} c_{(aij)}^{n_2(g'_a,g_i,g_j)} \times \\
& \prod_{ai} c_{(0ai)}^{\dagger n_2(g_0,g'_a,g_i)} \bar{c}_{(0ai)}^{\dagger n_2(g_0,g'_a,g_i)} \prod_{i=L}^{1} \bar{c}_{(0,i+1,i)}^{\dagger n_2(g_0,g_{i+1},g_i)}
\end{aligned}
$$

We see that ρ_E has a form

$$
\begin{aligned}
\rho_E &= \sum_{\{g_i\}} \rho_E(g_1, \dots, g_L) \otimes (\otimes_i |g_i\rangle\langle g_i|) \\
&= \sum_{\{g_i\}} |\{g_i\}_{\text{edge}}\rangle\langle\{g_i\}_{\text{edge}}| \qquad \text{(G18)}
\end{aligned}
$$

where

$$
\begin{aligned}
|\{g_i\}_{\text{edge}}\rangle = {} & |G_b|^{-1/2} \sum_{g_a} \prod_{\bigtriangledown} \nu_3(g_0, g_a, g_i, g_j) \times \\
& \prod_{i=1}^{L} \bar{c}_{(0,i+1,i)}^{n_2(g_0,g_{i+1},g_i)} \prod_{i} \bar{c}_{(0ai)}^{n_2(g_0,g_a,g_i)} c_{(0ai)}^{n_2(g_0,g_a,g_i)} \times \\
& \prod_{i} c_{(a,i+1,i)}^{\dagger n_2(g_a,g_{i+1},g_i)} c_{(0,a,i)}^{\dagger n_2(g_0,g_a,g_i)} \bar{c}_{(0,i+1,i)}^{\dagger n_2(g_0,g_{i+1},g_i)} \bar{c}_{(0,a,i+1)}^{\dagger n_2(g_0,g_a,g_{i+1})} \times \\
& |g_a\rangle \otimes (\otimes_i |g_i\rangle) \qquad \text{(G19)}
\end{aligned}
$$

We can simplify the above expression (G19). First we note that

$$
\begin{aligned}
&\prod_{i=1}^{L} \bar{c}_{(0,i+1,i)}^{n_2(g_0,g_{i+1},g_i)} \prod_{i} \bar{c}_{(0ai)}^{n_2(g_0,g_a,g_i)} c_{(0ai)}^{n_2(g_0,g_a,g_i)} \\
&= \prod_{i=1}^{L} \bar{c}_{(0ai)}^{n_2(g_0,g_a,g_i)} c_{(0ai)}^{n_2(g_0,g_a,g_i)} \bar{c}_{(0,i+1,i)}^{n_2(g_0,g_{i+1},g_i)} \\
&= (-)^{n_2(g_0,g_a,g_1)+n_2(g_0,g_a,g_1)\sum_{i=1}^{L} n_2(g_0,g_{i+1},g_i)} \times \\
&\quad \prod_{i=1}^{L} c_{(0ai)}^{n_2(g_0,g_a,g_i)} \bar{c}_{(0,i+1,i)}^{n_2(g_0,g_{i+1},g_i)} \bar{c}_{(0,a,i+1)}^{n_2(g_0,g_a,g_{i+1})}
\end{aligned} \tag{G20}
$$

Therefore,

$$
\begin{aligned}
&\prod_{i=1}^{L} \bar{c}_{(0,i+1,i)}^{n_2(g_0,g_{i+1},g_i)} \prod_{i} \bar{c}_{(0ai)}^{n_2(g_0,g_a,g_i)} c_{(0ai)}^{n_2(g_0,g_a,g_i)} \times \\
&\prod_{i} c_{(a,i+1,i)}^{\dagger n_2(g_a,g_{i+1},g_i)} c_{(0,a,i)}^{\dagger n_2(g_0,g_a,g_i)} \bar{c}_{(0,i+1,i)}^{\dagger n_2(g_0,g_{i+1},g_i)} \bar{c}_{(0,a,i+1)}^{\dagger n_2(g_0,g_a,g_{i+1})} \\
&= (-)^{n_2(g_0,g_a,g_1)+n_2(g_0,g_a,g_1)\sum_{i=1}^{L} n_2(g_0,g_{i+1},g_i)} \times \\
&\prod_{i=1}^{L} c_{(0ai)}^{n_2(g_0,g_a,g_i)} \bar{c}_{(0,i+1,i)}^{n_2(g_0,g_{i+1},g_i)} \bar{c}_{(0,a,i+1)}^{n_2(g_0,g_a,g_{i+1})} \times \\
&c_{(a,i+1,i)}^{\dagger n_2(g_a,g_{i+1},g_i)} c_{(0,a,i)}^{\dagger n_2(g_0,g_a,g_i)} \bar{c}_{(0,i+1,i)}^{\dagger n_2(g_0,g_{i+1},g_i)} \bar{c}_{(0,a,i+1)}^{\dagger n_2(g_0,g_a,g_{i+1})} \\
&= (-)^{n_2(g_0,g_a,g_1)+n_2(g_0,g_a,g_1)\sum_{i=1}^{L} n_2(g_0,g_{i+1},g_i)} \times \\
&(-)^{\sum_{i=1}^{L} n_2(g_0,g_a,g_i)n_2(g_0,g_a,g_{i+1})+n_2(g_0,g_{i+1},g_i)n_2(g_a,g_{i+1},g_i)} \times \\
&\quad \prod_{i=1}^{L} c_{(a,i+1,i)}^{\dagger n_2(g_a,g_{i+1},g_i)} .
\end{aligned} \tag{G21}
$$

We find

$$
\begin{aligned}
&|\{g_i\}_{\text{edge}}\rangle = |G_b|^{-1/2} \sum_{g_a} \prod_{i} \nu_3(g_0,g_a,g_{i+1},g_i) \times \\
&\quad (-)^{n_2(g_0,g_a,g_1)+n_2(g_0,g_a,g_1)\sum_{i=1}^{L} n_2(g_0,g_{i+1},g_i)} \times \\
&(-)^{\sum_{i=1}^{L} n_2(g_0,g_a,g_i)n_2(g_0,g_a,g_{i+1})+n_2(g_0,g_{i+1},g_i)n_2(g_a,g_{i+1},g_i)} \times \\
&\quad \prod_{i=1}^{L} c_{(a,i+1,i)}^{\dagger n_2(g_a,g_{i+1},g_i)} |g_a\rangle \otimes (\otimes_i |g_i\rangle)
\end{aligned} \tag{G22}
$$

From the above calculation, and using the relation between the entanglement density matrix and gapless edge excitations, we learn two things. 1) The low energy edge degrees of freedom are label by $\{g_1, ..., g_L\}$ on the boundary, since $\rho_E(g_1, ..., g_L)$ has one and only one non-zero eigenvalue for each $\{g_1, ..., g_L\}$. 2) The low energy edge degrees of freedom are entangled with the bulk degrees of freedom, since the states on the site-a (in the bulk) are different for different $\{g_1, ..., g_L\}$. Using the expression (G22), we calculation the low energy effective Hamiltonian H_{eff} on the edge from a physical Hamiltonian H_{edge} on the edge:

$$
(H_{\text{eff}})_{\{g_i'\},\{g_i\}} = \langle \{g_i'\}_{\text{edge}} | H_{\text{edge}} | \{g_i\}_{\text{edge}} \rangle . \tag{G23}
$$

4. Low energy effective edge Hamiltonian

Now let us assume that H_{edge} only changes g_k and only contains fermion operators $c_{(a,k+1,k)}$ and $c_{(a,k,k-1)}$. We find

$$
\begin{aligned}
(H_{\text{eff}})_{\{g_i'\},\{g_i\}} &= |G_b|^{-1} \sum_{g_a} \Big(\prod_{i\neq k} \delta_{g_i',g_i} \Big) \nu_3^*(g_0,g_a,g_{k+1},g_k') \nu_3^*(g_0,g_a,g_k',g_{k-1}) \nu_3(g_0,g_a,g_{k+1},g_k) \nu_3(g_0,g_a,g_k,g_{k-1}) \times \\
&(-)^{n_2(g_0,g_a,g_1)[n_2(g_0,g_{k+1},g_k)+n_2(g_0,g_k,g_{k-1})+n_2(g_0,g_{k+1},g_k')+n_2(g_0,g_k',g_{k-1})]} \times \\
&(-)^{n_2(g_0,g_a,g_k)n_2(g_0,g_a,g_{k+1})+n_2(g_0,g_{k+1},g_k)n_2(g_a,g_{k+1},g_k)} (-)^{n_2(g_0,g_a,g_k')n_2(g_0,g_a,g_{k+1})+n_2(g_0,g_{k+1},g_k')n_2(g_a,g_{k+1},g_k')} \times \\
&(-)^{n_2(g_0,g_a,g_{k-1})n_2(g_0,g_a,g_k)+n_2(g_0,g_k,g_{k-1})n_2(g_a,g_k,g_{k-1})} (-)^{n_2(g_0,g_a,g_{k-1})n_2(g_0,g_a,g_k')+n_2(g_0,g_k',g_{k-1})n_2(g_a,g_k',g_{k-1})} \times \\
&\delta_{n_2(g_a,g_k,g_{k-1})+n_2(g_a,g_{k+1},g_k)+n_2(g_a,g_k',g_{k-1})+n_2(g_a,g_{k+1},g_k')=\text{even}} \times \\
&\langle g_k' | c_{(a,k+1,k)}^{n_2(g_a,g_{k+1},g_k')} c_{(a,k,k-1)}^{n_2(g_a,g_k',g_{k-1})} H_{\text{edge}} c_{(a,k,k-1)}^{\dagger n_2(g_a,g_k,g_{k-1})} c_{(a,k+1,k)}^{\dagger n_2(g_a,g_{k+1},g_k)} | g_k \rangle .
\end{aligned} \tag{G24}
$$

Since

$$
\begin{aligned}
& n_2(g_a,g_k,g_{k-1}) + n_2(g_a,g_{k+1},g_k) + n_2(g_a,g_k',g_{k-1}) + n_2(g_a,g_{k+1},g_k') \\
&= n_2(g_0,g_k,g_{k-1}) + n_2(g_0,g_{k+1},g_k) + n_2(g_0,g_k',g_{k-1}) + n_2(g_0,g_{k+1},g_k') \bmod 2,
\end{aligned} \tag{G25}
$$

the factor $(-)^{n_2(g_0,g_a,g_1)[n_2(g_0,g_{k+1},g_k)+n_2(g_0,g_k,g_{k-1})+n_2(g_0,g_{k+1},g'_k)+n_2(g_0,g'_k,g_{k-1})]}$ can be dropped:

$$\begin{aligned}(H_{\text{eff}})_{\{g'_i\},\{g_i\}} = |G_b|^{-1}\sum_{g_a}\Big(\prod_{i\neq k}\delta_{g'_i,g_i}\Big)\nu_3^*(g_0,g_a,g_{k+1},g'_k)\nu_3^*(g_0,g_a,g'_k,g_{k-1})\nu_3(g_0,g_a,g_{k+1},g_k)\nu_3(g_0,g_a,g_k,g_{k-1})\times \\ (-)^{n_2(g_0,g_a,g_k)n_2(g_0,g_a,g_{k+1})+n_2(g_0,g_{k+1},g_k)n_2(g_a,g_{k+1},g_k)}(-)^{n_2(g_0,g_a,g'_k)n_2(g_0,g_a,g_{k+1})+n_2(g_0,g_{k+1},g'_k)n_2(g_a,g_{k+1},g'_k)}\times \\ (-)^{n_2(g_0,g_a,g_{k-1})n_2(g_0,g_a,g_k)+n_2(g_0,g_k,g_{k-1})n_2(g_a,g_k,g_{k-1})}(-)^{n_2(g_0,g_a,g_{k-1})n_2(g_0,g_a,g'_k)+n_2(g_0,g'_k,g_{k-1})n_2(g_a,g'_k,g_{k-1})}\times \\ \delta_{n_2(g_a,g_k,g_{k-1})+n_2(g_a,g_{k+1},g_k)+n_2(g_a,g'_k,g_{k-1})+n_2(g_a,g_{k+1},g'_k)=\text{even}}\times \\ \langle g'_k|c_{(a,k+1,k)}^{n_2(g_a,g_{k+1},g'_k)}c_{(a,k,k-1)}^{n_2(g_a,g'_k,g_{k-1})}H_{\text{edge}}c_{(a,k,k-1)}^{\dagger n_2(g_a,g_k,g_{k-1})}c_{(a,k+1,k)}^{\dagger n_2(g_a,g_{k+1},g_k)}|g_k\rangle.\end{aligned} \tag{G26}$$

Now let us consider that following three form of H_{edge}

$$H_{\text{edge}}^{00}=H^{00},\quad H_{\text{edge}}^{++}=H^{++}c^\dagger_{(a,k+,k)}c^\dagger_{(a,k,k-1)},\quad H_{\text{edge}}^{+-}=H^{+-}c^\dagger_{(a,k+1,k)}c_{(a,k,k-1)}, \tag{G27}$$

where H^{00}, H^{++}, and H^{+-} are bosonic operators that only act on $|g_k\rangle$. The corresponding H_{eff} are given by

$$\begin{aligned}(H^{00}_{\text{eff}})_{\{g'_i\},\{g_i\}} = |G_b|^{-1}\sum_{g_a}\Big(\prod_{i\neq k}\delta_{g'_i,g_i}\Big)\nu_3^*(g_0,g_a,g_{k+1},g'_k)\nu_3^*(g_0,g_a,g'_k,g_{k-1})\nu_3(g_0,g_a,g_{k+1},g_k)\nu_3(g_0,g_a,g_k,g_{k-1})\times \\ (-)^{n_2(g_0,g_a,g_k)n_2(g_0,g_a,g_{k+1})+n_2(g_0,g_{k+1},g_k)n_2(g_a,g_{k+1},g_k)}(-)^{n_2(g_0,g_a,g'_k)n_2(g_0,g_a,g_{k+1})+n_2(g_0,g_{k+1},g'_k)n_2(g_a,g_{k+1},g'_k)}\times \\ (-)^{n_2(g_0,g_a,g_{k-1})n_2(g_0,g_a,g_k)+n_2(g_0,g_k,g_{k-1})n_2(g_a,g_k,g_{k-1})}(-)^{n_2(g_0,g_a,g_{k-1})n_2(g_0,g_a,g'_k)+n_2(g_0,g'_k,g_{k-1})n_2(g_a,g'_k,g_{k-1})}\times \\ \delta_{n_2(g_a,g_k,g_{k-1}),n_2(g_a,g'_k,g_{k-1})}\delta_{n_2(g_a,g_{k+1},g_k),n_2(g_a,g_{k+1},g'_k)}H^{00}_{g'_k,g_k};\end{aligned} \tag{G28}$$

$$\begin{aligned}(H^{++}_{\text{eff}})_{\{g'_i\},\{g_i\}} = -|G_b|^{-1}\sum_{g_a}\Big(\prod_{i\neq k}\delta_{g'_i,g_i}\Big)\nu_3^*(g_0,g_a,g_{k+1},g'_k)\nu_3^*(g_0,g_a,g'_k,g_{k-1})\nu_3(g_0,g_a,g_{k+1},g_k)\nu_3(g_0,g_a,g_k,g_{k-1})\times \\ (-)^{n_2(g_0,g_a,g_k)n_2(g_0,g_a,g_{k+1})}(-)^{n_2(g_0,g_a,g'_k)n_2(g_0,g_a,g_{k+1})+n_2(g_0,g_{k+1},g'_k)}\times \\ (-)^{n_2(g_0,g_a,g_{k-1})n_2(g_0,g_a,g_k)}(-)^{n_2(g_0,g_a,g_{k-1})n_2(g_0,g_a,g'_k)+n_2(g_0,g'_k,g_{k-1})}\times \\ \delta_{n_2(g_a,g_k,g_{k-1}),0}\delta_{n_2(g_a,g_{k+1},g_k),0}\delta_{n_2(g_a,g'_k,g_{k-1}),1}\delta_{n_2(g_a,g_{k+1},g'_k),1}H^{++}_{g'_k,g_k};\end{aligned} \tag{G29}$$

$$\begin{aligned}(H^{+-}_{\text{eff}})_{\{g'_i\},\{g_i\}} = |G_b|^{-1}\sum_{g_a}\Big(\prod_{i\neq k}\delta_{g'_i,g_i}\Big)\nu_3^*(g_0,g_a,g_{k+1},g'_k)\nu_3^*(g_0,g_a,g'_k,g_{k-1})\nu_3(g_0,g_a,g_{k+1},g_k)\nu_3(g_0,g_a,g_k,g_{k-1})\times \\ (-)^{n_2(g_0,g_a,g_k)n_2(g_0,g_a,g_{k+1})}(-)^{n_2(g_0,g_a,g'_k)n_2(g_0,g_a,g_{k+1})+n_2(g_0,g_{k+1},g'_k)}\times \\ (-)^{n_2(g_0,g_a,g_{k-1})n_2(g_0,g_a,g_k)+n_2(g_0,g_k,g_{k-1})}(-)^{n_2(g_0,g_a,g_{k-1})n_2(g_0,g_a,g'_k)}\times \\ \delta_{n_2(g_a,g_k,g_{k-1}),1}\delta_{n_2(g_a,g_{k+1},g_k),0}\delta_{n_2(g_a,g'_k,g_{k-1}),0}\delta_{n_2(g_a,g_{k+1},g'_k),1}H^{+-}_{g'_k,g_k}.\end{aligned} \tag{G30}$$

Those expressions allow us to constructed simple edge effective Hamiltonians for the corresponding SPT phases.

To obtain more generic edge effective theory, let us assume that H_{edge} changes both g_k and g_{k+2} and contains fermion operators $c_{(a,k+1,k)}$, $c_{(a,k,k-1)}$, $c_{(a,k+3,k+2)}$, and $c_{(a,k+2,k+1)}$:

$$\begin{aligned}(H_{\text{eff}})_{\{g'_i\},\{g_i\}} = |G_b|^{-1}\sum_{g_a}\Big(\prod_{i\neq k}\delta_{g'_i,g_i}\Big)\nu_3^*(g_0,g_a,g_{k+1},g'_k)\nu_3^*(g_0,g_a,g'_k,g_{k-1})\nu_3(g_0,g_a,g_{k+1},g_k)\nu_3(g_0,g_a,g_k,g_{k-1})\times \\ \nu_3^*(g_0,g_a,g_{k+3},g'_{k+2})\nu_3^*(g_0,g_a,g'_{k+2},g_{k+1})\nu_3(g_0,g_a,g_{k+3},g_{k+2})\nu_3(g_0,g_a,g_{k+2},g_{k+1})\times \\ (-)^{n_2(g_0,g_a,g_k)n_2(g_0,g_a,g_{k+1})+n_2(g_0,g_{k+1},g_k)n_2(g_a,g_{k+1},g_k)}(-)^{n_2(g_0,g_a,g'_k)n_2(g_0,g_a,g_{k+1})+n_2(g_0,g_{k+1},g'_k)n_2(g_a,g_{k+1},g'_k)}\times \\ (-)^{n_2(g_0,g_a,g_{k-1})n_2(g_0,g_a,g_k)+n_2(g_0,g_k,g_{k-1})n_2(g_a,g_k,g_{k-1})}(-)^{n_2(g_0,g_a,g_{k-1})n_2(g_0,g_a,g'_k)+n_2(g_0,g'_k,g_{k-1})n_2(g_a,g'_k,g_{k-1})}\times \\ (-)^{n_2(g_0,g_a,g_{k+2})n_2(g_0,g_a,g_{k+3})+n_2(g_0,g_{k+3},g_{k+2})n_2(g_a,g_{k+3},g_{k+2})}(-)^{n_2(g_0,g_a,g'_{k+2})n_2(g_0,g_a,g_{k+3})+n_2(g_0,g_{k+3},g'_{k+2})n_2(g_a,g_{k+3},g'_{k+2})}\times \\ (-)^{n_2(g_0,g_a,g_{k-1})n_2(g_0,g_a,g_k)+n_2(g_0,g_k,g_{k-1})n_2(g_a,g_k,g_{k-1})}(-)^{n_2(g_0,g_a,g_{k-1})n_2(g_0,g_a,g'_k)+n_2(g_0,g'_k,g_{k-1})n_2(g_a,g'_k,g_{k-1})}\times \\ \delta_{n_2(g_a,g_k,g_{k-1})+n_2(g_a,g_{k+1},g_k)+n_2(g_a,g'_k,g_{k-1})+n_2(g_a,g_{k+1},g'_k)+n_2(g_a,g_{k+2},g_{k+1})+n_2(g_a,g_{k+3},g_{k+2})+n_2(g_a,g'_{k+2},g_{k+1})+n_2(g_a,g_{k+3},g'_{k+2})=\text{even}}\times \\ \langle g'_k|c_{(a,k+3,k+2)}^{n_2(g_a,g_{k+3},g'_{k+2})}c_{(a,k+2,k+1)}^{n_2(g_a,g'_{k+2},g_{k+1})}c_{(a,k+1,k)}^{n_2(g_a,g_{k+1},g'_k)}c_{(a,k,k-1)}^{n_2(g_a,g'_k,g_{k-1})}H_{\text{edge}}\times \\ c_{(a,k,k-1)}^{\dagger n_2(g_a,g_k,g_{k-1})}c_{(a,k+1,k)}^{\dagger n_2(g_a,g_{k+1},g_k)}c_{(a,k+2,k+1)}^{\dagger n_2(g_a,g_{k+2},g_{k+1})}c_{(a,k+3,k+2)}^{\dagger n_2(g_a,g_{k+3},g_{k+2})}|g_k\rangle.\end{aligned} \tag{G31}$$

Appendix H: Ideal Hamiltonian

After constructed the fermionic SPT state $|\Psi\rangle$ in the Section VI A, here we would like construct a local Hamiltonian

$$H=-\sum_i H_i \tag{H1}$$



FIG. 18: (a) The matrix elements of the Hamiltonian term H_i can be obtained from the fermionic path integral of the fixed-point action amplitude $\mathcal{V}_3$ on the complex $\langle g_1g_2g_3g_4g_5g_6g_ig_i'\rangle$. The branching structure on the complex is chosen to march that of the trianglar lattice in Fig. 10. (b) The action of H_i can be obtained by attatch the complex in (a) to the trianglar lattice in Fig. 10. The Grassmann numbers on the six overlapping triangles are integerated out.

that has the G_f symmetry, such that $|\Psi\rangle$ is the unique ground state of the Hamiltonian.

We note that the path integral of the fixed-point action amplitude not only gives rise to an ideal ground state wave function (as discussed above), it also gives rise to an ideal Hamiltonian. From the structure of the fixed-point path integral, we find that H_i has structure that it acts on a seven-spin cluster labeled by i, $1-6$ in blue in Fig. 10, and on the six-fermion cluster on the six triangles inside the hexagon $\langle 123456\rangle$ in Fig. 10.

The matrix elements of H_i can be obtained by evaluating the fixed-point action amplitude $\mathcal{V}_3^\pm$ on the complex in Fig. 18a, since the action of H_i can be realized by attaching the complex Fig. 18a to the triangular lattice and then performing the fermionic path integral on the new complex (see Fig. 18b). The evaluation of $\mathcal{V}_3^\pm$ on the complex give us

$$
\begin{aligned}
&(-)^{m_1(g_4,g_i')+m_1(g_i',g_1)}(-)^{m_1(g_i,g_2)+m_1(g_3,g_i)+m_1(g_5,g_i)+m_1(g_i,g_6)}\int\prod_\Sigma\mathcal{V}_3\\
=&(-)^{m_1(g_4,g_i')+m_1(g_i',g_1)+m_1(g_i,g_2)+m_1(g_3,g_i)+m_1(g_5,g_i)+m_1(g_i,g_6)}\int\prod_{j=1,2,6}\mathrm{d}\theta_{(ii'j)}^{n_2(g_i,g_i',g_j)}\mathrm{d}\bar\theta_{(ii'j)}^{n_2(g_i,g_i',g_j)}\prod_{j=3,4,5}\mathrm{d}\theta_{(jii')}^{n_2(g_j,g_i,g_i')}\mathrm{d}\bar\theta_{(jii')}^{n_2(g_j,g_i,g_i')}\\
&(-)^{m_1(g_4,g_i)+m_1(g_5,g_i')+m_1(g_i,g_6)+m_1(g_i',g_1)+m_1(g_i,g_2)+m_1(g_3,g_i')}(-)^{m_1(g_i,g_i')}\times\\
&\mathcal{V}_3^+(g_5,g_i,g_i',g_6)\mathcal{V}_3^+(g_4,g_5,g_i,g_i')\mathcal{V}_3^-(g_4,g_3,g_i,g_i')\mathcal{V}_3^-(g_3,g_i,g_i',g_2)\mathcal{V}_3^-(g_i,g_i',g_2,g_1)\mathcal{V}_3^+(g_i,g_i',g_6,g_1)\\
=&(-)^{m_1(g_4,g_i')+m_1(g_i',g_1)+m_1(g_i,g_2)+m_1(g_3,g_i)+m_1(g_5,g_i)+m_1(g_i,g_6)}\int\prod_{j=1,2,6}\mathrm{d}\theta_{(ii'j)}^{n_2(g_i,g_i',g_j)}\mathrm{d}\bar\theta_{(ii'j)}^{n_2(g_i,g_i',g_j)}\prod_{j=3,4,5}\mathrm{d}\theta_{(jii')}^{n_2(g_j,g_i,g_i')}\mathrm{d}\bar\theta_{(jii')}^{n_2(g_j,g_i,g_i')}\\
&(-)^{m_1(g_4,g_i)+m_1(g_5,g_i')+m_1(g_i,g_6)+m_1(g_i',g_1)+m_1(g_i,g_2)+m_1(g_3,g_i')}(-)^{m_1(g_i,g_i')}\times\\
&\nu_3(g_5,g_i,g_i',g_6)\theta_{(ii'6)}^{n_2(g_i,g_i',g_6)}\theta_{(5i6)}^{n_2(g_5,g_i,g_6)}\bar\theta_{(5i'6)}^{n_2(g_5,g_i',g_6)}\bar\theta_{(5ii')}^{n_2(g_5,g_i,g_i')}\nu_3(g_4,g_5,g_i,g_i')\theta_{(5ii')}^{n_2(g_5,g_i,g_i')}\theta_{(45i')}^{n_2(g_4,g_5,g_i')}\bar\theta_{(4ii')}^{n_2(g_4,g_i,g_i')}\bar\theta_{(45i)}^{n_2(g_4,g_5,g_i)}\\
&\nu_3^{-1}(g_4,g_3,g_i,g_i')\theta_{(43i)}^{n_2(g_4,g_3,g_i)}\theta_{(4ii')}^{n_2(g_4,g_i,g_i')}\bar\theta_{(43i')}^{n_2(g_4,g_3,g_i')}\bar\theta_{(3ii')}^{n_2(g_3,g_i,g_i')}\nu_3^{-1}(g_3,g_i,g_i',g_2)\theta_{(3ii')}^{n_2(g_3,g_i,g_i')}\theta_{(3i'2)}^{n_2(g_3,g_i',g_2)}\bar\theta_{(3i2)}^{n_2(g_3,g_i,g_2)}\bar\theta_{(ii'2)}^{n_2(g_i,g_i',g_2)}\\
&\nu_3^{-1}(g_i,g_i',g_2,g_1)\theta_{(ii'2)}^{n_2(g_i,g_i',g_2)}\theta_{(i21)}^{n_2(g_i,g_2,g_1)}\bar\theta_{(ii'1)}^{n_2(g_i,g_i',g_1)}\bar\theta_{(i'21)}^{n_2(g_i',g_2,g_1)}\nu_3(g_i,g_i',g_6,g_1)\theta_{(i'61)}^{n_2(g_i',g_6,g_1)}\theta_{(ii'1)}^{n_2(g_i,g_i',g_1)}\bar\theta_{(i61)}^{n_2(g_i,g_6,g_1)}\bar\theta_{(ii'6)}^{n_2(g_i,g_i',g_6)}.
\end{aligned}
\tag{H2}
$$

Just like the wave function, we have include an extra factor $(-)^{m_1(g_4,g_i')+m_1(g_i',g_1)}(-)^{m_1(g_i,g_2)+m_1(g_3,g_i)+m_1(g_5,g_i)+m_1(g_i,g_6)}$ in the path integral. With this extra factor, multiplying eqn. (H2) to the wave function can then be viewed as attaching the complex Fig. 18a to the triangular lattice (see Fig. 18b). Note that the factor $(-)^{m_1(g_i,g_2)+m_1(g_3,g_i)+m_1(g_5,g_i)+m_1(g_i,g_6)}$ plus the factor that we included in the wave function $(-)^{m_1(g_4,g_i)+m_1(g_i,g_1)}$ give us a factor $(-)^{m_1(g_4,g_i)+m_1(g_i,g_1)+m_1(g_i,g_2)+m_1(g_3,g_i)+m_1(g_5,g_i)+m_1(g_i,g_6)}$ on all the six interior edges $\{i,1\}$, $\{i,2\}$,..., $\{i,6\}$. This factor is needed in the path integral on the complex in Fig. 18b obtained by attaching the complex in Fig. 18a to the triangular lattice. Also note that the factor $(-)^{m_1(g_4,g_i')+m_1(g_i',g_1)}$ is what we need in the new wave function after the action of attaching the complex Fig. 18a to the triangular lattice (see Fig. 18b). By combining all the m_1 factors on the right-hand-side of eqn. (H2), we find that all those m_1 factors become

$(-)^{n_2(g_3,g_i,g_i')+n_2(g_4,g_i,g_i')+n_2(g_5,g_i,g_i')}$. So we find

$$
\begin{aligned}
&(-)^{m_1(g_4,g_i')+m_1(g_i',g_1)}(-)^{m_1(g_i,g_2)+m_1(g_3,g_i)+m_1(g_5,g_i)+m_1(g_i,g_6)}\int\prod_{\Sigma}\mathcal{V}_3 \\
&= (-)^{n_2(g_3,g_i,g_i')+n_2(g_4,g_i,g_i')+n_2(g_5,g_i,g_i')}\int\prod_{j=1,2,6}\mathrm{d}\theta_{(ii'j)}^{n_2(g_i,g_i',g_j)}\mathrm{d}\bar\theta_{(ii'j)}^{n_2(g_i,g_i',g_j)}\prod_{j=3,4,5}\mathrm{d}\theta_{(jii')}^{n_2(g_j,g_i,g_i')}\mathrm{d}\bar\theta_{(jii')}^{n_2(g_j,g_i,g_i')}\times \\
&(-)^{n_2(g_4,g_5,g_i)n_2(g_4,g_i,g_i')+n_2(g_4,g_i,g_i')n_2(g_4,g_3,g_i)+n_2(g_i',g_2,g_1)n_2(g_i,g_i',g_1)+n_2(g_i,g_i',g_1)n_2(g_i',g_6,g_1)} \\
&\nu_3(g_5,g_i,g_i',g_6)\theta_{(ii'6)}^{n_2(g_i,g_i',g_6)}\theta_{(5i6)}^{n_2(g_5,g_i,g_6)}\bar\theta_{(5i'6)}^{n_2(g_5,g_i',g_6)}\bar\theta_{(5ii')}^{n_2(g_5,g_i,g_i')}\nu_3(g_4,g_5,g_i,g_i')\theta_{(5ii')}^{n_2(g_5,g_i,g_i')}\theta_{(45i')}^{n_2(g_4,g_5,g_i')}\bar\theta_{(45i)}^{n_2(g_4,g_5,g_i)}\bar\theta_{(4ii')}^{n_2(g_4,g_i,g_i')} \\
&\nu_3^{-1}(g_4,g_3,g_i,g_i')\theta_{(4ii')}^{n_2(g_4,g_i,g_i')}\theta_{(43i)}^{n_2(g_4,g_3,g_i)}\bar\theta_{(43i')}^{n_2(g_4,g_3,g_i')}\bar\theta_{(3ii')}^{n_2(g_3,g_i,g_i')}\nu_3^{-1}(g_3,g_i,g_i',g_2)\theta_{(3ii')}^{n_2(g_3,g_i,g_i')}\theta_{(3i'2)}^{n_2(g_3,g_i',g_2)}\bar\theta_{(3i2)}^{n_2(g_3,g_i,g_2)}\bar\theta_{(ii'2)}^{n_2(g_i,g_i',g_2)} \\
&\nu_3^{-1}(g_i,g_i',g_2,g_1)\theta_{(ii'2)}^{n_2(g_i,g_i',g_2)}\theta_{(i21)}^{n_2(g_i,g_2,g_1)}\bar\theta_{(i'21)}^{n_2(g_i',g_2,g_1)}\bar\theta_{(ii'1)}^{n_2(g_i,g_i',g_1)}\nu_3(g_i,g_i',g_6,g_1)\theta_{(ii'1)}^{n_2(g_i,g_i',g_1)}\theta_{(i'61)}^{n_2(g_i',g_6,g_1)}\bar\theta_{(i61)}^{n_2(g_i,g_6,g_1)}\bar\theta_{(ii'6)}^{n_2(g_i,g_i',g_6)}.
\end{aligned}
\tag{H3}
$$

In the above, we have also rearranged the order of the Grassmann numbers within some simplices to bring, say, $\bar\theta_{(4ii')}^{n_2(g_4,g_i,g_i')}$ next to $\theta_{(4ii')}^{n_2(g_4,g_i,g_i')}$. Now we can integral out $\mathrm{d}\theta\mathrm{d}\bar\theta$'s and obtain

$$
\begin{aligned}
&(-)^{m_1(g_4,g_i')+m_1(g_i',g_1)}(-)^{m_1(g_i,g_2)+m_1(g_3,g_i)+m_1(g_5,g_i)+m_1(g_i,g_6)}\int\prod_{\Sigma}\mathcal{V}_3 = (-)^{n_2(g_i,g_i',g_6)}\times \\
&(-)^{n_2(g_3,g_i,g_i')+n_2(g_4,g_i,g_i')+n_2(g_5,g_i,g_i')+n_2(g_4,g_5,g_i)n_2(g_4,g_i,g_i')+n_2(g_4,g_i,g_i')n_2(g_4,g_3,g_i)+n_2(g_i',g_2,g_1)n_2(g_i,g_i',g_1)+n_2(g_i,g_i',g_1)n_2(g_i',g_6,g_1)} \\
&\nu_3(g_5,g_i,g_i',g_6)\theta_{(5i6)}^{n_2(g_5,g_i,g_6)}\bar\theta_{(5i'6)}^{n_2(g_5,g_i',g_6)}\nu_3(g_4,g_5,g_i,g_i')\theta_{(45i')}^{n_2(g_4,g_5,g_i')}\bar\theta_{(45i)}^{n_2(g_4,g_5,g_i)}\nu_3^{-1}(g_4,g_3,g_i,g_i')\theta_{(43i)}^{n_2(g_4,g_3,g_i)}\bar\theta_{(43i')}^{n_2(g_4,g_3,g_i')} \\
&\nu_3^{-1}(g_3,g_i,g_i',g_2)\theta_{(3i'2)}^{n_2(g_3,g_i',g_2)}\bar\theta_{(3i2)}^{n_2(g_3,g_i,g_2)}\nu_3^{-1}(g_i,g_i',g_2,g_1)\theta_{(i21)}^{n_2(g_i,g_2,g_1)}\bar\theta_{(i'21)}^{n_2(g_i',g_2,g_1)}\nu_3(g_i,g_i',g_6,g_1)\theta_{(i'61)}^{n_2(g_i',g_6,g_1)}\bar\theta_{(i61)}^{n_2(g_i,g_6,g_1)}
\end{aligned}
\tag{H4}
$$

where the factor $(-)^{n_2(g_i,g_i',g_6)}$ comes from bringing $\bar\theta_{(ii'6)}^{n_2(g_i,g_i',g_6)}$ in eqn. (H3) all the way from the back to the front. Let us rearrange the order of the Grassmann numbers to bring, say, $\bar\theta_{(5i'6)}^{n_2(g_5,g_i',g_6)}$ in front of $\theta_{(5i6)}^{n_2(g_5,g_i,g_6)}$:

$$
\begin{aligned}
&(-)^{m_1(g_4,g_i')+m_1(g_i',g_1)}(-)^{m_1(g_i,g_2)+m_1(g_3,g_i)+m_1(g_5,g_i)+m_1(g_i,g_6)}\int\prod_{\Sigma}\mathcal{V}_3 \\
&= (-)^{n_2(g_i,g_i',g_6)+n_2(g_3,g_i,g_i')+n_2(g_4,g_i,g_i')+n_2(g_5,g_i,g_i')}\times \\
&\quad (-)^{n_2(g_4,g_5,g_i)n_2(g_4,g_i,g_i')+n_2(g_4,g_i,g_i')n_2(g_4,g_3,g_i)+n_2(g_i',g_2,g_1)n_2(g_i,g_i',g_1)+n_2(g_i,g_i',g_1)n_2(g_i',g_6,g_1)}\times \\
&\quad \frac{\nu_3(g_4,g_5,g_i,g_i')\nu_3(g_5,g_i,g_i',g_6)\nu_3(g_i,g_i',g_6,g_1)}{\nu_3(g_i,g_i',g_2,g_1)\nu_3(g_3,g_i,g_i',g_2)\nu_3(g_4,g_3,g_i,g_i')}(-)^{n_2(g_5,g_i',g_6)n_2(g_5,g_i,g_6)+n_2(g_4,g_3,g_i')n_2(g_4,g_3,g_i)+n_2(g_i',g_2,g_1)n_2(g_i,g_2,g_1)}\times \\
&\qquad \bar\theta_{(5i'6)}^{n_2(g_5,g_i',g_6)}\theta_{(5i6)}^{n_2(g_5,g_i,g_6)}\theta_{(45i')}^{n_2(g_4,g_5,g_i')}\bar\theta_{(45i)}^{n_2(g_4,g_5,g_i)}\bar\theta_{(43i')}^{n_2(g_4,g_3,g_i')}\theta_{(43i)}^{n_2(g_4,g_3,g_i)}\times \\
&\qquad \theta_{(3i'2)}^{n_2(g_3,g_i',g_2)}\bar\theta_{(3i2)}^{n_2(g_3,g_i,g_2)}\bar\theta_{(i'21)}^{n_2(g_i',g_2,g_1)}\theta_{(i21)}^{n_2(g_i,g_2,g_1)}\theta_{(i'61)}^{n_2(g_i',g_6,g_1)}\bar\theta_{(i61)}^{n_2(g_i,g_6,g_1)}
\end{aligned}
\tag{H5}
$$

Indeed, the above expression can be regarded as the fermion coherent state representation of the ideal Hamiltonian. We can replace, for example, $\bar\theta_{(5i'6)}^{n_2(g_5,g_i',g_6)}\theta_{(5i6)}^{n_2(g_5,g_i,g_6)}$ as:

$$
\left(c_{(5i6)}^\dagger\right)^{n_2(g_5,g_i',g_6)}c_{(5i6)}^{n_2(g_5,g_i,g_6)} - (1-n_2(g_5,g_i',g_6))\,(1-n_2(g_5,g_i,g_6))\,c_{(5i6)}^\dagger c_{(5i6)}, \tag{H6}
$$

We note that the fermion coherent state is defined as $|\theta_{(5i6)}\rangle = |0\rangle - \theta_{(5i6)}c_{(5i6)}^\dagger|0\rangle$. It is easy to check

$$
\langle\bar\theta_{(5i'6)}|\left(c_{(5i6)}^\dagger\right)^{n_2(g_5,g_i',g_6)}c_{(5i6)}^{n_2(g_5,g_i,g_6)} - (1-n_2(g_5,g_i',g_6))\,(1-n_2(g_5,g_i,g_6))\,c_{(5i6)}^\dagger c_{(5i6)}|\theta_{(5i6)}\rangle = \bar\theta_{(5i'6)}^{n_2(g_5,g_i',g_6)}\theta_{(5i6)}^{n_2(g_5,g_i,g_6)} \tag{H7}
$$

However, a sign factor $(-)^{n_2(g_5,g_i,g_6)}$ is required when the Hamiltonian acts on the coherent state since $\mathrm{d}\theta_{(5i6)}^{n_2(g_5,g_i,g_6)}\mathrm{d}\bar\theta_{(5i6)}^{n_2(g_5,g_i,g_6)}$ should be reordered as $\mathrm{d}\bar\theta_{(5i6)}^{n_2(g_5,g_i,g_6)}\mathrm{d}\theta_{(5i6)}^{n_2(g_5,g_i,g_6)}$. Similarly, for the face $(45i)$, the fermion coherent state is defined as $|\bar\theta_{(45i)}\rangle = |0\rangle - \bar\theta_{(45i)}c_{(45i)}^\dagger|0\rangle$. In this case $\mathrm{d}\theta_{(45i)}^{n_2(g_4,g_5,g_i)}\mathrm{d}\bar\theta_{(45i)}^{n_2(g_4,g_5,g_i)}$ is the correct ordering and we don't need to introduce the extra sign factor.

By applying the same discussions to other triangles, we find that $43i$ and $i12$ also contribute sign factors $(-)^{n_2(g_4,g_3,g_i)}$ and $(-)^{n_2(g_i,g_2,g_1)}$. We also note that

$$\begin{aligned} &(-)^{n_2(g_i,g_i',g_6)+n_2(g_3,g_i,g_i')+n_2(g_4,g_i,g_i')+n_2(g_5,g_i,g_i')}(-)^{n_2(g_5,g_i,g_6)+n_2(g_4,g_3,g_i)+n_2(g_i,g_2,g_1)} \\ &= (-)^{n_2(g_4,g_3,g_i')+n_2(g_5,g_i',g_6)+n_2(g_i,g_2,g_1)}. \end{aligned} \tag{H8}$$

and finally obtain

$$H_i = \sum_{g_i,g_i'} |g_i', g_1g_2g_3g_4g_5g_6\rangle\langle g_i, g_1g_2g_3g_4g_5g_6| \ \frac{\nu_3(g_4,g_5,g_i,g_i')\nu_3(g_5,g_i,g_i',g_6)\nu_3(g_i,g_i',g_6,g_1)}{\nu_3(g_i,g_i',g_2,g_1)\nu_3(g_3,g_i,g_i',g_2)\nu_3(g_4,g_3,g_i,g_i')} \times \tag{H9}$$

$$\begin{aligned} &(-)^{n_2(g_4,g_5,g_i)n_2(g_4,g_i,g_i')+n_2(g_4,g_i,g_i')n_2(g_4,g_3,g_i)+n_2(g_i',g_2,g_1)n_2(g_i,g_i',g_1)+n_2(g_i,g_i',g_1)n_2(g_i',g_6,g_1)} \times \\ &(-)^{n_2(g_5,g_i',g_6)n_2(g_5,g_i,g_6)+n_2(g_4,g_3,g_i')n_2(g_4,g_3,g_i)+n_2(g_i',g_2,g_1)n_2(g_i,g_2,g_1)}(-)^{n_2(g_4,g_3,g_i')+n_2(g_5,g_i',g_6)+n_2(g_i,g_2,g_1)} \times \\ &\left[\left(c^\dagger_{(5i6)}\right)^{n_2(g_5,g_i',g_6)} c_{(5i6)}^{n_2(g_5,g_i,g_6)} - (1-n_2(g_5,g_i',g_6))\,(1-n_2(g_5,g_i,g_6))\, c^\dagger_{(5i6)}c_{(5i6)}\right] \times \\ &\left[\left(c^\dagger_{(45i)}\right)^{n_2(g_4,g_5,g_i')} c_{(45i)}^{n_2(g_4,g_5,g_i)} - (1-n_2(g_4,g_5,g_i'))\,(1-n_2(g_4,g_5,g_i))\, c^\dagger_{(45i)}c_{(45i)}\right] \times \\ &\left[\left(c^\dagger_{(43i)}\right)^{n_2(g_4,g_3,g_i')} c_{(43i)}^{n_2(g_4,g_3,g_i)} - (1-n_2(g_4,g_3,g_i'))\,(1-n_2(g_4,g_3,g_i))\, c^\dagger_{(43i)}c_{(43i)}\right] \times \\ &\left[\left(c^\dagger_{(3i2)}\right)^{n_2(g_3,g_i',g_2)} c_{(3i2)}^{n_2(g_3,g_i,g_2)} - (1-n_2(g_3,g_i',g_2))\,(1-n_2(g_3,g_i,g_2))\, c^\dagger_{(3i2)}c_{(3i2)}\right] \times \\ &\left[\left(c^\dagger_{(i21)}\right)^{n_2(g_i',g_2,g_1)} c_{(i21)}^{n_2(g_i,g_2,g_1)} - (1-n_2(g_i',g_2,g_1))\,(1-n_2(g_i,g_2,g_1))\, c^\dagger_{(i21)}c_{(i21)}\right] \times \\ &\left[\left(c^\dagger_{(i61)}\right)^{n_2(g_i',g_6,g_1)} c_{(i61)}^{n_2(g_i,g_6,g_1)} - (1-n_2(g_i',g_6,g_1))\,(1-n_2(g_i,g_6,g_1))\, c^\dagger_{(i61)}c_{(i61)}\right] \end{aligned} \tag{H10}$$

The above expression is valid in the subspace where the fermion occupation number n_{ijk} on each triangle (ijk) satisfies $n_{ijk} = n_2(g_i, g_j, g_k)$.

Next, we would like to show that H_i is hermitian. This property is very important and it makes the whole theory to be unitary. In the above, we express the matrix element $H_{i;g_i,g_i'}(g_1g_2g_3g_4g_5g_6)$ as:

$$(-)^{m_1(g_4,g_i')+m_1(g_i',g_1)+m_1(g_i,g_2)+m_1(g_3,g_i)+m_1(g_5,g_i)+m_1(g_i,g_6)} \int \prod_{\Sigma} \mathcal{V}_3, \tag{H11}$$

hence, its hermitian conjugate $H^*_{i;g_i',g_i}(g_1g_2g_3g_4g_5g_6)$ reads:

$$\begin{aligned} &(-)^{m_1(g_4,g_i)+m_1(g_i,g_1)+m_1(g_i',g_2)+m_1(g_3,g_i')+m_1(g_5,g_i')+m_1(g_i',g_6)} \int \prod_{j=1,2,6} \mathrm{d}\theta_{(ii'j)}^{n_2(g_i',g_i,g_j)} \mathrm{d}\bar\theta_{(ii'j)}^{n_2(g_i',g_i,g_j)} \prod_{j=3,4,5} \mathrm{d}\theta_{(jii')}^{n_2(g_j,g_i',g_i)} \mathrm{d}\bar\theta_{(jii')}^{n_2(g_j,g_i',g_i)} \\ &(-)^{m_1(g_4,g_i')+m_1(g_5,g_i)+m_1(g_i',g_6)+m_1(g_i,g_1)+m_1(g_i',g_2)+m_1(g_3,g_i)}(-)^{m_1(g_i',g_i)} \times \\ &\nu_3^{-1}(g_5,g_i',g_i,g_6)\theta_{(5ii')}^{n_2(g_5,g_i',g_i)}\theta_{(5i'6)}^{n_2(g_5,g_i,g_6)}\bar\theta_{(5i6)}^{n_2(g_5,g_i',g_6)}\bar\theta_{(ii'6)}^{n_2(g_i',g_i,g_6)}\nu_3^{-1}(g_4,g_5,g_i',g_i)\theta_{(45i)}^{n_2(g_4,g_5,g_i')}\theta_{(4ii')}^{n_2(g_4,g_i',g_i)}\bar\theta_{(45i')}^{n_2(g_4,g_5,g_i)}\bar\theta_{(5ii')}^{n_2(g_5,g_i',g_i)} \\ &\nu_3(g_4,g_3,g_i',g_i)\theta_{(3ii')}^{n_2(g_3,g_i',g_i)}\theta_{(43i')}^{n_2(g_4,g_3,g_i)}\bar\theta_{(4ii')}^{n_2(g_4,g_i',g_i)}\bar\theta_{(43i)}^{n_2(g_4,g_3,g_i')}\nu_3(g_3,g_i',g_i,g_2)\theta_{(ii'2)}^{n_2(g_i',g_i,g_2)}\theta_{(3i2)}^{n_2(g_3,g_i',g_2)}\bar\theta_{(3i'2)}^{n_2(g_3,g_i,g_2)}\bar\theta_{(3ii')}^{n_2(g_3,g_i',g_i)} \\ &\nu_3(g_i',g_i,g_2,g_1)\bar\theta_{(i'21)}^{n_2(g_i,g_2,g_1)}\bar\theta_{(ii'1)}^{n_2(g_i',g_i,g_1)}\theta_{(i21)}^{n_2(g_i',g_2,g_1)}\theta_{(ii'2)}^{n_2(g_i',g_i,g_2)}\nu_3^{-1}(g_i',g_i,g_6,g_1)\theta_{(ii'6)}^{n_2(g_i',g_i,g_6)}\theta_{(i61)}^{n_2(g_i',g_6,g_1)}\bar\theta_{(ii'1)}^{n_2(g_i',g_i,g_1)}\bar\theta_{(i'61)}^{n_2(g_i,g_6,g_1)}. \end{aligned} \tag{H12}$$

Note $\nu^* = \nu^{-1}$(ν is a $U(1)$ phase factor) and $(\theta_1\theta_2\bar\theta_3\bar\theta_4)^* = \theta_4\theta_3\bar\theta_2\bar\theta_1$. However, we can not directly compare the above expression with $H_{i;g_i,g_i'}(g_1g_2g_3g_4g_5g_6)$, since its a function of new pairs of Grassmann variable $\theta_{(5i'6)}^{n_2(g_5,g_i',g_6)}\bar\theta_{(5i6)}^{n_2(g_5,g_i,g_6)}, \bar\theta_{(45i')}^{n_2(g_4,g_5,g_i')}\theta_{(45i)}^{n_2(g_4,g_5,g_i)}, \theta_{(43i')}^{n_2(g_4,g_3,g_i')}\bar\theta_{(43i)}^{n_2(g_4,g_3,g_i)}\bar\theta_{(3i'2)}^{n_2(g_3,g_i',g_2)}\theta_{(3i2)}^{n_2(g_3,g_i,g_2)}, \theta_{(i'21)}^{n_2(g_i',g_2,g_1)}\bar\theta_{(i21)}^{n_2(g_i,g_2,g_1)}$ and $\bar\theta_{(i'61)}^{n_2(g_i',g_6,g_1)}\theta_{(i61)}^{n_2(g_i,g_6,g_1)}$. Such a difference is simply because the action of H_i maps, for example, $\bar\theta_{(5i6)}$ to $\bar\theta_{(5i'6)}$ while the action of $H_i^\dagger$ maps $\bar\theta_{(5i'6)}$ back to $\bar\theta_{(5i6)}$. Thus, to see whether they are the same mapping in the Hilbert space or not, we need to redefine $\bar{\theta}_{(5i6)}(\theta_{(5i6)})$ as $\bar{\theta}_{(5i'6)}(\theta_{(5i'6)})$ and$\bar{\theta}_{(5i'6)}(\theta_{(5i'6)})$ as $\bar{\theta}_{(5i6)}(\theta_{(5i6)})$. A simple way to do so is just replacing $i(i')$ by $i'(i)$ in the above expression:

$$
\begin{aligned}
&(-)^{m_1(g_4,g_i)+m_1(g_i,g_1)+m_1(g_i',g_2)+m_1(g_3,g_i')+m_1(g_5,g_i')+m_1(g_i',g_6)} \int \prod_{j=1,2,6} \mathrm{d}\theta_{(i'ij)}^{n_2(g_i',g_i,g_j)} \mathrm{d}\bar{\theta}_{(i'ij)}^{n_2(g_i',g_i,g_j)} \prod_{j=3,4,5} \mathrm{d}\theta_{(ji'i)}^{n_2(g_j,g_i',g_i)} \mathrm{d}\bar{\theta}_{(ji'i)}^{n_2(g_j,g_i',g_i)} \\
&(-)^{m_1(g_4,g_i')+m_1(g_5,g_i)+m_1(g_i',g_6)+m_1(g_i,g_1)+m_1(g_i',g_2)+m_1(g_3,g_i)} (-)^{m_1(g_i',g_i)} \times \\
&\nu_3^{-1}(g_5,g_i',g_i,g_6)\theta_{(5i'i)}^{n_2(g_5,g_i',g_i)}\theta_{(5i6)}^{n_2(g_5,g_i,g_6)}\bar{\theta}_{(5i'6)}^{n_2(g_5,g_i',g_6)}\bar{\theta}_{(i'i6)}^{n_2(g_i',g_i,g_6)}\nu_3^{-1}(g_4,g_5,g_i',g_i)\theta_{(45i')}^{n_2(g_4,g_5,g_i')}\theta_{(4i'i)}^{n_2(g_4,g_i',g_i)}\bar{\theta}_{(45i)}^{n_2(g_4,g_5,g_i)}\bar{\theta}_{(5i'i)}^{n_2(g_5,g_i',g_i)} \\
&\nu_3(g_4,g_3,g_i',g_i)\theta_{(3i'i)}^{n_2(g_3,g_i',g_i)}\theta_{(43i)}^{n_2(g_4,g_3,g_i)}\bar{\theta}_{(4i'i)}^{n_2(g_4,g_i',g_i)}\bar{\theta}_{(43i')}^{n_2(g_4,g_3,g_i')}\nu_3(g_3,g_i',g_i,g_2)\theta_{(i'i2)}^{n_2(g_i',g_i,g_2)}\theta_{(3i'2)}^{n_2(g_3,g_i',g_2)}\bar{\theta}_{(3i2)}^{n_2(g_3,g_i,g_2)}\bar{\theta}_{(3i'i)}^{n_2(g_3,g_i',g_i)} \\
&\nu_3(g_i',g_i,g_2,g_1)\bar{\theta}_{(i21)}^{n_2(g_i,g_2,g_1)}\bar{\theta}_{(i'i1)}^{n_2(g_i',g_i,g_1)}\theta_{(i'21)}^{n_2(g_i',g_2,g_1)}\theta_{(i'i2)}^{n_2(g_i',g_i,g_2)}\nu_3^{-1}(g_i',g_i,g_6,g_1)\theta_{(i'i6)}^{n_2(g_i',g_i,g_6)}\theta_{(i'61)}^{n_2(g_i',g_6,g_1)}\bar{\theta}_{(i'i1)}^{n_2(g_i',g_i,g_1)}\bar{\theta}_{(i61)}^{n_2(g_i,g_6,g_1)} \\
&= (-)^{m_1(g_4,g_i)+m_1(g_i,g_1)+m_1(g_i',g_2)+m_1(g_3,g_i')+m_1(g_5,g_i')+m_1(g_i',g_6)} \int \prod_{j=1,2,6} \mathrm{d}\theta_{(i'ij)}^{n_2(g_i',g_i,g_j)} \mathrm{d}\bar{\theta}_{(i'ij)}^{n_2(g_i',g_i,g_j)} \prod_{j=3,4,5} \mathrm{d}\theta_{(ji'i)}^{n_2(g_j,g_i',g_i)} \mathrm{d}\bar{\theta}_{(ji'i)}^{n_2(g_j,g_i',g_i)} \\
&(-)^{m_1(g_4,g_i')+m_1(g_5,g_i)+m_1(g_i',g_6)+m_1(g_i,g_1)+m_1(g_i',g_2)+m_1(g_3,g_i)} (-)^{m_1(g_i',g_i)} \times \\
&\mathcal{V}_3^-(g_5,g_i',g_i,g_6)\mathcal{V}_3^-(g_4,g_5,g_i',g_i)\mathcal{V}_3^+(g_4,g_3,g_i',g_i)\mathcal{V}_3^+(g_3,g_i',g_i,g_2)\mathcal{V}_3^+(g_i',g_i,g_2,g_1)\mathcal{V}_3^-(g_i',g_i,g_6,g_1) \\
&= (-)^{m_1(g_4,g_i)+m_1(g_i,g_1)+m_1(g_i',g_2)+m_1(g_3,g_i')+m_1(g_5,g_i')+m_1(g_i',g_6)} \int \prod_{\Sigma} \mathcal{V}_3. \qquad \text{(H13)}
\end{aligned}
$$

We note that in the last line we evaluate the complex Fig.18a in a different way(by choosing opposite orientation for the internal link). The topological invariance of $\mathcal{V}_3$ path integral implies the two different ways must give out the same results. It is also not hard to see that the new sign factor $(-)^{m_1(g_4,g_i)+m_1(g_i,g_1)+m_1(g_i',g_2)+m_1(g_3,g_i')+m_1(g_5,g_i')+m_1(g_i',g_6)}$ is equivalent to $(-)^{m_1(g_4,g_i')+m_1(g_i',g_1)+m_1(g_i,g_2)+m_1(g_3,g_i)+m_1(g_5,g_i)+m_1(g_i,g_6)}$, since:

$$
\begin{aligned}
&m_1(g_4,g_i)+m_1(g_i,g_1)+m_1(g_i',g_2)+m_1(g_3,g_i')+m_1(g_5,g_i')+m_1(g_i',g_6)+ \\
&m_1(g_4,g_i')+m_1(g_i',g_1)+m_1(g_i,g_2)+m_1(g_3,g_i)+m_1(g_5,g_i)+m_1(g_i,g_6) \\
=&n_2(g_5,g_i',g_6)+n_2(g_5,g_i,g_6)+n_2(g_4,g_5,g_i')+n_2(g_4,g_5,g_i)+n_2(g_4,g_3,g_i')+n_2(g_4,g_3,g_i)+ \\
&n_2(g_3,g_i',g_2)+n_2(g_3,g_i,g_2)+n_2(g_i',g_2,g_1)+n_2(g_i,g_2,g_1)+n_2(g_i',g_6,g_1)+n_2(g_i,g_6,g_1) \\
=&0 \bmod 2. \qquad \text{(H14)}
\end{aligned}
$$

Thus, we have proved that the matrix element $H_{i;g_i,g_i'}(g_1g_2g_3g_4g_5g_6)$ is the same as $H^*_{i;g_i',g_i}(g_1g_2g_3g_4g_5g_6)$, implying H_i is a hermitian operator.

Finally, we note that the Hamiltonian only depends on ν_3 and n_2. It does not depend on m_1. So the Hamiltonian is symmetric under the G_f symmetry. By construction, the ideal ground state wave function is an eigenstate of H_i with eigenvalue 1. The topological invariance of $\mathcal{V}_3$ path integral implies that $H_i^2 = H_i$ and $H_iH_j = H_jH_i$. Thus H_i is a hermitian projection operator, and the set $\{H_i\}$ is a set of commuting projectors. Therefore $H = -\sum_i H_i$ is an exactly soluble Hamiltonian which realizes the fermionic SPT state described by $(\nu_3, n_2, u_2^g) \in \mathscr{H}^3[G_f, U_T(1)]$.

---

[1] L. D. Landau, Phys. Z. Sowjetunion **11**, 26 (1937)

[2] V. L. Ginzburg and L. D. Landau, Zh. Ekaper. Teoret. Fiz. **20**, 1064 (1950)

[3] L. D. Landau and E. M. Lifschitz, *Statistical Physics - Course of Theoretical Physics Vol 5* (Pergamon, London, 1958)

[4] X.-G. Wen, Phys. Rev. B **40**, 7387 (1989)

[5] X.-G. Wen and Q. Niu, Phys. Rev. B **41**, 9377 (1990)

[6] X.-G. Wen, Int. J. Mod. Phys. B **4**, 239 (1990)

[7] M. Levin and X.-G. Wen, Phys. Rev. Lett. **96**, 110405 (2006), cond-mat/0510613

[8] A. Kitaev and J. Preskill, Phys. Rev. Lett. **96**, 110404 (2006)

[9] X. Chen, Z.-C. Gu, and X.-G. Wen, Phys. Rev. B **82**, 155138 (2010), arXiv:1004.3835

[10] M. Levin and X.-G. Wen, Phys. Rev. B **71**, 045110 (2005), cond-mat/0404617

[11] F. Verstraete, J. I. Cirac, J. I. Latorre, E. Rico, and M. M. Wolf, Phys. Rev. Lett. **94**, 140601 (2005)

[12] G. Vidal, Phys. Rev. Lett. **99**, 220405 (2007)

[13] D. C. Tsui, H. L. Stormer, and A. C. Gossard, Phys. Rev. Lett. **48**, 1559 (1982)

[14] R. B. Laughlin, Phys. Rev. Lett. **50**, 1395 (1983)

[15] V. Kalmeyer and R. B. Laughlin, Phys. Rev. Lett. **59**, 2095

(1987)

[16] X.-G. Wen, F. Wilczek, and A. Zee, Phys. Rev. B **39**, 11413 (1989)

[17] N. Read and S. Sachdev, Phys. Rev. Lett. **66**, 1773 (1991)

[18] X.-G. Wen, Phys. Rev. B **44**, 2664 (1991)

[19] R. Moessner and S. L. Sondhi, Phys. Rev. Lett. **86**, 1881 (2001)

[20] G. Moore and N. Read, Nucl. Phys. B **360**, 362 (1991)

[21] X.-G. Wen, Phys. Rev. Lett. **66**, 802 (1991)

[22] R. Willett, J. P. Eisenstein, H. L. Strörmer, D. C. Tsui, A. C. Gossard, and J. H. English, Phys. Rev. Lett. **59**, 1776 (1987)

[23] I. P. Radu, J. B. Miller, C. M. Marcus, M. A. Kastner, L. N. Pfeiffer, and K. W. West, Science **320**, 899 (2008)

[24] M. Freedman, C. Nayak, K. Shtengel, K. Walker, and Z. Wang, Ann. Phys. (NY) **310**, 428 (2004), cond-mat/0307511

[25] Z.-C. Gu, Z. Wang, and X.-G. Wen(2010), arXiv:1010.1517

[26] Y.-M. Lu, X.-G. Wen, Z. Wang, and Z. Wang, Phys. Rev. B **81**, 115124 (2010), arXiv:0910.3988

[27] B. Blok and X.-G. Wen, Phys. Rev. B **42**, 8145 (1990)

[28] N. Read, Phys. Rev. Lett. **65**, 1502 (1990)

[29] J. Fröhlich and T. Kerler, Nucl. Phys. B **354**, 369 (1991)

[30] X.-G. Wen and A. Zee, Phys. Rev. B **46**, 2290 (1992)

[31] F. D. M. Haldane, Physics Letters A **93**, 464 (1983)

[32] I. Affleck, T. Kennedy, E. H. Lieb, and H. Tasaki, Commun. Math. Phys. **115**, 477 (1988)

[33] Z.-C. Gu and X.-G. Wen, Phys. Rev. B **80**, 155131 (2009), arXiv:0903.1069

[34] F. Pollmann, E. Berg, A. M. Turner, and M. Oshikawa(2009), arXiv:0909.4059

[35] C. L. Kane and E. J. Mele, Phys. Rev. Lett. **95**, 226801 (2005), cond-mat/0411737

[36] B. A. Bernevig and S.-C. Zhang, Phys. Rev. Lett. **96**, 106802 (2006)

[37] C. L. Kane and E. J. Mele, Phys. Rev. Lett. **95**, 146802 (2005), cond-mat/0506581

[38] J. E. Moore and L. Balents, Phys. Rev. B **75**, 121306 (2007), cond-mat/0607314

[39] L. Fu, C. L. Kane, and E. J. Mele, Phys. Rev. Lett. **98**, 106803 (2007), cond-mat/0607699

[40] X.-L. Qi, T. Hughes, and S.-C. Zhang, Phys. Rev. B **78**, 195424 (2008), arXiv:0802.3537

[41] H. Yao and S. A. Kivelson(2010), arXiv:1008.1065

[42] X. Chen, Z.-C. Gu, and X.-G. Wen, Phys. Rev. B **83**, 035107 (2011), arXiv:1008.3745

[43] N. Schuch, D. Perez-Garcia, and I. Cirac(2011), arXiv:1010.3732

[44] X.-G. Wen(2011), arXiv:1111.6341

[45] L. Fidkowski and A. Kitaev, Phys. Rev. B **81**, 134509 (2010), arXiv:0904.2197

[46] L. Fidkowski and A. Kitaev(2010), arXiv:1008.4138

[47] A. M. Turner, F. Pollmann, and E. Berg(2010), arXiv:1008.4346

[48] X. Chen, Z.-C. Gu, and X.-G. Wen(2011), arXiv:1103.3323

[49] X. Chen, Z.-X. Liu, and X.-G. Wen(2011), arXiv:1106.4752

[50] X. Chen, Z.-C. Gu, Z.-X. Liu, and X.-G. Wen(2011), arXiv:1106.4772

[51] For an introduction to group cohomology, see
"Group cohomology" on Wiki;
Romyar Sharifi, "AN INTRODUCTION TO GROUP COHOMOLOGY" on
http://math.arizona.edu/~sharifi/groupcoh.pdf

[52] X.-G. Wen, Phys. Rev. B **65**, 165113 (2002), cond-mat/0107071

[53] J. Wess and B. Zumino, Phys. Lett. B **37**, 95 (1971)

[54] E. Witten, Nucl. Phys. B **223**, 422 (1983)

[55] A. Kitaev, the Proceedings of the L.D.Landau Memorial Conference "Advances in Theoretical Physics"(2008), arXiv:0901.2686

[56] A. P. Schnyder, S. Ryu, A. Furusaki, and A. W. W. Ludwig, Phys. Rev. B **78**, 195125 (2008), arXiv:0803.2786

[57] A. Y. Kitaev, Phys.-Usp. **44**, 131 (2001), arXiv:cond-mat/0010440

[58] Z. C. Gu and M. Levin(2012), in preparation

[59] M. Levin and Z. C. Gu(2012), to appear

[60] A. Y. Kitaev, Annals of Phys. **321**, 2 (2006), arXiv:cond-mat/0506438

[61] M. Levin and C. P. Nave, Phys. Rev. Lett. **99**, 120601 (2007)

[62] Z.-C. Gu, F. Verstraete, and X.-G. Wen(2010), arXiv:1007.2022

[63] F. Costantino, Math. Z. **251**, 427 (2005), arXiv:math/0403014

[64] N. E. Steenrod, The Annals of Mathematics **48**, 290 (1947)

[65] N. E. Steenrod, Proceedings of the National Academy of Sciences of the United States of America **39**, 213 (1953)

[66] David Joyner, (2007) arXiv:math/0706.0549.

[67] Robert Greenblatt, Homology, Homotopy and Applications, **8**, 91, (2006).
http://www.intlpress.com/HHA/v8/n2/a5/pdf.

[68] Chris Gerig (mathoverflow.net/users/12310), Kuenneth-formula for group cohomology with nontrivial action on the coefficient, http://mathoverflow.net/questions/75485 (version: 2011-09-15)

[69] In a superconductor, the $U(1)$ symmetry is broken down to Z_2 symmetry, hence the vortex of a superconductor can be regarded as Z_2-flux.